

M. W. Kalinowski

**University of Economics
and Computer Science
Warsaw, Poland**

**Scalar Fields
in The Nonsymmetric
Kaluza–Klein (Jordan–Thiry)
Theory**

*To the memory
of my teacher
Professor Adam Bielecki*

Motto

W fizyce teoretycznej
potrzebujemy każdej matematyki,
a głównie takiej,
której jeszcze nie ma.

Adam Bielecki

(In theoretical physics
we need all the mathematics,
especially such a mathematics
which now does not exist.)

M. W. KALINOWSKI

Scalar fields in the nonsymmetric Kaluza–Klein (Jordan–Thiry) theory

Abstract

In this paper we construct the Nonsymmetric Jordan–Thiry Theory unifying N.G.T., the Yang–Mills’ field, the Higgs’ fields and scalar forces in a geometric manner. In this way we get masses from higher dimensions. We discuss spontaneous symmetry breaking, the Higgs’ mechanism and a mass generation in the theory. The scalar field Ψ (as in the classical Jordan–Thiry Theory) is connected to the effective gravitational constant. This field is massive and has Yukawa-type behaviour. We discuss the relation between $\mathbb{R}_+$ invariance and $\mathbf{U}(1)_F$ from G.U.T. within Einstein λ -transformation, and fermion number conservation. In this way we connect $\overline{W}_\mu$ -field from N.G.T. with a gauge field $\tilde{A}_\mu^F$ from G.U.T. We derive the equation of motion for a test particle from conservation laws in the hydrodynamic limit. We consider a truncation procedure for a tower of massive ρ_k (or Ψ_k) scalar fields using Friedrichs’ theory and an approximation procedure for the lagrangian involving Higgs’ field up to the second order with respect to $h_{\mu,\nu} = g_{\mu,\nu} - \eta_{\mu,\nu}$ and $\zeta k_{\tilde{a}\tilde{b}}^0$. The geodetic equations on the Jordan–Thiry manifold are considered with an emphasis to terms involving Higgs’ field. We consider also field equations in linear approximation and an influence of electromagnetic and $\overline{W}_\mu$ fields on field equations.

We consider a dynamics of Higgs’ field in the framework of cosmological models involving the scalar field Ψ . The field Ψ plays here a rôle of a quintessence field. We consider phase transition in cosmological models of the second and the first order. Due to a tower of massive scalar fields Ψ_k derived from the theory we are able to get a

warp factor known from some modern approaches. Using a warp factor we build a time-machine in the theory.

We consider a mass of a quintessence particle, various properties of a quintessence field. We calculate a speed of sound in a quintessence and fluctuations of a quintessence caused by primordial metric fluctuations.

Contents

Introduction	1
1. Elements of Geometry	7
2. The Nonsymmetric Tensor on a Lie Group	12
3. The Nonsymmetric Metrization of the Bundle $\underline{P}$	21
4. Preliminary Remarks	30
5. The Nonsymmetric Jordan–Thiry Theory over $V = E \times G/G_0$	43
6. Dimensional Reduction Procedure	52
7. Conformal Transformation of $g_{\mu\nu}$. Transformation of the Scalar Field ρ	63
8. Symmetry Breaking. Higgs Mechanism. Mass Generation. Cosmological Constant	66
9. Variational Principle. Equations of Fields. Interpretation of the Scalar Field Ψ	89
10. Properties of the Scalar Field Ψ . Cosmological Drifting of the Mass Scale	105
11. Fermion Number, $\mathbb{R}_+$ and $\mathbf{U}(1)_F$ Invariance	109
12. Fermion Number as the Second Gravitational Charge. $\overline{W}_\mu$ and $\tilde{A}_\mu^F$ Potentials	114
13. Equations of Motion for a Test Particle in the Nonsymmetric Jordan–Thiry Theory	123
14. On a Cosmological Origin of the Mass of the Scalar Field Ψ (or ρ). Cosmological Models with Quintessence and Inflation	127
15. Geodetic Equations on the Manifold $\underline{P}$	206
16. A Tower of Scalar Fields in the Nonsymmetric–Nonabelian Jordan–Thiry Theory. The Warp Factor	227
17. The Approximation Procedure for the Lagrangian of the Higgs Field in the Nonsymmetric Jordan–Thiry Theory	268
18. On some Cosmological Consequences of the Nonsymmetric Kaluza–Klein (Jordan–Thiry) Theory	286
19. Properties of a Quintessence	314
Conclusions	361
References	362
Acknowledgement	373

Introduction

In this work we develop a unification of the Nonsymmetric Gravitational Theory and gauge fields (Yang–Mills’ fields) including spontaneous symmetry breaking and the Higgs’ mechanism with scalar forces connected to the gravitational constant and cosmological terms appearing as the so-called quintessence. The theory is geometric and unifies tensor-scalar gravity with massive gauge theory using a multidimensional manifold in a Jordan–Thiry manner (see Refs. [1–70]). We use a nonsymmetric version of this theory (see [23], [24], [25], [26], [27]). The general scheme is the following. We introduce the principal fibre bundle over the base $V = E \times G/G_0$ with the structural group H , where E is a space-time, G is a compact semisimple Lie group, G_0 is its compact subgroup and H is a semisimple compact group. The manifold $M = G/G_0$ has an interpretation as a “vacuum states manifold” if G is broken to G_0 (classical vacuum states). We define on the space-time E , the nonsymmetric tensor $g_{\alpha\beta}$ from N.G.T., which is equivalent to the existence of two geometrical objects

$$\begin{aligned}\bar{g} &= g_{(\alpha\beta)} \bar{\theta}^\alpha \otimes \bar{\theta}^\beta \\ \underline{g} &= g_{[\alpha\beta]} \bar{\theta}^\alpha \wedge \bar{\theta}^\beta\end{aligned}$$

the symmetric tensor $\bar{g}$ and the 2-form $\underline{g}$. Simultaneously we introduce on E two connections from N.G.T. $\bar{W}_{\beta\gamma}^\alpha$ and $\tilde{\Gamma}_{\beta\gamma}^\alpha$. On the homogeneous space M we define the nonsymmetric metric tensor

$$g_{\tilde{a}\tilde{b}} = h_{\tilde{a}\tilde{b}}^0 + \zeta k_{\tilde{a}\tilde{b}}^0$$

where ζ is the dimensionless constant, in a geometric way. Thus we really have the nonsymmetric metric tensor on or $V = E \times G/G_0$.

$$\gamma_{AB} = \left(\begin{array}{c|c} g_{\alpha\beta} & 0 \\ \hline 0 & r^2 g_{\tilde{a}\tilde{b}} \end{array} \right)$$

r is a parameter which characterizes the size of the manifold $M = G/G_0$. Now on the principal bundle $\underline{P}$ we define the connection ω , which is the 1-form with values in the Lie algebra of H .

After this we introduce the nonsymmetric metric on $\underline{P}$ right-invariant with respect to the action of the group H , introducing scalar field ρ in a Jordan–Thiry manner (see Ref [18]). The only difference is that now our base space has more dimensions than four. It is $(n_1 + 4)$ -dimensional, where $n_1 = \dim(M) = \dim(G) - \dim(G_0)$. In other words, we combine the nonsymmetric tensor γ_{AB} on V with the right-invariant nonsymmetric tensor on the group H using the connection ω and the scalar field ρ .

We suppose that the factor ρ depends on a space-time point only. This condition can be abandoned and we consider a more general case where $\rho = \rho(x, y)$, $x \in E$, $y \in M$ resulting in a tower of massive scalar field $\rho_k, k = 1, 2, \dots$. This is really the Jordan–Thiry theory in the nonsymmetric version but with $(n_1 + 4)$ -dimensional “space-time”. After this we act in the classical manner as in Refs. [1], [18], [22], [23], [24], [26], [27]. We introduce the linear connection which is compatible with this nonsymmetric metric. This connection is the multidimensional analogue of the connection $\tilde{T}^\alpha_{\beta\gamma}$ on the space-time E . Simultaneously we introduce the second connection W . The connection W is the multidimensional analogue of the $\overline{W}$ -connection from N.G.T. and Einstein’s Unified Field Theory. It is the same as the connection from Refs. [23], [24]. Now we calculate the Moffat–Ricci curvature scalar $R(W)$ for the connection W and we get the following result. $R(W)$ is equal to the sum of the Moffat–Ricci curvature on the space-time E (the gravitational lagrangian in Moffat’s theory of gravitation), plus $(n_1 + 4)$ -dimensional lagrangian for the Yang–Mills’ field from the Nonsymmetric Kaluza–Klein Theory plus the Moffat–Ricci curvature scalar on the homogeneous space G/G_0 and the Moffat–Ricci curvature scalar on the group H plus the lagrangian for the scalar field ρ . The only difference is that our Yang–Mills’ field is defined on $(n_1 + 4)$ -dimensional “space-time” and the existence of the Moffat–Ricci curvature scalar of the connection on the homogeneous space G/G_0 . All of these terms (including $R(\overline{W})$) are multiplied by some factors depending on the scalar field ρ .

This lagrangian depends on the point of $V = E \times G/G_0$ i.e. on the point of the space-time E and on the point of $M = G/G_0$. The curvature scalar on G/G_0 also depends on the point of M .

We now go to the group structure of our theory. We assume G invariance of the connection ω on the principal fibre bundle $\underline{P}$, the so called Wang-condition (see Refs. [59], [71]–[76]). According to the Wang-theorem and the Ref. [72] the connection ω decomposes into the connection $\tilde{\omega}_E$ on the principal bundle Q over space-time E with structural group G and the multiplet of scalar fields Φ . Due to this decomposition the multidimensional Yang–Mills’ lagrangian decomposes into: a 4-dimensional Yang–Mills’ lagrangian with the gauge group G from the Nonsymmetric Kaluza–Klein Theory, plus a polynomial of 4th order with respect to the fields Φ , plus a term which is quadratic with respect to the gauge derivative of Φ (the gauge derivative with respect to the connection $\tilde{\omega}_E$ on a space-time E) plus a new term which is of 2nd order in the Φ , and is linear with respect to the Yang–Mills’ field strength. After this we perform the dimensional reduction procedure for the Moffat–Ricci scalar curvature on the manifold $\underline{P}$. We average $R(W)$ with respect to the homogeneous space $M = G/G_0$ as in Refs. [73], [74], [75]. In this way we get the lagrangian of our theory. It is the sum of the Moffat–Ricci curvature scalar on E (gravitational lagrangian) plus a Yang–Mills’ lagrangian with gauge group G from the Nonsymmetric Kaluza–Klein Theory (see [23]), plus a kinetic term for the scalar field Φ , plus a potential $V(\Phi)$ which is of 4th order with respect to Φ , plus $\mathcal{L}_{\text{int}}$ which describes a nonminimal interaction between the scalar field Φ and the Yang–Mills’ field, plus cosmological terms, plus lagrangian

for scalar field ρ . All of these terms (including $\overline{R}(\overline{W})$) are multiplied of course by some factors depending on the scalar field ρ . We redefine tensor $g_{\mu\nu}$ and ρ (as in [18], [19], [50]) and pass from scalar field ρ to Ψ

$$\rho = e^{-\Psi}$$

After this we get lagrangian which is the sum of gravitational lagrangian, Yang–Mills’ lagrangian, Higgs’ field lagrangian, interaction term $\mathcal{L}_{\text{int}}$ and lagrangian for scalar field Ψ plus cosmological terms. These terms depend now on the scalar field Ψ . In this way we have in our theory a multiplet of scalar fields (Ψ, Φ) . As in the Nonsymmetric–Nonabelian Kaluza–Klein Theory we get a polarization tensor of the Yang–Mills’ field induced by the skewsymmetric part of the metric on the space-time and on the group G . We get an additional term in the Yang–Mills’ lagrangian induced by the skewsymmetric part of the metric $g_{\alpha\beta}$ (see Ref. [23]). We get also $\mathcal{L}_{\text{int}}$, which is absent in the dimensional reduction procedure known up to now (see [73]–[76], [77]–[80]). Simultaneously, our potential for the scalar–Higgs’ field really differs from the analogous potential from Refs. [73]–[80]. Due to the skewsymmetric part of the metric on G/G_0 and on H it has a more complicated structure. This structure offers two kinds of critical points for the minimum of this potential: Φ_{crt}^0 and Φ_{crt}^1 . The first is known in the classical, symmetric dimensional reduction procedure (see [73], [80]) and corresponds to the trivial Higgs’ field (“pure gauge”). This is the “true” vacuum state of the theory. The second, Φ_{crt}^1 , corresponds to a more complex configuration. This is only a local (no absolute) minimum of V . It is a “false” vacuum. The Higgs’ field is not a “pure” gauge here. In the first case the unbroken group is always G_0 . In the second case, it is in general different and strongly depends on the details of the theory: groups G_0 , G , H , tensors ℓ_{ab} , $g_{\tilde{a}\tilde{b}}$ and the constants ζ , ξ . It results in a different spectrum of mass for intermediate bosons. However, the scale of the mass is the same and it is fixed by a constant r (“radius” of the manifold $M = G/G_0$). In the first case $V(\Phi_{\text{crt}}^0) = 0$, in the second case it is, in general, not zero $V(\Phi_{\text{crt}}^1) \neq 0$. Thus, in the first case, the cosmological constant is a sum of the scalar curvature on H and G/G_0 , and in the second case, we should add the value $V(\Phi_{\text{crt}}^1)$. We proved that using the constant ξ we are able in some cases to make the cosmological constant as small as we want (it is almost zero, from the observational data point of view). Here we can perform the same procedure for the second term in the cosmological constant using the constant ζ . In the first case we are able to make the cosmological constant sufficiently small but this is not possible in general for the second case.

The transition from “false” to “true” vacuum occurs as a second order phase transition (see [81]–[85]). We discuss this transition in context of the first order phase transition in models of the Universe (see [81]–[85]). In this paper the interesting point is that there exists an effective scale of masses, which depends on the scalar field Ψ .

Using Palatini variational principle we get an equation for fields in our theory. We find a gravitational equation from N.G.T. with Yang–Mills’, Higgs’ and scalar sources

(for scalar field Ψ) with cosmological terms. This gives us an interpretation of the scalar field Ψ as an effective gravitational constant

$$G_{\text{eff}} = G_N e^{-(n+2)\Psi}$$

We get an equation for this scalar field Ψ . Simultaneously we get equations for Yang–Mills’ and Higgs’ field. We also discuss the change of the effective scale of mass, m_{eff} with a relation to the change of the gravitational constant G_{eff} .

In the “true” vacuum case we get that the scalar field Ψ is massive and has Yukawa-type behaviour. In this way the weak equivalence principle is satisfied. In the “false” vacuum case the situation is more complex. It seems that there are possible some scalar forces with infinite range. Thus the two worlds constructed over the “true” vacuum and the “false” vacuum seem to be completely different: with different unbroken groups, different mass spectrum for the broken gauge and Higgs’ bosons, different cosmological constants and with different behaviour for the scalar field Ψ . The last point means that in the “false” vacuum case the weak equivalence principle could be violated and the gravitational constant (Newton’s constant) would increase in distance between bodies.

We discuss in the paper $\mathbb{R}_+$ and $\mathbf{U}(1)_F$ invariance in a continuation of our consideration from Refs. [26], [27]. We develop the first possibility from Section 8 of Ref. [26]. In this way we decide that $\mathbf{U}(1)_F$ invariance from G.U.T. is a local invariance. Due to a geometrical construction we are able to identify $\bar{W}_\mu$ from Moffat’s theory of gravitation with the four-potential $\tilde{A}_\mu^F$ corresponding to the $\mathbf{U}(1)_F$ group (internal rotations connected to fermion charge). In this way, the fermion number is conserved and plays the role of the second gravitational charge. Due to the Higgs’ mechanism $\tilde{A}_\mu^F$ is massive and its strength, $\tilde{H}_{\mu\nu}^F$, is of short range with Yukawa-type behaviour. This has important consequences. The Lorentz-like force term (or Coriolis-like force term) in the equation of motion for a test particle is of short range with Yukawa-type behaviour. The range of this force is smaller than the range of the weak interactions. Thus it is negligible in the equation of motion for a test particle. We discuss the possibility of the cosmological origin of the mass of the scalar field Ψ and geodetic equations on $\underline{P}$. We consider an infinite tower of scalar fields $\Psi_k(x)$ coming from the expansion of the field $\Psi(x, y)$ on the manifold $M = G/G_0$ into harmonics of the Beltrami–Laplace operator. Due to Friedrichs’ theory we can diagonalize an infinite matrix of masses for Ψ_k transforming them into new fields Ψ'_k . The truncation procedure means here to take a zero mass mode Ψ_0 and equal it to Ψ from the preceding section.

We discuss here cosmological models involving field Ψ which plays a rôle of a quintessence field. We find inflationary models of the Universe and we discuss a dynamics of the Higgs field. Higgs’ field dynamics undergoes a second order phase transition which causes a phase transition in an evolution of the Universe. This ends an inflationary epoch and changes an evolution of the field Ψ . Afterwards we consider the field Ψ as a quintessence field building some cosmological models with a quintessence and even with a K -essence. A dynamics of a Higgs field in several approximations gives us an

amount of an inflation. We consider also a fluctuation spectrum of a primordial fluctuations caused by a Higgs field and a speed up factor of an evolution. We speculate on a future of the Universe based on our simple model with a special behaviour of a quintessence.

We consider a possibility to take seriously additional dimensions in our theory in a framework similar to a Randall–Sundrum scenario. In our theory the additional dimensions connecting to the manifold M (a vacuum manifold) could be considered in such a way. They are not directly observable because the size of the manifold is very small. In order to see them it is necessary to excite massive modes of the scalar field (a tower of these fields). We find an interesting toy model (a 5-dimensional model) which can describe a possibility to travel with speed higher than the speed of light using the fifth dimension. This dimension has nothing to do with the fifth dimension in Kaluza–Klein theory. We do some analysis on an energy to excite a scalar field Ψ to get this special solution to the theory. We also discuss some quantitative relations involving traveling signals in the model via the fifth dimension. We consider the various possibilities to excite a warp factor due to fluctuations of a tower of scalar fields finding a density of an excitation energy. Eventually we find a zero energy (or almost zero) condition for such an excitation.

We also develop the approximation procedure for the lagrangian involving the Higgs' field up to the second order of expansion with respect to $h_{\mu\nu} = g_{\mu\nu} - \eta_{\mu\nu}$ and ζ . We find $V(\Phi)$, $\mathcal{L}(\overset{\text{gauge}}{\nabla}\Phi)$, $\mathcal{L}_{\text{int}}(\tilde{A}, \Phi)$ up to this order.

The paper is organized as follows. In section 1 we introduce the notation and the definitions of geometrical quantities which we use throughout this part. In sec. 2 we define nonsymmetric tensor on a group Lie. The third section deals with a nonsymmetric metrization of the fibre bundle. We define the skewsymmetric tensor on $M = G/G_0$ and examine its properties in the fourth section. Section 5 is devoted to the Nonsymmetric Jordan–Thiry Theory over the manifold $V = E \times G/G_0$. We introduce the connections ω^A_B and W^A_B which are analogues of the connections $\bar{W}^\alpha_\beta$ and $\bar{\omega}^\alpha_\beta$ from N.G.T. We calculate Moffat–Ricci curvature scalar for W^A_B and the density for this scalar. In section 6 we perform the dimensional reduction procedure for $R(W)$ using results from section 5. We find the integral of action in our theory and the lagrangian. In section 7 we perform a conformal transformation for the nonsymmetric tensor $g_{\mu\nu}$ and a transformation for the scalar field ρ . We pass from ρ to Ψ . In section 8 we discuss symmetry breaking in our theory with two kinds of critical points corresponding to two minimas, to the “true” and “false” vacuums. We deal also with the Higgs' mechanism, mass generation in the theory and the scale of mass for intermediate bosons. We also discuss a problem of cosmological terms and phase transition in the models of Universe based on our theory. We consider a possibility of nontrivial gauge configurations in our theory with topological magnetic charges (monopoles). Section 9 is devoted to the Palatini variational principle and field equations. We find the interpretation of the scalar field Ψ as a gravitational “constant”. In section 10 we discuss the properties of the scalar field Ψ . We prove that this field is massive. We discuss also cosmological

drifting of the effective scale of masses for the intermediate bosons. In section 11 we discuss $\mathbb{R}_+$ and $\mathbf{U}(1)_F$ invariance. In section 12 we discuss the geometrical construction which allows us to identify $\overline{W}_\mu$ with the four-potential $\tilde{A}_\mu^F$ connected to $\mathbf{U}(1)_F$ (fermion number subgroup of G). In section 13 we derive equation of motion for a test particle from conservation laws in the hydrodynamic limit. Section 14 is devoted to the problem of a cosmological origin of the mass of a scalar field Ψ (or ρ). We consider here cosmological models with a quintessence and a dynamics of Higgs' field with phase transitions. Section 15 deals with geodetic equations on $\underline{P}$ and geodetic equation deviation. In Sec. 16 we consider a tower of scalar fields coming from the factor ρ depending on a point of $M = G/G_0$. We consider here a toy model with a warp factor and we find a zero energy condition to excite the factor using a tower of scalar fields and (n_1+4) -dimensional equations for cosmological solutions. We consider a possibility to use a warp factor in order to send an effective superluminal signal and to create a time machine. Section 17 is devoted to the approximation procedure for the lagrangian involving Higgs' field and geodetic equations on the manifold $\underline{P}$. We consider also linear approximation of full field equations and an influence of $\overline{W}_\mu$ -field and electromagnetic field on field equations.

Section 18 is devoted to examination of cosmological consequences of our theory. We calculate spectral functions of primordial fluctuations and masses of quintessence particles in both de Sitter phases of an evolution of the Universe. In Section 19 we consider properties of a quintessence. We calculate a mass of a quintessence particle of order 10^{-5} eV, a speed of sound for a quintessence and various solutions of quintessence field equation. We consider also primordial fluctuations of a quintessence field.

1. Elements of Geometry

In this section we introduce notations and define geometric quantities used in the paper. We use a smooth principal fibre bundle $\underline{P}$, which includes in its definition the following list of differentiable manifolds and smooth maps:

a total (bundle) space $\underline{P}$

a base space E ; in our case it is a space time (or $E \times G/G_0$ as in sec. 4)

a projection $\pi : \underline{P} \rightarrow E$

a map $\Phi : \underline{P} \times G \rightarrow \underline{P}$ defining the action of G on $\underline{P}$; if $a, b \in G$ and $\varepsilon \in G$ is the unit element then $\Phi(a) \circ \Phi(b) = \Phi(ba)$ and $\Phi(\varepsilon) = \text{id}$ and $\Phi(a)p = \Phi(p, a) = R_ap = pa$, moreover $\pi \circ \Phi(a) = \pi$. ω is a 1-form of a connection on $\underline{P}$ with values in the Lie algebra of the group G .

Let $\Phi'(a)$ be the tangent map to $\Phi(a)$ whereas $\Phi^*(a)$ is contragradient to $\Phi(a)$ at the point a . The form ω is a form of Ad-type i.e.:

$$\Phi^*(a)\omega = \text{Ad}_{a^{-1}}\omega, \quad (1.1)$$

where $\text{Ad}_a \in \text{GL}(\mathfrak{g})$ is the tangent map to the internal automorphism of the group G (i.e. it is an adjoint representation of a group G)

$$\text{ad}_a(b) = aba^{-1}.$$

Due to the form ω we can introduce the distribution field of linear elements H_r , $r \in \underline{P}$, where $H_r \subset T_r(\underline{P})$ is a subspace of the space tangent to $\underline{P}$ at a point r and

$$v \in H_r \Leftrightarrow \omega(v) = 0. \quad (1.2)$$

We have

$$T_r(\underline{P}) = V_r \oplus H_r, \quad (1.3)$$

where H_r is called a subspace of horizontal vectors and V_r of vertical vectors. For vertical vectors $v \in V_r$ we have $\pi'(v) = 0$.

This means that v is tangent to fibres. Let us define

$$v = \text{hor}(v) + \text{ver}(v), \quad \text{hor}(v) \in H_r, \text{ ver}(v) \in V_r. \quad (1.4)$$

It is well known that the distribution H_r is equivalent to a choice of the connection ω . We can reproduce the connection form ω demanding that $\pi'|_{H_r} : H_r \rightarrow T_{\pi(r)}(E)$ is a vector space isomorphism ($\dim H_r = \dim E = 4$), $H_{\Phi(r,g)} = \Phi'(g)H_r$, $(T_{\pi(r)}(E))$ is a

tangent space to space-time E at a point $\pi(r)$). We use the operation “hor” for forms, i.e.:

$$(\text{hor } \beta)(X, Y) = \beta(\text{hor } X, \text{hor } Y), \quad (1.5)$$

where $X, Y \in T_r(\underline{P})$. The 2-form of curvature of the connection ω is:

$$\Omega = \text{hor } d\omega. \quad (1.6)$$

It is also a form of Ad-type like ω . Ω obeys the structural Cartan’s equation

$$\Omega = d\omega + \frac{1}{2}[\omega, \omega], \quad (1.7)$$

where

$$[\omega, \omega](X, Y) = [\omega(X), \omega(Y)].$$

Bianchi’s identity for ω is:

$$\text{hor } d\Omega = 0. \quad (1.8)$$

For the principal fibre bundle we use the following convenient scheme (Fig. 1A). The map $e : U \rightarrow \underline{P}$, $U \subset E$ (U open), so that $e \circ \pi = \text{id}_U$ is called a local section. From the physical point of view it means choosing the gauge. Thus

$$\begin{aligned} e^* \omega &= e^*(\omega^a X_a) = A_\mu^a \bar{\theta}^\mu X_a = A, \\ e^* \Omega &= e^*(\Omega^a X_a) = \frac{1}{2} F_{\mu\nu}^a \bar{\theta}^\mu \wedge \bar{\theta}^\nu X_a. \end{aligned} \quad (1.9)$$




Fig. 1A. The principal fibre bundle $\underline{P}$

Further we introduce a notation

$$\Omega^a = \frac{1}{2} H_{\mu\nu}^a \theta^\mu \wedge \theta^\nu, \quad (1.10)$$

where $\theta^\mu = \pi^*(\bar{\theta}^\mu)$ and

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + C_{bc}^a A_\mu^b A_\nu^c$$

X_a , ($a = 1, 2 \dots \dim G = n$) are generators of the Lie algebra $\mathfrak{g}$ of the group G and

$$[X_a, X_b] = C_{ab}^c X_c.$$

Analogically we can introduce a second local section $f : U \rightarrow P$, and corresponding to it $\bar{A} = f^* \omega$, $\bar{F} = f^* \Omega$. For every $x \in U \subset E$ there is an element $g(x) \in G$ such that $f(x) = e(x)g(x) = R_{g(x)}e(x) = \Phi(e(x), g(x))$. Due to Eq. (1.1) and an analogical formula for Ω one gets $\bar{A} = \text{Ad}_{g^{-1}} A + g^{-1} dg$ and $\bar{F} = \text{Ad}_{g^{-1}} F$. These formulae give a geometrical meaning of gauge transformation.

In this paper we use also a linear connection on manifolds $\underline{P}$ and E using the formalism of differential forms. So the basic quantity is a 1-form of a connection ω^A_B . This is an R -valued (coefficient) connection form and it is referred to the principal fibre bundle of frames with $\underline{P}$ or E as a base. The 2-form of curvature is the following

$$\Omega^A_B = d\omega^A_B + \omega^A_C \wedge \omega^C_B \quad (1.11)$$

and the 2-form of torsion

$$\Theta^A = D\theta^A, \quad (1.12)$$

where θ^A are basic forms, D means the exterior covariant derivative with respect to ω^A_B . The following relations define the interrelation between our symbols and generally used ones

$$\begin{aligned} \omega^A_B &= \Gamma^A_{BC} \theta^C, \\ \Theta^A &= \frac{1}{2} Q^A_{BC} \theta^B \wedge \theta^C, \\ \Omega^A_B &= \frac{1}{2} R^A_{BCD} \theta^C \wedge \theta^D. \end{aligned} \quad (1.13)$$

Where Γ^A_{BC} are coefficients of the connection (they do not have to be symmetric in indices B and C), R^A_{BCD} is a tensor of curvature and Q^A_{BC} is a tensor of torsion. Covariant exterior differentiation with respect to ω^A_B is given by the formula

$$\begin{aligned} D\Xi^A &= d\Xi^A + \omega^A_C \wedge \Xi^C, \\ D\Sigma^A_B &= d\Sigma^A_B + \omega^A_C \wedge \Sigma^C_B - \omega^C_B \wedge \Sigma^A_C. \end{aligned} \quad (1.14)$$

The forms of curvature Ω^A_B and torsion Θ^A obey Bianchi's identities

$$\begin{aligned} D\Omega^A_B &= 0, \\ D\Theta^A &= \Omega^A_B \wedge \theta^B. \end{aligned} \quad (1.15)$$

In the paper we use also Einstein's $+$ and $-$ differentiations for the nonsymmetric metric tensor g_{AB} .

$$Dg_{A+B-} = Dg_{AB} - g_{AD}Q^D_{BC}\theta^C, \quad (1.16)$$

where D is the covariant exterior derivative with respect to ω^A_B and Q^D_{BC} is the tensor of torsion for ω^A_B . In a homolonomic system of coordinates we easily get:

$$Dg_{A+B-} = g_{A+B-,C}\theta^C = [g_{AB,C} - g_{DB}\Gamma^D_{AC} - g_{AD}\Gamma^D_{CB}]\theta^C. \quad (1.17)$$

All quantities introduced in this section and their precise definitions can be found in Refs. [51], [59], [60], [61].

Finally let us connect a general formalism of the principal fibre bundle with a formalism of a linear connection on E or $\underline{P}$.

Let M be a m -dimensional pseudo-Riemannian manifold with metric g of arbitrary signature. Let $T(M)$ be the tangent bundle and $O(M, g)$ the principal fibre bundle of frames (orthonormal frames) over M . The structure group of $O(M, g)$ is the group $\text{Gl}(m, \mathbb{R})$ or the subgroup of $\text{Gl}(m, \mathbb{R})$, $O(m-p, p)$ which leaves the metric invariant. Let Π be the projection of $O(M, g)$ onto M . Let X be a tangent vector at a point x in $O(M, g)$. The canonical or soldering form $\tilde{\theta}$ is an R^m -valued form on $O(M, g)$ whose A th component $\tilde{\theta}^A$ at x of X is the A th component of $\Pi'(X)$ in the frame x . The connection form $\tilde{\omega} = \omega^A_B X^B_A$ is a 1-form on $O(M, g)$ which takes its values in the Lie algebra $\mathfrak{gl}(m, \mathbb{R})$ of $\text{Gl}(m, \mathbb{R})$ or in $\mathfrak{o}(m-p, p)$ of $O(m-p, p)$ and satisfies the structure equations

$$d\tilde{\omega} + \frac{1}{2}[\tilde{\omega}, \tilde{\omega}] = \tilde{\Omega} = \widetilde{\text{Hor}} d\tilde{\omega}, \quad (1.18)$$

where $\widetilde{\text{Hor}}$ is understood in the sense of $\tilde{\omega}$ and $\tilde{\Omega} = \tilde{\Omega}^A_B X^B_A$ is a $\mathfrak{gl}(m, \mathbb{R})(\mathfrak{o}(m-p, p))$ valued 2-form of the curvature. We can write Eq (1.18) using R^{2m} valued forms and commutation relations of the Lie algebra $\mathfrak{gl}(m, \mathbb{R})(\mathfrak{o}(m, m-p))$

$$\tilde{\Omega}^A_B = d\tilde{\omega}^A_B + \tilde{\omega}^A_C \wedge \tilde{\omega}^C_B. \quad (1.19)$$

Taking any local section of $O(M, g)$ e , one can get forms of coefficients of the connection, torsion, curvature, basic forms

$$\begin{aligned} e^* \tilde{\omega}^A_B &= \omega^A_B, \\ e^* \tilde{\Omega}^A_B &= \Omega^A_B, \\ e^* \tilde{\theta}^A &= \theta^A, \\ e^* \tilde{\Theta}^A &= \Theta^A. \end{aligned} \quad (1.20)$$

The forms of the right-hand side of the Eq. (1.20) are the forms defined in Eq. (1.11), (1.12), (1.13), (1.14) etc. We call this formalism a linear (affine, metric, Riemannian, Einstein) connection on M .




Fig. 1B. Principal fibre bundles $\underline{P}$, P' , P'' and P''' ,
 P''' is principal fibre bundle of frames over G .

In our theory it is necessary to consider at least four principal fibre bundles. A principal fibre bundle $\underline{P}$ over E with, a structural group G (a gauge group), connection ω and horizontality operator “hor”, a principal fibre bundle P' of frames over (E, g) with the connection $\tilde{\omega}^\alpha_\beta X^\beta_\alpha = \overline{\mathcal{W}}$, a structural group $\text{Gl}(4, \mathbb{R})(O(1, 3))$ an operator of horizontality “hor”, a principal fibre bundle P'' of frames over $(\underline{P}, \gamma)$ (a metrized fibre bundle $\underline{P}$) with a structural group $\text{Gl}(4+n, \mathbb{R})(O(n+3, 1))$, a connection $\tilde{\omega}^A_B X^B_A = \tilde{\omega}$ and an operator of horizontality “hor” and a principal fibre bundle of frames P''' over G with a projection Π''' , operator of horizontality (hor)''', a connection ω and the structural group $\text{Gl}(n, \mathbb{R})$. Moreover in order to simplify considerations we are using in the work the formalism of linear connection coefficients on manifolds (E, g) , $(\underline{P}, \gamma)$ and a principal fibre bundle formalism for $\underline{P}$ i.e. (a principal fibre bundle over E with the structural group G a gauge group). In the meaning of the author of this work this is a way to make the formalism more natural and readable (see Fig. 1B).

2. The Nonsymmetric Tensor on a Lie Group

Let G be a Lie group and let us define on G a tensor field $h = h_{ab}v^a \otimes v^b$ and a field of a 2-form $k = k_{ab}v^a \wedge v^b$, where

$$dv^a = -\frac{1}{2}C^a_{bc}v^b \wedge v^c, \quad (2.1)$$

v^a is a usual left-invariant frame on G , C^a_{bc} are structure constants. Let X_a be generators of a Lie algebra $G - \mathfrak{g}$, X_a are left-invariant vector fields on G and they are dual to the forms v^a .

$$[X_a, X_b] = C^c_{ab}X_c. \quad (2.2)$$

Using h and k we construct a tensor field on G

$$\ell_{ab} = h_{ab} + \mu k_{ab}, \quad (2.3)$$

where μ is a real number. Let us remind that the left-invariant vector fields on G are infinitesimal transformations of a right action of G on G . The symbol $\text{Ad}_G(g)$ means a matrix of the adjoint representation of the group G . Shortly we denote it $\text{Ad } g$. R -means a right action of the group G on G , L , a left action ($R(g), L(g), g \in G$).

We are looking for the following h and k

$$R^*(g)h = h, \quad (2.4)$$

$$R^*(g)k = k, \quad (2.5)$$

or in terms of the tensor ℓ_{ab}

$$R^*(g)\ell = \ell. \quad (2.6)$$

The condition (2.5) can be rewritten

$$(R^*(g))k_{g_1}(X_{g_1}, Y_{g_1}) = k_{g_1g}(X_{g_1}g', Y_{g_1}g') = k_{g_1}(X_{g_1}, Y_{g_1}), \quad (2.5a)$$

where $g, g_1 \in G$.

Moreover X, Y are left-invariant vector fields on G . Thus $X_g = X_\varepsilon = X$, $Y_g = Y_\varepsilon = Y$ and

$$(R^*(g))k_{g_1}(X, Y) = k_{g_1g}(Xg', Yg') = k_{g_1}(X, Y), \quad (2.5b)$$

where $\varepsilon \in G$, is a unit element of G .

In order to find h and k satisfying (2.4) and (2.5) we define a linear connection on G such that

$$\widehat{\omega}^a_b = -C^a_{bc}v^c. \quad (2.7)$$

Let the covariant differentiation with respect to $\hat{\omega}^a_b$ be $\hat{\nabla}_c$ and an exterior covariant differentiation $\hat{D}$. It is easy to see that this connection is flat

$$\hat{\Omega}^a_b = d\hat{\omega}^a_b + \hat{\omega}^a_c \wedge \hat{\omega}^c_b = 0 \quad (2.8)$$

with non-zero torsion

$$\hat{\Theta}^a = \hat{D}v^a = dv^a + \hat{\omega}^a_b \wedge v^b = \frac{1}{2}C^a_{bc}v^b \wedge v^c \quad (2.8a)$$

and with a tensor of torsion

$$\hat{Q}^a_{bc} = C^a_{bc}. \quad (2.9)$$

This connection is also metric. It means that the Killing–Cartan tensor on the group G is absolutely parallel with respect to $\hat{\omega}^a_b$. A parallel transport according to this connection is a right action of the group G on G .

One can easily find that (2.4), (2.5), (2.6) are equivalent to the condition

$$\hat{\nabla}_c \ell_{ab} = 0. \quad (2.10)$$

Thus in order to find h and k we should solve the Eqs. (2.10) on the group G . The tensor ℓ_{ab} satisfies an equation equivalent to (2.10)

$$X_f \ell_{cd} + \ell_{nd} C^n_{cf} + \ell_{cn} C^n_{df} = 0. \quad (2.11)$$

It is easy to see that a bi-invariant tensor h on G satisfies (2.11) identically (for example a Killing–Cartan tensor).

Thus one gets for a tensor k_{ab}

$$\hat{\nabla}_c k_{ab} = X_c k_{ab} + k_{nb} C^n_{ac} + k_{an} C^n_{bc} = 0. \quad (2.12)$$

It is easy to see that if k_{ab} satisfies (2.12) $b \cdot k_{ab}$ satisfies this condition as well for $b = \text{const}$.

In the case of an abelian group k is bi-invariant on G .

The interesting case in our theory is a semisimple group G . In this case k_{ab} can not be bi-invariant. The only bi-invariant 2-form on the semisimple Lie group G is a zero form. Moreover the Eq. (2.12) has always a solution on a semisimple group and k is right-invariant. Moreover we suppose that the symmetric part of ℓ is bi-invariant (left and right-invariant) and k only right-invariant.

We can also define k in a special way

$$k(A, B) = h([A, B], V), \quad A = A^a X_a, \quad B = B^a X_a, \quad (2.13)$$

where

$$\hat{\nabla}_c V_d = 0, \quad (2.14)$$

$V = V_d \otimes v^d$ is a covector field on G (it is right-invariant) and h is a Killing–Cartan tensor on G .

In order to be more familiar with the notion of a tensor k we find it for the group $\text{SO}(3)$. In this case we have left-invariant vector fields

$$\begin{aligned} e_1 &= \cos \psi \frac{\partial}{\partial \theta} - \sin \psi \left(\cos \theta \frac{\partial}{\partial \phi} - \frac{1}{\sin \theta} \frac{\partial}{\partial \phi} \right), \\ e_2 &= \sin \psi \frac{\partial}{\partial \theta} + \cos \psi \left(\cot \theta \frac{\partial}{\partial \psi} - \frac{1}{\sin \theta} \frac{\partial}{\partial \phi} \right), \\ e_3 &= \frac{\partial}{\partial \psi}, \end{aligned} \tag{2.15}$$

such that

$$[e_a, e_b] = -\varepsilon_{abc} e_c; \quad a, b, c = 1, 2, 3, \tag{2.16}$$

θ, ϕ, ψ are Euler angles—usual parametrization of $\text{SO}(3)$

$$\begin{aligned} 0 &\leq \theta \leq \pi, \\ 0 &\leq \psi \leq 2\pi, \\ 0 &\leq \phi \leq 2\pi \end{aligned} \tag{2.17}$$

and $\varepsilon_{123} = 1$ and ε_{abc} is a Levi–Civita symbol (see [62])

In this case one can easily integrate (2.14) and finds

$$\begin{aligned} V_1(\theta, \phi, \psi) &= a(\cos \phi \cos \psi - \cos \theta \sin \phi \sin \psi) + b \sin \psi \sin \theta \\ &\quad - c(\sin \phi \cos \psi + \cos \theta \cos \phi \sin \psi), \\ V_2(\theta, \phi, \psi) &= a(\sin \psi \cos \phi + \cos \theta \cos \psi \sin \phi) - b \cos \psi \sin \theta \\ &\quad + c(\cos \theta \cos \phi \cos \psi - \sin \phi \sin \psi), \\ V_3(\theta, \phi, \psi) &= a \sin \phi \sin \theta + b \cos \theta + c \sin \phi \sin \theta, \quad a, b, c = \text{const}. \end{aligned} \tag{2.18}$$

In a simpler case $a = c = 0$, $b \neq 0$, one gets:

$$\begin{aligned} V_1 &= b \sin \theta \sin \psi, \\ V_2 &= -b \sin \theta \cos \psi, \\ V_3 &= b \cos \theta, \quad b = \text{const}. \end{aligned} \tag{2.19}$$

For

$$k_{ab} = \varepsilon_{abc} V_c \tag{2.20}$$

we get

$$k_{ab} = \begin{pmatrix} 0 & (a \sin \psi \sin \theta + b \cos \theta + c \sin \phi \sin \theta) & -[a(\sin \psi \cos \phi + \cos \theta \cos \psi \sin \phi) + b \cos \psi \sin \theta + c(\cos \theta \cos \phi \cos \psi - \sin \phi \sin \psi)] \\ -(a \sin \phi \sin \theta + b \cos \theta + c \sin \phi \sin \theta) & 0 & [a(\alpha \phi \cos \psi - \cos \theta \sin \phi \sin \psi) + b \sin \psi \sin \theta + c(\sin \phi \cos \psi + \cos \theta \cos \phi \sin \psi)] \\ a(\sin \psi \cos \phi + \cos \theta \cos \psi \sin \phi) + b \cos \psi \sin \theta + c(\cos \theta \cos \phi \cos \psi - \sin \phi \sin \psi) & -[a(\cos \phi \cos \psi - \cos \theta \sin \phi \sin \psi) + b \sin \psi \sin \theta + c(\sin \phi \cos \psi + \cos \theta \cos \phi \sin \psi)] & 0 \end{pmatrix}, \quad (2.21)$$

In a simpler case for $a = c = 0$, $b \neq 0$ one gets:

$$k_{ab} = \begin{pmatrix} 0 & , & b \cos \theta & , & -b \sin \theta \cos \psi \\ -b \cos \theta & , & 0 & , & b \sin \theta \cos \psi \\ b \sin \theta \cos \psi & , & -b \sin \theta \sin \psi & , & 0 \end{pmatrix} \quad (2.22)$$

Thus if we choose for h a Killing–Cartan tensor on $\text{SO}(3)$ (this is a unique bi-invariant tensor on $\text{SO}(3)$ *modulo* constant factor)

$$h_{ab} = -2\delta_{ab} \quad (2.23)$$

we easily get:

$$\ell_{ab} = \begin{pmatrix} -2 & \mu(a \sin \psi \sin \theta + \beta \cos \theta + \gamma \sin \phi \sin \theta) & -\mu[\alpha(\sin \psi \cos \phi + \cos \theta \cos \psi \sin \phi) + \beta \cos \psi \sin \theta + \gamma(\cos \theta \cos \phi \cos \psi - \sin \phi \sin \psi)] \\ -2 & \mu[\alpha(\cos \phi \cos \psi - \cos \theta \sin \phi \sin \psi) + \beta \sin \psi \sin \theta + \gamma(\sin \phi \cos \psi + \cos \theta \cos \phi \sin \psi)] & -2 \\ \mu(\sin \psi \cos \phi + \cos \theta \cos \psi \sin \phi) + \beta \cos \psi \sin \theta + \gamma(\cos \theta \cos \phi \cos \psi - \sin \phi \sin \psi) & -\mu[\alpha(\cos \phi \cos \psi - \cos \theta \sin \phi \sin \psi) + \beta \sin \psi \sin \theta + \gamma(\sin \phi \cos \psi + \cos \theta \cos \phi \sin \psi)] & -2 \end{pmatrix} \quad (2.24)$$

where $\mu = \eta\sqrt{a^2 + b^2 + c^2}$, $\eta^2 = 1$, $\alpha = \frac{a}{\mu}$, $\beta = \frac{b}{\mu}$, $\gamma = \frac{c}{\mu}$.

In a simpler case i.e. for $a = c = 0$, $b \neq 0$ one gets (absorbing β by μ):

$$\ell_{ab} = \begin{pmatrix} -2 & , & \mu \cos \theta & , & -\mu \sin \theta \cos \psi \\ -\mu \cos \theta & , & -2 & , & \mu \sin \theta \cos \psi \\ \mu \sin \theta \cos \psi & , & -\mu \sin \theta \sin \psi & , & -2 \end{pmatrix} \quad (2.24a)$$

For an inverse tensor ℓ^{ab} such that

$$\ell^{ab} \ell_{ac} = \ell^{ba} \ell_{ca} = \delta_c^b \quad (2.25)$$

we have

$$\ell^{ab} = \frac{\Delta^{ab}}{\Delta}. \quad (2.26)$$

where $\Delta = \det(\ell_{ab}) = -2(4 + \mu^2)$ Δ^{ab} is a factor matrix and

$$\begin{aligned} \Delta^{11} &= 4 + \mu^2 \sin^2 \theta \sin^2 \psi, \\ \Delta^{12} &= -(2\mu \cos \theta + \mu^2 \sin^2 \theta \sin \psi \cos \psi), \\ \Delta^{13} &= (\mu^2 \cos \theta \sin \theta \sin \psi - 2\mu \sin \theta \cos \psi), \\ \Delta^{21} &= (2\mu \cos \theta - \mu^2 \sin^2 \theta \sin \psi \cos \psi), \\ \Delta^{22} &= (4 + \mu^2 \sin^2 \theta \cos^2 \psi), \\ \Delta^{23} &= -(2\mu \sin \theta \sin \psi + \mu^2 \cos \theta \sin \theta \cos \psi), \\ \Delta^{31} &= (\mu^2 \cos \theta \sin \theta \sin \psi + 2\mu \sin \theta \cos \psi), \\ \Delta^{32} &= (2\mu \sin \theta \sin \psi - \mu^2 \cos \theta \sin \theta \cos \psi), \\ \Delta^{33} &= (4 + \mu^2 \cos^2 \theta), \end{aligned} \quad (2.27)$$

In the case of $\text{SO}(3)$ Eq. (2.21) is the most general tensor satisfying (2.5) except a constant factor in front. Thus this tensor is unique for $\text{SO}(3)$ *modulo* a constant factor.

In the case of any $\text{SO}(n)$ one can find k and ℓ similarly using Euler angles parametrization and so for classical groups $\text{SU}(n)$, $\text{Sp}(2n)$, G_2 , F_4 , E_6 , E_7 , E_8 . In the case of solvable and nilpotent groups we can also try to find a bi-invariant skew-symmetric tensors.

Finally we suggest a general form of the tensor k_{ab} on a semi-simple group G i.e. such that Eq. (2.4) is satisfied. The solution of the Eq. (2.10) and (2.12) are as follows

$$\ell_{ab}(e^c) = \ell_{a'b'}(\varepsilon)(e^{\text{Ad}' c})_a^{a'}(e^{\text{Ad}' c})_b^{b'}$$

and

$$k_{ab}(e^c) = k_{a'b'}(\varepsilon)(e^{\text{Ad}' c})_a^{a'}(e^{\text{Ad}' c})_b^{b'}.$$

One writes

$$k_{ab}(g) = f_{a'b'} U^{a'}_a(g) U^{b'}_b(g), \quad g \in G, \quad (2.28)$$

where $U(g) = \text{Ad}_G(g)$ is an adjoint representation of the group G . It is easy to see that for (2.28) we have

$$\widehat{\nabla}_c k_{ab} = 0, \quad (2.29)$$

$$f_{ab} = -f_{ba} = \text{const} \quad (2.30)$$

and it is defined in the representation space of the adjoint representation of the group G . In the case of the group $\text{SO}(3)$ one has

$$f_{ab} = \varepsilon_{abc} f_c, \quad (2.31)$$

$$k_{ab} = \varepsilon_{abc} V_c \quad (2.31a)$$

and

$$V_a = f_{c'} U^{c'}{}_a(g). \quad (2.32)$$

If we choose $f_c = (0, 0, b)$ we get Eq. (2.19). Moreover it is always possible because an orthogonal ($\text{SO}(3)$) transformation can transform any vector f into $(0, 0, \pm\|f\|)$, where $\|f\|$ is the length of f .

The semisimple Lie group G can be considered a Riemannian manifold equipped with a bi-invariant tensor h (a Killing–Cartan tensor) and a connection induced by this tensor. This Riemannian manifold has a constant curvature. Such a manifold has a maximal group of isometries H of a dimension $\frac{1}{2}n(n+1)$, $n = \dim G$ (see Ref [59]). (The isometry is here understood in a sense of the metric measured along geodetic lines in a Riemannian geometry induced by a Killing–Cartan tensor.) This group is a Lie group. It is easy to see that for $G = \text{SO}(3)$ we have $H = \text{SO}(3) \rtimes \text{SO}(3)$ and $\dim \text{SO}(3) \rtimes \text{SO}(3) = 6$, $\dim \text{SO}(3) = 3$. The group $\text{SO}(3)$ leaves the Killing–Cartan tensor h_{ab} invariant

$$h_{a'b'} A^{a'}{}_a A^{b'}{}_b = h_{ab}, \quad (2.33)$$

where $A \in \text{SO}(3)$.

Moreover f_{ab} has exactly 3 arbitrary parameters and solutions of Eq. (2.12) have the same freedom in arbitrary constants. This suggests that the tensor (2.28) could be in some sense unique *modulo* an isometry on $\text{SO}(3)$ and a constant factor b . In this case the classification of k_{ab} tensors on a $\text{SO}(3)$ could be reduced to the classification of skewsymmetric tensors f_{ab} with respect to the action of the group $\text{SO}(3)$. In general the situation is more complex, because $\text{SO}(n)$, $n = \dim G$ does not leave the commutator (Lie bracket) invariant.

Let us suppose that G is compact. In this case we should find all inequivalent f_{ab} -tensors with respect to an orthogonal transformation $A \in \text{SO}(n)$. It means we should transform f_{ab} to a canonical form via an orthogonal matrix i.e.

$$(f_{ab}) = f \rightarrow f' = (f'_{ab}) = A^T f A = A^{-1} f A. \quad (2.34)$$

For skewsymmetric matrices we have the following canonical forms, the so-called block-diagonal matrices For $n = 2m$

$$f = \begin{bmatrix} 0 & \xi^1 & & & \mathbf{0} \\ -\xi^1 & 0 & & & \\ & & \ddots & & \\ \mathbf{0} & & & 0 & \xi^m \\ & & & -\xi^m & 0 \end{bmatrix} \quad (2.35)$$

or for $n = 2m + 1$

$$f = \begin{bmatrix} 0 & \xi^1 & & & \mathbf{0} \\ -\xi^1 & 0 & & & \\ & & \ddots & & \\ & \mathbf{0} & & 0 & \xi^m & 0 \\ & & & -\xi^m & 0 & 0 \\ & & & 0 & 0 & 0 \end{bmatrix}, \quad (2.36)$$

where $\xi^1, \xi^2 \dots \xi^m$ are real numbers. In order to find them we should solve a secular equation for f

$$\det(\mu I_n - f) = \mu^n + a_1(f)\mu^{n-2} + a_2(f)\mu^{n-2} + \dots \quad (2.37)$$

$$(I_n = (\delta_j^i)_{i,j=1,2\dots n}).$$

The coefficients $a_1, a_2 \dots$ are invariant with respect to an action of the group $O(n)$ ($SO(n)$) and they are functions of $\xi^1, \dots \xi^m$. Thus in the case of a compact semisimple Lie group the skewsymmetric tensor k_{ab} on G is defined as follows

$$k_{ab}(g) = b \cdot \tilde{f}_{a'b'} U^{a'}_a(g) U^{b'}_b(g), \quad (2.38)$$

where b is a constant real factor and $(\tilde{f}_{ab}) = \tilde{f}$ is given by

$$\tilde{f} = A^T \begin{bmatrix} 0 & 1 & & & \mathbf{0} \\ -1 & 0 & & & \\ & & 0 & \xi^1 & \\ & & -\xi^1 & 0 & \\ & & & \ddots & \\ \mathbf{0} & & & & 0 & \xi^{m-1} \\ & & & & -\xi^{m-1} & 0 \end{bmatrix} A, \quad (2.39)$$

for $n = 2m$, or

$$\tilde{f} = A^T \begin{bmatrix} 0 & 1 & & & \mathbf{0} \\ -1 & 0 & & & \\ & & 0 & \xi^1 & \\ & & -\xi^1 & 0 & \\ & & & \ddots & \\ \mathbf{0} & & & & 0 & \xi^{m-1} \\ & & & & -\xi^{m-1} & 0 \end{bmatrix} A, \quad (2.39a)$$

for $n = 2m + 1$.

Thus one gets

$$k_{ab}(g) = b \cdot C^c_{ab} f^0_{c'} U^{c'}_c(g), \quad (2.43)$$

where b is a constant factor and

$$(f^0_{c'}) = f^0 = \underbrace{(0, 0, \dots, 1)}_{n \text{ times}}. \quad (2.44)$$

Thus we can write k in a more compact form

$$k(A, B)(g) = b \cdot h([A, B], \text{Ad}_g f^0), \quad (2.45)$$

where $A = A^a X_a$, $B = B^a X_a$

Using a bi-invariancy of the Killing–Cartan tensor one can write

$$k(A, B)(g) = b \cdot h(\text{Ad}_{g^{-1}}[A, B], f^0). \quad (2.45a)$$

Moreover if there is $\tilde{g} \in G$ such that $\tilde{g}^2 = g$ we get

$$k(A, B)(g^2) = b \cdot h(\text{Ad}_{\tilde{g}^{-1}}[A, B], \text{Ad}_{\tilde{g}} f^0). \quad (2.46)$$

We find an interpretation of a factor b for k given by formulae (2.45–46).

One gets

$$k_{ab} k^{ab} = h^{aa'} h^{bb'} k_{ab} k_{a'b'} = b^2 \|\text{Ad}_g f^0\|^2 = b^2. \quad (2.47)$$

Thus we have

$$b = \pm \sqrt{k_{ab} k^{ab}}. \quad (2.48)$$

Finally let us notice that we can repeat the considerations changing right(left)-invariant to left(right)-invariant in all the places. In this case we can consider left-invariant 2-form k and left-invariant nonsymmetric tensor on a Lie group G .

In general we can simply say that our right invariant tensor on a group G can be constructed as follows. We define a skew-symmetric tensor at a unit element of a group and afterwards propagate it on G via group translations (if the group is simply connected). If it is not simply connected, we define it for a universal covering group or only for simply connected part containing a unit element.

3. The Nonsymmetric Metrization of the Bundle $\underline{P}$

Let us introduce the principal fibre bundle $\underline{P}$ over the space-time E with the structural group G and with the projection π . Let us suppose that (E, g) is a manifold with a nonsymmetric metric tensor of the signature $(-, -, -, +)$

$$g_{\mu\nu} = g_{(\mu\nu)} + g_{[\mu\nu]} \quad (3.1)$$

Let us introduce a natural frame on $\underline{P}$

$$\theta^A = (\pi^*(\bar{\theta}^\alpha), \theta^a = \lambda\omega^a), \quad \lambda = \text{const}. \quad (3.2)$$

It is convenient to introduce the following notations. Capital Latin indices A, B, C run $1, 2, 3, \dots, n+4$, $n = \dim G$. Lower Greek indices $\alpha, \beta, \gamma, \delta = 1, 2, 3, 4$ and lower Latin cases $a, b, c, d = 5, 6, \dots, n+4$. The symbol “ $-$ ” over θ^α and over other quantities indicates that these quantities are defined on E .

It is easy to see that the existence of the nonsymmetric metric on E is equivalent to introducing two independent geometrical quantities on E .

$$\overline{g} = g_{\alpha\beta} \bar{\theta}^\alpha \otimes \bar{\theta}^\beta = g_{(\alpha\beta)} \bar{\theta}^\alpha \otimes \bar{\theta}^\beta, \quad (3.3)$$

$$\underline{g} = g_{\alpha\beta} \bar{\theta}^\alpha \wedge \bar{\theta}^\beta = g_{[\alpha\beta]} \bar{\theta}^\alpha \wedge \bar{\theta}^\beta, \quad (3.4)$$

i.e. the symmetric metric tensor $\overline{g}$ on E and 2-form $\underline{g}$. On the group G we can introduce a bi-invariant symmetric tensor called the Killing–Cartan tensor.

$$h(A, B) = \text{Tr}(\text{Ad}'_A \circ \text{Ad}'_B), \quad (3.5)$$

where $\text{Ad}_A(C) = [A, C]$ (it is tangent to Ad i.e. it is an “infinitesimal” Ad -transformation). It is easy to see that

$$h(A, B) = h_{ab} A^a \cdot B^b, \quad (3.6)$$

where

$$h_{ab} = C^c_{ad} C^d_{bc}, \quad h_{ab} = h_{ba}, \quad A = A^a X_a, \quad B = B^a X_a.$$

This tensor is distinguished by the group structure, but there are of course other bi-invariant tensors on G . Normally it is supposed that G is semisimple. It means that $\det(h_{ab}) \neq 0$. In this construction we use $\ell_{(ab)} = h_{ab}$ (the bi-invariant tensor on G) in order to get a proper limit (i.e. the Nonabelian Kaluza–Klein Theory) for $\mu = 0$.

For a natural 2-form k on G , or a natural skewsymmetric right-invariant tensor we choose k described in sec. 2, k is zero for $\mathbf{U}(1)$. Let us turn to the nonsymmetric

natural metrization of $\underline{P}$. Let us suppose that:

$$\overline{\gamma}(X, Y) = \overline{g}(\pi'X, \pi'Y) + \lambda^2 \rho^2 h(\omega(X), \omega(Y)), \quad (3.7)$$

$$\underline{\gamma}(X, Y) = \underline{g}(\pi'X, \pi'Y) + \mu \lambda^2 \rho^2 k(\omega(X), \omega(Y)), \quad (3.8)$$

$\mu = \text{const}$ and is dimensionless, $X, Y \in \text{Tan}(\underline{P})$ and $\rho = \rho(x)$ is a scalar field defined on E . The first formula (2.9) was introduced by A. Trautman (in the case with $\rho = 1$) for the symmetric natural metrization of $\underline{P}$ and it was used to construct the Kaluza–Klein Theory for $\mathbf{U}(1)$ and nonabelian generalizations of this theory. It is easy to see that

$$\overline{\gamma} = \pi^* \overline{g} \oplus \rho^2 h_{ab} \theta^a \otimes \theta^b, \quad (3.9)$$

$$\underline{\gamma} = \pi^* \underline{g} + \mu \rho^2 k_{ab} \theta^a \wedge \theta^b, \quad (3.10)$$

or

$$\gamma_{(AB)} = \left(\begin{array}{c|c} g_{(\alpha\beta)} & 0 \\ \hline 0 & \rho^2 h_{ab} \end{array} \right), \quad (3.11)$$

$$\gamma_{[AB]} = \left(\begin{array}{c|c} g_{[\alpha\beta]} & 0 \\ \hline 0 & \mu \rho^2 k_{ab} \end{array} \right). \quad (3.12)$$

For

$$\gamma_{AB} = \gamma_{(AB)} + \gamma_{[AB]}$$

one easily gets

$$\gamma_{AB} = \left(\begin{array}{c|c} g_{\alpha\beta} & 0 \\ \hline 0 & \rho^2 \ell_{ab} \end{array} \right), \quad (3.13)$$

where $\ell_{ab} = h_{ab} + \mu k_{ab}$. Tensor γ_{AB} has this simple form in the natural frame on $\underline{P}, \theta^A$. This frame is unholonomical, because:

$$d\theta^a = \frac{\lambda}{2} \left(H^a_{\mu\nu} \theta^\mu \wedge \theta^\nu - \frac{1}{\lambda^2} C^a_{bc} \theta^b \wedge \theta^c \right) \neq 0 \quad (3.14)$$

γ is invariant with respect to the right action of the group on $\underline{P}$. In the case with $k_{ab} = 0$ we have

$$\gamma_{AB} = \left(\begin{array}{c|c} g_{\alpha\beta} & 0 \\ \hline 0 & \rho^2 h_{ab} \end{array} \right). \quad (3.15)$$

For the electromagnetic case ($G = \mathbf{U}(1)$) one easily finds

$$\gamma_{AB} = \left(\begin{array}{c|c} g_{\alpha\beta} & 0 \\ \hline 0 & -\rho^2 \end{array} \right). \quad (3.16)$$

Now let us take a section $e : E \rightarrow \underline{P}$ and attach to it a frame v^a , $a = 5, 6 \dots n+4$, selecting $X^\mu = \text{const}$ on a fibre in such a way that e is given by the condition $e^* v^a = 0$ and the fundamental fields ζ_a such that $v^a(\zeta_b) = \delta^a_b$ satisfy $[\zeta_b, \zeta_a] = \frac{1}{\lambda} C^c_{ab} \zeta_c$.

Thus we have

$$\omega = \frac{1}{\lambda} v^a X_a + \pi^*(A^a{}_\mu \bar{\theta}^\mu) X_a,$$

where

$$e^* \omega = A = A^a{}_\mu \bar{\theta}^\mu X_a.$$

In this frame the tensor γ takes the form

$$\gamma_{AB} = \left(\begin{array}{c|c} g_{\alpha\beta} + \lambda^2 \rho^2 \ell_{ab} A^a{}_\alpha A^b{}_\beta & \lambda \rho^2 \ell_{cb} A^c{}_\alpha \\ \hline \lambda \rho^2 \ell_{cb} A^c{}_\beta & \rho^2 \ell_{ab} \end{array} \right), \quad (3.17)$$

where

$$\ell_{ab} = h_{ab} + \mu k_{ab}.$$

This frame is also unholonomic. One easily finds

$$dv^a = \frac{-1}{2\lambda} C^a{}_{bc} v^b \wedge v^c. \quad (3.18)$$

The nonsymmetric theory of gravitation uses the nonsymmetric metric $g_{\mu\nu}$ such that

$$g_{\mu\nu} g^{\beta\nu} = g_{\nu\mu} g^{\nu\beta} = \delta_\mu^\beta, \quad (3.19)$$

where the order of indices is important. If G is semisimple and $k_{ab} = 0$

$$\ell_{ab} = h_{ab}, \quad \det(h_{ab}) \neq 0$$

and

$$h_{ab} h^{bc} = \delta_a^b. \quad (3.20)$$

Thus one easily finds in this case:

$$\gamma_{AC} \gamma^{BC} = \gamma_{CA} \gamma^{CB} = \delta^B{}_A. \quad (3.21)$$

where the order of indices is important. We have the same for the electromagnetic case ($G = \mathbf{U}(1)$). In general if $\det(\ell_{ab}) \neq 0$ then

$$\ell_{ab} \ell^{ac} = \ell_{ba} \ell^{ca} = \delta_b^c, \quad (3.22)$$

where the order of indices is important. From (3.22) we have (3.21) for the general nonsymmetric metric γ .

It is easy to see that

$$\begin{aligned} \Phi'(g) \bar{\gamma} &= \bar{\gamma}, \\ \Phi'(g) \underline{\gamma} &= \underline{\gamma} \end{aligned} \quad (3.23)$$

and γ_{AB} is invariant tensor with respect to the right-action of the group G on $\underline{P}$ (this means it is covariant and right-invariant).

In the case of any abelian group the condition (3.23) is stronger and we get that γ_{AB} is bi-invariant. Thus in the case of $G = \mathbf{U}(1)$ (electromagnetic case)

$$\xi_5 \overline{\gamma} = 0 = \xi_5 \underline{\gamma}, \quad (3.24)$$

where ξ_A is a dual base

$$\theta^A(\xi_B) = \delta^A_B, \quad (3.25)$$

$A, B = 1, 2, 3, 4, 5$

$$\xi_A = (\xi_\alpha, \xi_5). \quad (3.26)$$

Let us come back to the connection $\widehat{\omega}^a_b$ defined on the group G . For a typical fibre is diffeomorphic to G we can define $\widehat{\omega}^a_b$ on every fibre $F_x \simeq G$, $x \in E$. Due to a local trivialization of the bundle P we can define $\widehat{\omega}^a_b$ on every set $U \times G$, where $U \subset E$ and is open. Thus we get a linear connection on P such that

$$\widehat{\omega}^A_B = \left(\begin{array}{c|c} 0 & 0 \\ \hline 0 & -\frac{1}{\lambda} c^a_{bc} \theta^c \end{array} \right) \quad (3.27)$$

defined in a frame $\theta^A = (\pi^*(\overline{\theta}^\alpha), \theta^a)$ where $\overline{\theta}^\alpha$ is a frame on E and θ^a is a horizontal lift base.

This connection can be examined in a systematic way. Let us introduce a metric on $\underline{P}$ in a following way

$$p = \pi^* \eta \oplus h_{ab} \theta^a \otimes \theta^b, \quad (3.28)$$

where $\eta = \eta_{\mu\nu} \overline{\theta}^\mu \otimes \overline{\theta}^\nu$ is a Minkowski tensor and h_{ab} is a Killing–Cartan tensor on G . One gets

$$p_{AB} = \left(\begin{array}{c|c} \eta_{\alpha\beta} & 0 \\ \hline 0 & h_{ab} \end{array} \right) \text{ and } p^{AB} = \left(\begin{array}{c|c} \eta^{\alpha\beta} & 0 \\ \hline 0 & h^{ab} \end{array} \right). \quad (3.29)$$

The connection $\widehat{\omega}^A_B$ can be defined as

$$\widehat{\omega}^A_B = \left(\begin{array}{c|c} \pi^* \widehat{\omega}^\alpha_\beta & 0 \\ \hline 0 & \phi_x^0 \widehat{\omega}^a_b \end{array} \right), \quad (3.27a)$$

where $\widehat{\omega}^\alpha_\beta$ is a trivial connection on the Minkowski space, $\widehat{\omega}^a_b$ is the connection defined in sec. 2 and ϕ_x is a diffeomorphism $\phi_x : F_x \rightarrow G$, $x \in U$.

It is easy to check that

$$\widehat{D}p_{AB} = 0 = \widehat{D}p^{AB}, \quad (3.30)$$

where $\widehat{D}$ is an exterior covariant differential with respect to $\widehat{\omega}^A_B$. One can easily calculate the torsion for $\widehat{\omega}^A_B$

$$\widehat{Q}^a_{\mu\nu} = \lambda H^a_{\mu\nu}, \quad (3.31a)$$

$$\widehat{Q}^a_{bc} = \frac{1}{\lambda} C^a_{bc} \quad (3.31b)$$

and the curvature tensor

$$\widehat{R}^a{}_{b\mu\nu} = \lambda X_b H^a{}_{\mu\nu} \quad (3.32)$$

(the remaining torsion and curvature components are zero).

The connection $\widehat{\omega}^A{}_B$ is neither flat nor torsionless. Moreover it is still metric as a connection $\widehat{\omega}^a{}_b$ from sec. 2.

The covariant differentiation with respect to this connection is connected to the right action of the group G on $\underline{P}$. Thus the condition of the right-invariance of the p -form $\Xi^{A_1 \dots A_l}{}_{B_1 \dots B_m}$ on $\underline{P}$ is equivalent to

$$\widehat{\nabla}_a \Xi^{A_1 \dots A_l}{}_{B_1 \dots B_m} = 0, \quad (3.33)$$

where $\widehat{\nabla}_k$ is a covariant derivative with respect to $\widehat{\omega}^A{}_B$ in vertical directions on $\underline{P}$. This means right-invariance of Ξ . This can be written

$$\widehat{\nabla}_{\text{ver}(X)} \Xi = 0, \quad (3.33a)$$

ver is understood in the sense of ω .

$$\Phi'(g) \Xi = \Xi, \quad (3.34)$$

where $g \in G$ and $\Xi = (\Xi)^{A_1 \dots A_l B_1 \dots B_m} = (p^{B_1 B'_1} p^{B_2 B'_2} \dots p^{B_m B'_m} \Xi^{A_1 \dots A_l}{}_{B_1 \dots B_m})$.

For a connection on a bundle $\underline{P}$, ω , its curvature Ω one gets

$$\widehat{\nabla}_k \omega = \widehat{\nabla}_k \Omega = 0. \quad (3.34*)$$

Thus we can rewrite Eq. (3.23)

$$\widehat{\nabla}_a \underline{\gamma} = \widehat{\nabla}_a \overline{\gamma} = 0. \quad (3.35)$$

This means that

$$\widehat{\nabla}_a \gamma_{AB} = 0, \quad (3.36)$$

or

$$\widehat{\nabla}_{\text{ver}(X)} \gamma = 0. \quad (3.36a)$$

For every linear connection $\omega^A{}_B$ defined on $\underline{P}$ compatible in some sense with γ_{AB} we get

$$\Phi^*(g) \omega_{AB} = \text{Ad}_g \omega_{AB}, \quad (3.37)$$

which means that ω_{AB} is right-invariant with respect to the right-action of the group G on P . The same we say for a 2-form of torsion and 2-form of curvature derived for $\omega^A{}_B$ i.e.

$$\widehat{\nabla}_a \Omega^A{}_B = \widehat{\nabla}_a \Theta^A = 0. \quad (3.38)$$

The curvature scalar is invariant with respect to the right action of the group G on P .

$$0 = \widehat{\nabla}_a R = X_a R = \zeta_a R. \quad (3.39)$$

The condition (3.37) is the same as in the classical Kaluza–Klein (Jordan–Thiry) theory in a nonabelian case. A parallel transport with respect to the connection $\hat{\omega}^A_B$ means simply a right action of the group G on $\underline{P}$.

Our subject of investigations consists in looking for a generalization of the geometry from Einstein Unified Field Theory (the so called Einstein–Kaufmann Theory (see [4,5], [61])) defined on P i.e. for a connection ω^A_B such that

$$D\gamma_{AB} = \gamma_{AD}Q^D_{BE}\theta^E, \quad (3.40)$$

where D is an exterior, covariant differential with respect to the connection ω^A_B and Q^D_{BE} is a tensor of torsion for ω^A_B . We suppose that this connection is right-invariant with respect the right action of the group G .

We can write Eqs. (3.37–39) for a torsion, curvature and a scalar of curvature for ω^A_B . In this way we consider Einstein–Kaufmann G -structure on the bundle of linear frames over the manifold $\underline{P}$ (i.e. a right G -structure).

We can repeat all the considerations changing right(left)-invariant into left(right)-invariant in all places.

In this section we define ω^A_B as a collection of 1-forms defined on the manifold $\underline{P}$ (a gauge bundle manifold) and we choose for ω^A_B a lift horizontal frame (connected to the connection ω on the gauge bundle).

The collection of 1-forms ω^A_B becomes a linear connection on $\underline{P}$ iff it satisfies the following transformation properties

$$\omega^{A'}_{B'} = \Sigma^{-1A'}_A(p)\omega^A_B\Sigma^{-1B}_{B'}(p) - \Sigma^{-1A'}_A(p)d\Sigma^A_{B'}(p), \quad (3.41)$$

where

$$\Sigma(p) \in \text{GL}(n+4, \mathbb{R}), \quad p \in U_p \subset P$$

and

$$\theta^C = \Sigma^C_{C'}(p)\theta^{C'} \quad (3.42)$$

is a simultaneous transformation property of a frame. Having ω^A_B with (3.41–42) transformation properties we can lift it on a principal fibre bundle of frames over $\underline{P}$ with the structural group $GL(n+4, \mathbb{R})$ getting a 1-form of connection $\tilde{\omega}$

$$\tilde{\omega}_p = \text{Ad}_{\text{GL}(n+4, \mathbb{R})}(g_p^{-1})[\Pi^*(\omega^A_B X^B_A) - g_p^{-1}dg_p], \quad (3.43)$$

where Π is a projection defined on this principal fibre bundle of frames and

$$g_p : z \in \Pi^{-1}(U_p) \rightarrow g_p(z) = (\text{pr}_{\text{GL}(n+4, \mathbb{R})}\Psi_p(z))^{-1} \in \text{GL}(n+4, \mathbb{R}), \\ p \in U_p \subset \underline{P},$$

pr means a projection on $\text{GL}(n+4, \mathbb{R})$ in a local trivialization of the bundle P'' , Ψ is an action of $\text{GL}(n+4, \mathbb{R})$ on principal fibre bundle of frames over $\underline{P}$, $\Psi \rightarrow \text{GL}(n+4, \mathbb{R})$

$4, \mathbb{R}) \times P'' \rightarrow P''$, Ψ_p is defined for $\text{GL}(n+4, \mathbb{R}) \times U_p$. In this way we have an action of $\text{GL}(n+4, \mathbb{R})$ on the bundle and for $\tilde{\omega}$.

$$\Psi^*(g)\tilde{\omega} = \text{Ad}_{\text{GL}(n+4, \mathbb{R})}(g^{-1})\tilde{\omega}, \quad (3.44)$$

X^A_B are generators of the Lie algebra $\mathfrak{gl}(n+4, \mathbb{R})$ of $\text{GL}(n+4, \mathbb{R})$ and $g \in \text{GL}(n+4, \mathbb{R})$. For a soldering forms $\tilde{\theta}^A$ one gets

$$\tilde{\theta}^A = g_p \Pi^*(\theta^A). \quad (3.45)$$

Taking any two sections of the principal fibre bundle of P'' frames E and F such that

$$\begin{aligned} E^*\tilde{\omega} &= \omega^{A'}_{B'} X^{B'}_{A'}, \\ F^*\tilde{\omega} &= \omega^A_B X^B_A, \end{aligned} \quad (3.46)$$

$$\begin{aligned} E^*\tilde{\theta}^A &= \theta^{A'}, \\ F^*\tilde{\theta}^A &= \theta^A, \end{aligned} \quad (3.47)$$

one gets transformation properties (3.41) and (3.42). In such a way that

$$E(p) = F(p)\Sigma(p). \quad (3.48)$$

Eq. (3.40) can be rewritten in a more compact form

$$\nabla\gamma = S, \quad (3.49)$$

where

$$\begin{aligned} S(X, Y, Z) &= [\text{Tr}(\gamma \otimes Q)](X, Y, Z) = \sum_A \gamma(X, e_A) \theta^A(Q(Y, Z)), \\ Q(Y, Z) &= -Q(Z, Y), \end{aligned}$$

is a torsion of the connection $\tilde{\omega}$, X, Y, Z are contravariant vector fields and θ^A, e_B

$$\theta^A(e_B) = \delta^A_B \text{ are dual bases.}$$

Or in a different shape

$$\nabla_Z \gamma(X, Y) = S(X, Y, Z), \quad (3.50)$$

∇ is a covariant derivative with respect to the connection $\tilde{\omega}$ on the fibre bundle of frames.

Moreover now we consider γ, Q, X, Y, Z etc. as geometrical objects living on appropriate associate fibre bundles to the fibre bundle of frames. The condition (3.50) gives us Einstein–Kaufmann connection $\tilde{\omega}$ on the principal fibre bundle of frames over P . For $\tilde{\omega}$ is right-invariant with respect to the action of group G on this bundle of frames (lifted to this bundle from $\underline{P}$, the condition (3.50) is also right-invariant. Let us consider the right-invariance of the Einstein connection on $(\underline{P}, \gamma)$ from a different point of view.

Let $\Phi : G \times \underline{P} \rightarrow \underline{P}$ be a right action of the group G on $\underline{P}$ and let $\Phi^*(g)$ be a contragradient map to $\Phi'(g)$, a tangent map to Φ at $g \in G$. Let $\sum : G \rightarrow \text{GL}(n+4, \mathbb{R})$ be a homomorphism of groups and let us consider a connection $\tilde{\omega}$ on a fibre bundle of

frames over P with a structural group $\text{GL}(n+4, \mathbb{R})$ compatible in Einstein-Kaufmann sense with the nonsymmetric tensor γ . The connection is right-invariant with respect to the action of the group G on P if one has $\hat{\Phi}^*(g)\tilde{\omega} = \text{Ad}_{\text{GL}(n+4, R)}(\Sigma(g^{-1}))\tilde{\omega}$ ($\hat{\Phi}$ is an action of G on P lifted to this bundle) or for any local section E of the bundle of frames over $\underline{P}$

$$\Phi^*(g)\Gamma = \text{Ad}_{\text{GL}(n+4, R)}\left(\Sigma(g^{-1})\right)\Gamma + \Sigma^{-1}(g)d\Sigma(g), \quad (3.51)$$

where $\Gamma = E^*\tilde{\omega}$

$$\Gamma = \Gamma^A_{BC}\theta^C X^B_A \text{ and}$$

Thus one gets

$$\begin{aligned} \Phi^*(g)\Gamma^{A'}_{B'C'}, &= \Sigma^{A'}_A(g^{-1})\Gamma^A_{BC'}(g^{-1})\Sigma^B_{B'}(g^{-1})\Sigma^C_{C'}(g^{-1}) + \\ &+ \Sigma^{-1A'}_A(g)d_C\Sigma^A_{B'}(g)\Sigma^C_{C'}(g), \end{aligned} \quad (3.52)$$

where d_C is a vector field dual to θ^C . The reper transforms

$$\Phi^*(g)\theta^C = \Sigma^C_{C'}(g^{-1})\theta^{C'}. \quad (3.52')$$

Let us take the lift horizontal basis. In this case one gets

$$\Sigma^A_B(g) = \left(\begin{array}{c|c} \delta^\alpha_\beta & 0 \\ \hline 0 & \overline{U}^a_b(g) \end{array} \right), \quad \Sigma(g) = \text{id}_E \otimes \overline{U}(g), \quad (3.53)$$

where $U^a_b(g) = (\text{Ad}_G(g^{-1}))^a_b$ is a matrix of the adjoint representation of G . Thus we have

$$\begin{aligned} \Phi^*(g)\theta^\alpha &= \theta^\alpha, \\ \Phi^*(g)\theta^a &= \overline{U}^a_{a'}(g)\theta^{a'} \end{aligned} \quad (3.54)$$

$$\begin{aligned} \Phi^*(g)\Gamma^\alpha_{\beta\gamma} &= \Gamma^\alpha_{\beta\gamma}, \\ \Phi^*(g)\Gamma^{a'}_{\beta'\gamma'} &= \overline{U}^{a'}_a(g^{-1})\Gamma^a_{\beta'\gamma'}, \end{aligned} \quad (3.55)$$

$$\begin{aligned} \Phi^*(g)\Gamma^{a'}_{\beta'c'} &= \overline{U}^{a'}_a(g^{-1})\Gamma^a_{\beta'c}U^c_{c'}(g), \\ \Phi^*(g)\Gamma^{a'}_{b'\gamma} &= \overline{U}^{a'}_a(g^{-1})\Gamma^a_{b\gamma}U^b_{b'}(g^{-1}), \\ \Phi^*(g)\Gamma^{\alpha'}_{\beta'a'} &= \Gamma^\alpha_{\beta'a}\overline{U}^a_{a'}(g^{-1}). \end{aligned}$$

For $\omega^a_b = \tilde{\Gamma}^a_{bc}\theta^c$ one gets

$$\Phi^*(g)\omega^{a'}_{b'} = \overline{U}^{a'}_a(g^{-1})\omega^a_b\overline{U}^b_{b'}(g^{-1}). \quad (3.56)$$

In this way $\tilde{\Gamma}^a_{bc}$ has tensorial properties in the lift horizontal basis (Ad-type).

Moreover Eq. (3.56) has a natural interpretation a right-invariancy of the connection $\tilde{\omega}^a_b$ with respect to the right action of the group G on G . The second equation of

Eq. (3.54) means Ad-property of the connection on the principal fibre bundle $\underline{P}$ (a gauge bundle).

Eq. (3.56) can be rewritten in a more familiar form

$$R^*(g)\tilde{\Gamma} = \text{Ad}\left(\widetilde{\sum}(g^{-1})\right)\tilde{\Gamma} + \widetilde{\sum}(g)d\widetilde{\sum}(g), \quad (3.57)$$

where $\tilde{\Gamma} = \tilde{\omega}^a_b Y^b_a$ and Y^b_a are generators of the Lie algebra $gl(n, R)$ of the group $\text{GL}(n, \mathbb{R})$ and $\text{Ad}_{\text{GL}(n, R)}$ is an adjoint representation of $\text{GL}(n, \mathbb{R})$ and R is a right action of G on G .

$$\widetilde{\sum} : G \rightarrow \text{GL}(n, \mathbb{R}) \quad (3.58)$$

is a smooth homomorphism of groups such that

$$\widetilde{\sum}^a_b(g) = (\text{Ad}_G(g^{-1}))^a_b = \bar{U}^a_b(g). \quad (3.59)$$

In this way we are coming to the notion of the fibre bundle of frames over a group G and to the right invariant connection defined on this bundle. Eq. (3.57) can be rewritten:

$$\hat{R}^*(g)\hat{\omega} = \hat{\omega}, \quad (3.60)$$

where $\hat{\omega}$ is a connection on the principal fibre bundle of frames over G with the structural group $\text{GL}(n, \mathbb{R})$ and

$$\tilde{\Gamma} = f^*\hat{\omega}, \quad (3.61)$$

f is a local section of this bundle. $\hat{R}$ is an action of G on G lifted to the bundle of frames.

Let us notice that our considerations are valid for any connection defined on a fibre bundle of frames over $\underline{P}$, not only for Einstein–Kaufmann one. The same we can say for a connection on a fibre bundle of frames over G . The above considerations justify some Ad_G -properties of ω^A_B defined on manifold $\underline{P}$ (a gauge bundle) and $\tilde{\omega}^a_b$ defined on G (a group manifold). They are treated there as some 1-forms defined on P or G according to our conventions. From Eq. (3.51) one easily gets transformation laws for the curvature

$$\hat{\Phi}^*(g)\tilde{\Omega} = \tilde{\Omega} \quad (3.62)$$

and from (3.60)

$$\hat{R}^*(g)\tilde{\Omega} = \tilde{\Omega}. \quad (3.63)$$

For the curvature scalars we get

$$\hat{\Phi}^*(g)R(\tilde{\omega}) = R(\tilde{\omega}), \quad (3.64)$$

$$\hat{R}^*(g)R(\tilde{\omega}) = R(\tilde{\omega}). \quad (3.65)$$

In this work we consider a non-symmetric metrization of the principal fibre bundle defined over a base manifold $V = E \times G/G_0$, where E is a space-time and $M = G/G_0$ is a certain homogeneous space.

4. Preliminary Remarks

Let us introduce the principal fibre bundle P over the $V = E \times M$ with the structural group H and with the projection π , where $M = G/G_0$ is a homogeneous space, G is a compact Lie group (semisimple), $G_0 \subset G$ is a semisimple compact subgroup of G and let $\omega = \omega^a X_a$ is a connection defined on P (Fig. 2A). Notice that now we have in the place of E , $E \times G/G_0$ and in the place of G , H . Let us suppose that (V, γ) is a manifold with a nonsymmetric metric tensor

$$\gamma_{AB} = \gamma_{(AB)} + \gamma_{[AB]}. \quad (4.1)$$




Fig. 2A. Principal fibre bundles P and Q

The signature of the tensor γ is $(+ - - -, \underbrace{- - - \dots -}_{n_1})$. Let us introduce a natural frame on $\underline{P}$

$$\theta^{\tilde{A}} = (\pi^*(\theta^A), \theta^a = \lambda \omega^a), \quad \lambda = \text{const}. \quad (4.2)$$

It is convenient to introduce the following notations. Capital Latin indices with tilde $\tilde{A}, \tilde{B}, \tilde{C}$ run $1, 2, 3, \dots, m+4$, $m = \dim H + \dim M = n + \dim M = n + n_1$, $n_1 = \dim M$, $n = \dim H$. Lower Greek indices $\alpha, \beta, \gamma, \delta = 1, 2, 3, 4$ and lower Latin cases $a, b, c, d = n_1 + 5, n_1 + 6, \dots, m+4$. Capital Latin indices $A, B, C = 1, 2, 3, \dots, n_1 + 4$, lower Latin cases with tilde $\tilde{a}, \tilde{b}, \tilde{c}$ run $5, 6, \dots, n_1 + 4$. The symbol “ $-$ ” over θ^A and over other quantities indicates that these quantities are defined on V . We have of course $\dim H = n$, $\dim M = \dim(G/G_0) = n_1$, $\dim G = n_2$, $\dim G_0 = n_2 - n_1$, and

$m = n_1 + n$. It is easy to see that the existence of the nonsymmetric metric on V is equivalent to introducing the two independent geometrical quantities on E .

$$\bar{\gamma} = \gamma_{AB} \bar{\theta}^A \otimes \bar{\theta}^B = \gamma_{(AB)} \bar{\theta}^A \otimes \bar{\theta}^B, \quad (4.3)$$

$$\underline{\gamma} = \gamma_{AB} \bar{\theta}^A \wedge \bar{\theta}^B = \gamma_{[AB]} \bar{\theta}^A \wedge \bar{\theta}^B, \quad (4.4)$$

i.e. the symmetric metric tensor $\bar{\gamma}$ on V and the 2-form $\underline{\gamma}$. On the group H we can introduce a bi-invariant symmetric tensor called the Killing–Cartan tensor.

$$h(A, B) = h_{ab} A^a \cdot B^b, \quad (4.5)$$

where

$$h_{ab} = h_{ba}, \quad A = A^a X_a, \quad B = B^a X_a,$$

X_a —are generators of a Lie algebra of H , $(\mathfrak{h})$.

It is supposed that H is semisimple; it means that $\det(h_{ab}) \neq 0$. Let us introduce a skewsymmetric right-invariant tensor on H

$$k(A, B) = k_{bc} A^b \cdot B^c, \quad k_{bc} = -k_{cb}. \quad (4.6)$$

Let us turn to the nonsymmetric metrization of $\underline{P}$. Let us suppose that

$$\bar{\varkappa}(X, Y) = \bar{\gamma}(\pi' X, \pi' Y) + \rho^2 \lambda^2 h(\omega(X), \omega(Y)), \quad (4.7)$$

$$\underline{\varkappa}(X, Y) = \underline{\gamma}(\pi' X, \pi' Y) + \rho^2 \xi \lambda^2 k(\omega(X), \omega(Y)), \quad (4.8)$$

$\xi = \text{const}$ and is dimensionless, $X, Y \in \text{Tan}(\underline{P})$ and $\rho = \rho(x)$ is a scalar field on E . It is easy to see that:

$$\bar{\varkappa} = \pi^* \bar{\gamma} \oplus \rho^2 h_{ab} \theta^a \otimes \theta^b, \quad (4.9)$$

$$\underline{\varkappa} = \pi^* \underline{\gamma} \oplus \rho^2 \xi k_{ab} \theta^a \wedge \theta^b, \quad (4.10)$$

For

$$\varkappa_{\tilde{A}\tilde{B}} = \varkappa_{(\tilde{A}\tilde{B})} + \varkappa_{[\tilde{A}\tilde{B}]}$$

one easily gets

$$\varkappa_{\tilde{A}\tilde{B}} = \left(\begin{array}{c|c} \gamma_{AB} & 0 \\ \hline 0 & \rho^2 \ell_{ab} \end{array} \right), \quad (4.11)$$

where $\ell_{ab} = h_{ab} + \xi k_{ab}$ and $\det(\ell_{ab}) \neq 0$. One easily writes:

$$\varkappa_{\tilde{A}\tilde{C}} \varkappa^{\tilde{B}\tilde{C}} = \varkappa_{\tilde{C}\tilde{A}} \varkappa^{\tilde{C}\tilde{B}} = \delta^{\tilde{B}}_{\tilde{A}}, \quad (4.12)$$

where the order of indices is important. The signature of the tensor $\varkappa$ is $(+ \dots, \underbrace{\dots}_{n_1}, \underbrace{\dots}_{n})$. If $\det(\ell_{ab}) \neq 0$ then

$$\ell_{ab} \ell^{ac} = \ell_{ba} \ell^{ca} = \delta_b^c, \quad (4.13)$$

where the order of indices is important.

$$[X_a, X_b] = C_{ab}^c X_c \quad (4.13a)$$

are commutation relation for the Lie algebra of H , $(\mathfrak{h})$.

Let us pass to the nonsymmetric metrization of the homogeneous space G/G_0 . Let M be a homogeneous space for G (with left action of group G) (see [86], [87]). Let us suppose that the Lie algebra of G , $\mathfrak{g}$ has the following reductive decomposition

$$\mathfrak{g} = \mathfrak{g}_0 \dot{+} \mathfrak{m}, \quad (4.14)$$

where $\mathfrak{g}_0$ is a Lie algebra of G_0 (a subalgebra of $\mathfrak{g}$) and $\mathfrak{m}$ (the complement to the subalgebra $\mathfrak{g}_0$) is $\text{Ad } G_0$ invariant, $\dot{+}$ means a direct sum. Such a decomposition might not be unique in general, but we shall assume that one has been chosen. Sometimes one assumes a stronger condition for $\mathfrak{m}$, the so called symmetry requirement

$$[\mathfrak{m}, \mathfrak{m}] \subset \mathfrak{g}_0.$$

Let us introduce the following notation for generation of $\mathfrak{g}$.

$$Y_i \in \mathfrak{g}, \quad Y_{\bar{i}} \in \mathfrak{m}, \quad Y_{\bar{a}} \in \mathfrak{g}_0.$$

This is a decomposition of a basics of $\mathfrak{g}$ according to (4.14).

If G is a connected Lie group and G_0 is a closed subgroup, G/G_0 is the space of cosets $\{gG_0\}$, $\varphi : G \rightarrow G/G_0$ is defined by $\varphi(g) = \{gG_0\}$, $g \in G$. Let us suppose that G is a compact Lie group and define a field of 2-form $k^0(x)$, $x \in M$ on G/G_0 . Let us notice that $G \xrightarrow{\varphi} M = G/G_0$ gives us a principal fibre bundle over M with a projection φ and a structural group G_0 . We use the complement $\mathfrak{m}$ in order to introduce a coordinate system on G/G_0 . Let $g \in G$ and let “exp” be an exponential mapping $\exp : \mathfrak{g} \rightarrow G$. Thus

$$(x_5, x_6, \dots, x_{n_1+4}) \rightarrow \varphi\left(g \exp \sum_{\bar{a}} x_{\bar{a}} Y_{\bar{a}}\right) \quad (*)$$

is a diffeomorphism of a neighbourhood of zero in $\mathfrak{m}$ on a neighbourhood of $\varphi(g)$ in G/G_0 . An inverse mapping is a local coordinate system in a neighbourhood of $\varphi(g) \in G/G_0$ on $M = G/G_0$

The mapping $\varphi : G \rightarrow G/G_0$ has a tangent mapping φ' which maps $\mathfrak{g}$ in a tangent space to $M = G/G_0$ at a point $o = \{eG_0\}$. A kernel of φ' is equal to $\mathfrak{g}_0$. The operator of left action $\tau(g) : hG_0 \rightarrow ghG_0$ satisfies $\varphi o L(g) = \tau(g) o \varphi$, $h \in G$.

For $\varphi o R_{g_0} = \varphi$, $\text{Ad}_G(g)Y = R'_{g^{-1}} o L'_g(Y)$ we get $\varphi' o \text{Ad}_g(g_0)Y = \tau'(g_0)_o o \varphi'(Y)$, $Y \in \mathfrak{g}$.

Thus $\mathfrak{m} \simeq \mathfrak{g}/\mathfrak{g}_0 \simeq \text{Tan}_0(G/G_0)$ and using the left action of the group G on M we can get vector fields at any point $x \in M$ from $\text{Tan}_0(M) \simeq \mathfrak{m}$. Thus let $Y_{\bar{a}} \in \mathfrak{m}$. Due to an isomorphism of $\mathfrak{m}$ and $\text{Tan}_0(M)$ we get

$$\tilde{Y}_{\bar{a}}(x) = L'_g(o), \quad x = go, \quad g \in G,$$

$Y(x)$ is left invariant on M . If $k(g)$ is a field of 2-forms on G and being left-invariant we define a field of 2-form $k^0(x)$ on M in the following way: Let us restrict it to the subspace $\mathfrak{m}$ (a complement)

$$k'(g) = k(g)|_{\mathfrak{m}},$$

i.e. $\bigwedge_{g \in G} k(g)|_{\mathfrak{m}}(x, y) = 0$ if x or y do not belong to $\mathfrak{m}$ and consider a 2-form $k^0(x)$.

The form $k(g)$ is left-invariant on G . The form $k(g)|_{\mathfrak{m}}$ is G_0 Ad'-invariant, for $\mathfrak{m}$ is Ad'_{G_0} invariant. Let us consider a local section $\sigma_1 : U_1 \rightarrow G$, $U_1 \subset M$ of the bundle G . We get

$$k^{\sigma_1}(x) = \sigma_1^* k(\sigma_1(x))|_{\mathfrak{m}}, \quad x \in U_1. \quad (4.15)$$

Let us take a different local section $\sigma_2 : U_2 \rightarrow G$, $U_2 \subset M$

$$k^{\sigma_2}(x) = \sigma_2^* k(\sigma_2(x))|_{\mathfrak{m}}. \quad (4.16)$$

One has

$$\sigma_2(x) = p(x)\sigma_1(x), \quad (4.17)$$

where $p(x) \in G_0$

$$x \in U_1 \cap U_2.$$

Thus one obtains

$$k^{\sigma_2}(x) = R_{p(x)}^* k^{\sigma_1}(x) = k^{\sigma_1}(x). \quad (4.18)$$

Thus we get

$$k^0(x) = k^{\sigma}(x), \quad (4.19)$$

for any local section of G . Let us examine a transformation properties of k^0 with respect to the left action of the group G on M . There is a natural action of G on left cosets, such that G acts with an isotropy subgroup G_0 . One gets

$$L_g^* k^0(x) = k^0(x). \quad (4.21)$$

In this way $k^0(x)$ is left invariant with respect to the action of the group G on $M = G/G_0$.

This fact allows us to define k being left-invariant on G in such a way that $k = \varphi^* k^0$ and k^0 is left-invariant 2-form on $M = G/G_0$. This gives us a construction of a left-invariant 2-form on G if we have a left-invariant 2-form on $M = G/G_0$.

In the same way we can define a tensor field $h^0(x)$ on G/G_0 , $x \in G/G_0$ using tensor field h on G . Moreover if we suppose that h is a biinvariant (Ad-invariant) metric on G (a Killing–Cartan tensor) we have a simpler construction.

The complement $\mathfrak{m}$ is a tangent space to the point $\{\varepsilon G_0\}$ of M , ε is a unit element of G . We restrict h to the space $\mathfrak{m}$ only. Thus we have $h^0(\{\varepsilon G_0\})$ at the one point of M . Now we propagate $h^0(\{fG_0\})$ using a left action of the group G .

$$h^0(\{fG_0\}) = (L_f^{-1})^*(h^0(\{\varepsilon G_0\})), \quad (4.22)$$

$h^0(\{\varepsilon G_0\})$ is of course $\text{Ad } G_0$ invariant tensor defined on $\mathfrak{m}$ and

$$L_f^* h^0 = h^0,$$

where $f \in G$.

Let us find $k^0(x)$ in the case of $\text{SO}(3)/\text{SO}(2) \simeq S^2$, where $\text{SO}(2)$ subgroup is defined by the exponentiation of e_3 , i.e. rotations around an axis z and $\mathfrak{m} \cong \text{Span}\{e_1, e_2\}$. One finds

$$k(g)|_{\mathfrak{m}} = k'(g) = 2k_{12}v^1 \wedge v^2,$$

where v^1 and v^2 are dual with respect to e_1 and e_2 being right-invariant vector fields on $\text{SO}(3)$. Thus

$$k'_{\tilde{a}\tilde{b}}(\vartheta, \phi, \psi) = \begin{pmatrix} 0 & \cos \vartheta \\ -\cos \vartheta & 0 \end{pmatrix}, \quad (4.23)$$

$$k^0_{\tilde{a}\tilde{b}}(\vartheta, \phi) = \begin{pmatrix} 0 & \cos \vartheta \\ -\cos \vartheta & 0 \end{pmatrix}. \quad (4.23c)$$

We can also try to find and classify such tensors for all homogeneous spaces. This will be done elsewhere. The classification of homogeneous spaces for which G and G_0 are nilpotent or solvable will be done elsewhere as well. In this case the same construction as for h^0 is applicable for k^0 (coming from bi-invariant form defined on a nilpotent or a solvable Lie algebra).

Let f^i_{jk} are the structure constants of the Lie algebra $\mathfrak{g}$, i.e.

$$[Y_j, Y_k] = f^i_{jk} Y_i, \quad (4.24)$$

Y_j are generators of Lie algebra $\mathfrak{g}$. Let us construct a natural left-invariant frame on $M = G/G_0$. To do this we choose the local section $\sigma : U \rightarrow G/G_0$, of the natural bundle $G \rightarrow G/G_0$ where $U \subset M = G/G_0$ (see Fig. 2B).




Fig. 2B. The principal fibre bundle G over $G/G_0 = M$ and some details on the constructions of the form $k^0(x)$ on M

The local section σ can be considered an introduction of coordinate system on M i.e (*). Let σ^* means a pull-back of σ and let us define

$$\omega_{\text{MC}}^\sigma = \sigma^* \omega_{\text{MC}}, \quad (4.25)$$

where ω_{MC} is a left-invariant Maurer–Cartan form. Using the decomposition (4.14) we have

$$\omega_{\text{MC}}^\sigma = \omega_0^\sigma + \omega_{\mathfrak{m}}^\sigma = \theta^i Y_i + \bar{\theta}^{\tilde{a}} Y_{\tilde{a}}. \quad (4.26)$$

It is easy to see that $\theta^{\tilde{a}}$ is the natural left-invariant frame on M and we have

$$h^0 = h_{\tilde{a}\tilde{b}}^0 \bar{\theta}^{\tilde{a}} \otimes \bar{\theta}^{\tilde{b}}, \quad (4.27)$$

$$k^0 = k_{\tilde{a}\tilde{b}}^0 \bar{\theta}^{\tilde{a}} \wedge \bar{\theta}^{\tilde{b}}. \quad (4.28)$$

According to our notation we have $\tilde{a}, \tilde{b} = 5, 6, \dots, n_1+4$. Thus we have a nonsymmetric metric on M

$$\gamma_{\tilde{a}\tilde{b}} = r^2(h_{\tilde{a}\tilde{b}}^0 + \zeta k_{\tilde{a}\tilde{b}}^0) = r^2 g_{\tilde{a}\tilde{b}}, \quad (4.29)$$

where ζ is a real dimensionless parameter and r has length-dimension. r characterizes the size of the manifold M and plays the role of a “radius”. The tensor field $g_{\tilde{a}\tilde{b}} \bar{\theta}^{\tilde{a}} \otimes \bar{\theta}^{\tilde{b}} = (h_{\tilde{a}\tilde{b}}^0 + \zeta k_{\tilde{a}\tilde{b}}^0) \bar{\theta}^{\tilde{a}} \otimes \bar{\theta}^{\tilde{b}}$ is left-invariant on $M = G/G_0$. For k^0 is left-invariant with respect to the action of the group G on M one gets

$$Y_\ell k^0 = 0 \quad (4.30)$$

and in particular

$$\begin{aligned} Y_{\tilde{\ell}} k^0 &= 0, \\ Y_{\tilde{\ell}} &\in \mathfrak{m} = \mathfrak{g}/\mathfrak{g}_0. \end{aligned} \quad (4.31)$$

Let $\widehat{\widehat{D}}$ be an exterior covariant derivative with respect to the Riemannian connection on M induced by the tensor field h^0 .

If

$$\widehat{\widehat{D}} k^0 = dk^0 = 0 \quad (4.32)$$

the form k^0 is closed and it can define a symplectic structure on M (of course if k^0 is nondegenerate and $\dim M = 2k$, i.e. it is even). In this case we have

$$\widehat{\widehat{\nabla}}_{[\tilde{a}\tilde{b}\tilde{c}]} k^0 = 0. \quad (4.33)$$

In this case we can define an almost complex structure compatible with the Riemannian and the symplectic structure on M i.e

$$J : T(M) \rightarrow T(M), \quad J^2 = -\text{id}, \quad (4.34)$$

such that

$$h^0(JX, Y) = k^0(X, Y), \quad (4.35)$$

for any $X, Y \in T(M)$.

If those three structure are compatible we have:

$$\widehat{\nabla}_z J = 0, \quad z \in T(M). \quad (4.36)$$

The almost complex structure J can be integrated into a complex structure (a unique one) if the torsion (torsion of J , not of a nonriemannian connection) vanishes i.e.

$$S(X, Y) = [X, Y] + J[JX, Y] + J[X, JY] - [JX, JY] = 0, \quad X, Y \in T(M). \quad (4.37)$$

In this case we get Kählerian structure on the manifold M . If $\dim M = 2$, this can be always achieved.

Thus in the case of $\dim G/G_0 = 2$ we can always integrate the left-invariant symmetric tensor $g_{\tilde{a}\tilde{b}}$ introduced in this section if $\widehat{\nabla}_{\tilde{c}} k_{\tilde{a}\tilde{b}}^0 = 0$ into Kählerian geometry.

In particular we get in this way for $M = S^2 = \text{SO}(3)/\text{SO}(2)$ a Kählerian geometry which is holomorphically equivalent to the standard one. Holomorphically means here also conformally.

In the case of Kählerian structure, one easily gets that

$$\widehat{D}g_{\tilde{a}\tilde{b}} = 0. \quad (4.38)$$

Eq. (4.38) has important consequences.

The Kählerian structure on $M = G/G_0$ (G -invariant) induces a left-invariant 2-form on M , k^0 . Moreover k^0 induces a skew-symmetric form on G . This has some important consequences, because we can induce on an G a left-invariant nonsymmetric tensor coming from Kählerian structure on G/G_0 .

Thus we are able to write down the nonsymmetric metric on $V = E \times M = E \times G/G_0$.

$$\gamma_{AB} = \left(\begin{array}{c|c} g_{\alpha\beta} & 0 \\ \hline 0 & r^2 g_{\tilde{a}\tilde{b}} \end{array} \right), \quad (4.39)$$

where

$$g_{\alpha\beta} = g_{(\alpha\beta)} + g_{[\alpha\beta]},$$

$$g_{\tilde{a}\tilde{b}} = h_{\tilde{a}\tilde{b}}^0 + \zeta k_{\tilde{a}\tilde{b}}^0,$$

$$k_{\tilde{a}\tilde{b}}^0 = -k_{\tilde{a}\tilde{b}}^0, \quad h_{\tilde{a}\tilde{b}}^0 = h_{\tilde{a}\tilde{b}}^0,$$

$\alpha, \beta = 1, 2, 3, 4$, $\tilde{a}, \tilde{b} = 5, 6, \dots, n_1 + 4 = \dim M + 4 = \dim G - \dim G_0 + 4$. The frame $\bar{\theta}^{\tilde{a}}$ is unholonomic one:

$$d\bar{\theta}^{\tilde{a}} = \frac{1}{2} \varkappa_{\tilde{b}\tilde{c}}^{\tilde{a}} \bar{\theta}^{\tilde{b}} \wedge \bar{\theta}^{\tilde{c}}, \quad (4.40)$$

where $\varkappa^{\tilde{a}}_{\tilde{b}\tilde{c}}$ are coefficients of nonholonomicity and they depend on the point of the manifold $M = G/G_0$ (they are not constant in general). They depend on the section σ and on the constants $f^{\tilde{a}}_{\tilde{b}\tilde{c}}$.

Let us pass to the group structure of our theory. We have there three groups H , G and G_0 which have up to now been disconnected. Let us suppose that there exists a homomorphism μ between G_0 and H

$$\mu : G_0 \rightarrow H, \quad (4.41)$$

such that the centralizer of $\mu(G_0)$ in H , C^μ is isomorphic to G . C^μ , the centralizer of $\mu(G_0)$ in H is the set of all elements of H which commute with elements of $\mu(G_0)$ (it is a subgroup of H , of course).

$$C^\mu = G, \quad (4.42)$$

This means that H has the following structure:

$$\mu(G_0) \otimes G \subset H. \quad (4.43)$$

If we suppose that μ is an isomorphism between G_0 and $\mu(G_0)$ we get:

$$G_0 \otimes G \subset H. \quad (4.44)$$

Let us denote μ' as the tangent map to μ at a unit element. Thus μ' is the differential of μ acting on the Lie algebra elements. Let us suppose that the connection ω on the fibre bundle P is invariant under group action of G on the manifold $V = E \times G/G_0$. According to Ref. [23] this means the following. Let e be a local section of P , $e : U \subset V \rightarrow P$ and $A = e^*\omega$. Then for every $g \in G$ there exists a gauge transformation ρ_g such that:

$$f^*(g)A = \text{Ad}_{\rho_g^{-1}} A + \rho_g^{-1} d\rho_g, \quad (4.45)$$

f^* means a pull-back of the action f of the group G on manifold V . According to Refs. [71], [72], [73] we are able to write a general form for such an ω . Following Ref. [72] we have:

$$\omega = \tilde{\omega}_E + \mu' \circ \omega_0^\sigma + \Phi \circ \omega_m^\sigma, \quad (4.46)$$

where $\omega_0^\sigma + \omega_m^\sigma = \omega_{\text{MC}}^\sigma$ are components of the pull-back of the Maurer–Cartan form from the decomposition (4.14), $\tilde{\omega}_E$ is the connection defined on the fibre bundle Q over space-time E with structural group C^μ and the projection π_E . Moreover $C^\mu = G$, and $\tilde{\omega}_E$ is a 1-form with values in the Lie algebra $\mathfrak{g}$. This connection describes the ordinary Yang–Mills' field with gauge group $G = C^\mu$ on the space-time E . Φ is a function on E with values in the space $\tilde{S}$ of linear maps:

$$\Phi : \mathfrak{m} \rightarrow \mathfrak{h} \quad (4.47)$$

satisfying

$$\Phi([X_0, X]) = [\mu' X_0, \Phi(X)], \quad X_0 \in \mathfrak{g}_0 \quad (4.48)$$

thus

$$\begin{aligned}\tilde{\omega}_E &= \tilde{\omega}_E^i Y_i, & Y_i &\in \mathfrak{g}, \\ \omega_0^\sigma &= \theta^{\hat{i}} Y_{\hat{i}}, & Y_{\hat{i}} &\in \mathfrak{g}_0, \\ \omega_{\mathfrak{m}}^\sigma &= \bar{\theta}^{\tilde{a}} Y_{\tilde{a}}, & Y_{\tilde{a}} &\in \mathfrak{m}.\end{aligned}$$

Let us write condition (4.46) in the base of left-invariant form $\theta^{\hat{i}}$ and $\bar{\theta}^{\tilde{a}}$, which span respectively dual spaces to $\mathfrak{g}_0$ and $\mathfrak{m}$.

It is easy to see that

$$\Phi \circ \omega_{\mathfrak{m}}^\sigma = \Phi_a^a(x) \bar{\theta}^{\tilde{a}} X_a, \quad X_a \in \mathfrak{h} \quad (4.49)$$

and

$$\mu' = \mu_{\hat{i}}^a \theta^{\hat{i}} X_a, \quad (4.50)$$

where $\Phi_a^a(x)$ is the scalar field on E and $\mu_{\hat{i}}^a$ is a constant matrix. From (4.48) one gets

$$\Phi_b^c(x) f_{\hat{i}\tilde{a}}^{\tilde{b}} = \mu_{\hat{i}}^a \Phi_{\tilde{a}}^b(x) C^c_{ab}, \quad (4.51)$$

where $f_{\hat{i}\tilde{a}}^{\tilde{b}}$ are structure constants of the Lie algebra $\mathfrak{g}$ and C^c_{ab} are structure constants of the Lie algebra $\mathfrak{h}$. (4.51) is a constraint on the scalar field $\Phi_a^a(x)$. For a curvature of ω one gets:

$$\begin{aligned}\Omega &= \frac{1}{2} H_{AB} \theta^A \wedge \theta^B X_c = \\ &= \frac{1}{2} \tilde{H}^i_{\mu\nu} \theta^\mu \wedge \theta^\nu \alpha_i^c X_c + \overset{\text{gauge}}{\nabla}_\mu \Phi_a^c \theta^\mu \wedge \theta^{\tilde{a}} X_c + \\ &\quad + \frac{1}{2} C^c_{ab} \Phi_a^a \Phi_b^b \theta^{\tilde{a}} \wedge \theta^{\tilde{b}} X_c - \frac{1}{2} \mu_{\hat{i}}^c f_{\tilde{a}\tilde{b}}^{\hat{i}} \theta^{\tilde{a}} \wedge \theta^{\tilde{b}} X_c - \frac{1}{2} \Phi_d^c f_{\tilde{a}\tilde{b}}^{\tilde{d}} \theta^{\tilde{a}} \wedge \theta^{\tilde{b}} X_c.\end{aligned} \quad (4.52)$$

Thus we have:

$$H_{\mu\nu}^c = \alpha_i^c \tilde{H}_{\mu\nu}^i, \quad (4.53)$$

$$H_{\tilde{a}\tilde{b}}^c = \overset{\text{gauge}}{\nabla}_\mu \Phi_a^c = -H_{\tilde{a}\mu}^c, \quad (4.54)$$

$$H_{\tilde{a}\tilde{b}}^c = C_{ab}^c \Phi_a^a \Phi_b^b - \mu_{\hat{i}}^c f_{\tilde{a}\tilde{b}}^{\hat{i}} - \Phi_d^c f_{\tilde{a}\tilde{b}}^{\tilde{d}}, \quad (4.55)$$

where $\overset{\text{gauge}}{\nabla}_\mu$ means gauge derivative with respect to the connection $\tilde{\omega}_E$ defined on the bundle $Q(E, G)$,

$$Y_i = \alpha_i^c X_c \quad (4.56)$$

and $\tilde{H}_{\mu\nu}^i$ is the curvature of the connection $\tilde{\omega}_E$ in the base $\{Y_i\}$, generators of the Lie algebra of the Lie group G , $\mathfrak{g}$, α_i^c is the matrix which connects $\{Y_i\}$ with $\{X_c\}$. Now we would like to remind that indices a, b, c refer to the Lie algebra $\mathfrak{h}$, $\tilde{a}, \tilde{b}, \tilde{c}$ to the space $\mathfrak{m}$ (tangent space to M), $\hat{i}, \hat{j}, \hat{k}$ to the Lie algebra $\mathfrak{g}_0$ and i, j, k to the Lie algebra

of the group G , $\mathfrak{g}$. The matrix α_i^c establishes a direct relation between generators of the Lie algebra of the subgroup of the group H isomorphic to the group G .

Finally let us consider a different way to introduce a G -invariant (left-invariant) skewsymmetric form on $M = G/G_0$. Let us suppose that G is a connected compact simple Lie group with center $\{\varepsilon\}$ and G_0 has nondiscrete center and is a maximal connected proper subgroup of G . Really we need suppose that a center of G_0 , Z_{G_0} is isomorphic to $U(1)$ (a circle group). In this case (see the first position of Ref. [86] p. 382) there is an Hermitian structure (really Kählerian) invariant with respect to the action of the group G .

Let us denote the unique G -invariant almost complex structure on M by $J^2 = -1$. J is integrable to complex structure and let us denote by h^0 a G -invariant Riemannian structure on M , which is **Hermitian** $h^0(JX, JY) = h^0(X, Y)$, $X, Y \in \text{Tan}(M)$ and $\widehat{\nabla}_x J = 0$, $\widehat{\nabla}$ is covariant derivative induced by a Riemannian connection on M .

This allows us to introduce a skewsymmetric G -invariant 2-form on M . One has

$$k^0(X, Y) \stackrel{\text{df}}{=} h^0(JX, Y). \quad (4.57)$$

One easily gets that

$$k^0(X, Y) = -k^0(Y, X), \quad (4.58)$$

$$\widehat{\nabla}_z k^0 = 0 \quad (4.59)$$

and k^0 is G -invariant.

Thus one can write

$$\widetilde{g}(X, Y) = h^0(X, Y) + \zeta k^0(X, Y) = h^0(X, Y) + \zeta h^0(JX, Y). \quad (4.60)$$

Let us notice that we get a stronger result, for a symmetric space then for a group G . The form k^0 is defined uniquely for any G -invariant metric. We can define the metric h^0 using a bi-invariant tensor on G (a Killing-Cartan tensor), supposing the reductive decomposition of $\mathfrak{g}$ or even the symmetry requirement. For the group G is compact any bi-invariant tensor on G is proportional to Killing-Cartan tensor. In this way any nonsymmetric metric on $M = G/G_0$ is proportional to (4.60). To be safe we can use in the place of h^0 , $\frac{1}{2}(h^0(X, Y) + h^0(JX, JY))$.

Thus we get a left-invariant (G -invariant) nonsymmetric tensor on G , such that

$$\widehat{\nabla}_x g = 0. \quad (4.61)$$

In particular let us notice that S^2 is an Hermitian symmetric space (and Kählerian) with $\text{SO}(3)$ invariant scalar product and $\text{SO}(3)$ invariant symplectic form.

In the first position of Ref. [86] p. 518 S.Helgason quoted all the Kählerian structures on the symmetric spaces of compact type:

$$\begin{aligned}
& \mathrm{SU}(p+q)/\mathrm{S}(U_p \times U_q), \quad \dim M = 2pq, \quad p, q = 1, 2, \dots, \\
& \mathrm{SO}(2n)/\mathrm{U}(n), \quad \dim M = n(n-1), \quad n = 1, 2, \dots, \\
& \mathrm{SO}(p+2)/\mathrm{SO}(p) \times \mathrm{SO}(2), \quad \dim M = 2p, \quad p = 2, 3, \dots, \\
& \mathrm{Sp}(n)/\mathrm{U}(n), \quad \dim M = n(n+1), \quad n = 1, 2, \dots, \\
& (\mathfrak{e}_{6(-78)}, \mathfrak{so}(10) + R), \quad \dim M = 32, \\
& (\mathfrak{e}_{7(-133)}, \mathfrak{e}_6 + R), \quad \dim M = 54.
\end{aligned} \tag{4.62}$$

On every of these manifold of even dimension (real) and compact there is a Kählerian structure. Thus there is a symmetric metric and a skewsymmetric 2-form (nondegenerate) inducing a nonsymmetric metric which is covariantly constant with respect to the Riemannian connection (induced by a symmetric metric tensor) and left-invariant with respect to the action of the group G on M . In this way a torsion of the connection compatible with nonsymmetric tensor on M vanishes. This has some interesting consequences, which we use in further sections. The skewsymmetric G -invariant 2-form on M induces in a natural way a skewsymmetric left- G -invariant form on G , which together with the Killing–Cartan form on G form a nonsymmetric metric on G . Now, however the covariant derivative of the skewsymmetric tensor G with respect to the Riemannian connection on G (induced by a Killing–Cartan tensor) does not vanish in general.

In this way the torsion of the connection compatible with the nonsymmetric tensor on G does not vanish. This is different as for the Hermitian homogeneous space with left-invariant 2-form. In sec. 5 we find a general shape for such a connection which can be applied for our case i.e for the skewsymmetric form obtained from the left-invariant 2-form on M .

Let us calculate how many nonsymmetric (Einstein–Kaufmann) left-invariant structure we get in this way from (4.62). One gets

$$\begin{aligned}
& E_6 - 1, \quad E_7 - 1, \\
& \mathrm{Sp}(n) - 1, \quad n = 1, 2, \dots, \\
& \mathrm{SU}(n) - \left[\frac{n+1}{2} \right], \quad n = 2, 3, \dots, \\
& \mathrm{SO}(2n+1) - 1, \quad n = 1, 2, \dots, \\
& \mathrm{SO}(2n) - 2, \quad n = 1, 2, \dots,
\end{aligned} \tag{4.63}$$

Thus except $\mathrm{SU}(n)$ we have really 1 or 2 inequivalent Einstein–Kaufmann connections on G obtained from Kählerian structure on a homogeneous space (being a symmetric space).

Moreover for lower dimensions we have some coincidences, i.e.

$$\begin{aligned}
\mathrm{SU}(2)/S(U_1 \times U_1) &\simeq \mathrm{Sp}(1)/U(1), \\
\mathrm{SO}(5)/\mathrm{SO}(3) \times \mathrm{SO}(2) &\simeq \mathrm{Sp}(2)/U(2), \\
\mathrm{SU}(4)/S(U(2) \times U(2)) &\simeq \mathrm{SO}(6)/U(3), \\
\mathrm{SO}(8)/\mathrm{SO}(4) \times \mathrm{SO}(4) &\simeq \mathrm{SO}(8)/U(4).
\end{aligned} \tag{4.64}$$

Notice the absence in (4.63) G_2 , F_4 and E_8 . Moreover we can introduce on those groups left-invariant skewsymmetric forms. Let us notice that in (4.62) we have the symmetric space $\mathrm{SU}(4)/S(U_3 \times U_1)$ which can have some physical applications. The same we can say for $\mathrm{SO}(10)/U(5)$ and for the last two symmetric spaces in (4.62).

We can introduce almost complex (almost Kählerian) structures on homogeneous spaces which are not symmetric spaces. They have the following shape: G/K , where K is a fixed point set of an automorphism of G of order 3. If $\dim(G/K) \geq 6$ these manifolds are not complex (Kählerian). For example we have the following possibilities:

1. $G = \mathrm{SU}(n)/Z_n$,
 $K = S(U(r_1) \times U(r_2) \times U(r_3))/Z_n$,
 $0 \leq r_1 \leq r_2 \leq r_3$, $r_1 + r_2 + r_3 = n$,
 $0 < r_2$ (4.65)
2. $G = \mathrm{SO}(2n+1)$,
 $K = U(r) \times \mathrm{SO}(2n-2r+1)$,
 $1 \leq r \leq n$.
3. $G = \mathrm{Sp}(n)/Z_2$,
 $K = \{U(r) \times \mathrm{Sp}(n-r)\}/Z_2$, $1 \leq r \leq n$.
4. $G = \mathrm{SO}(2n)/Z_2$, $n \geq 3$,
 $K = \{U(r) \times \mathrm{SO}(2n-2r)\}/Z_2$, $1 \leq r \leq n$.

Except the above example we have the following

$$\begin{aligned}
G &= G_2, & K &= U(2), \\
G &= F_4, & K &= (\mathrm{Spin}(7) \times T^1)/Z_2, \\
G &= F_4, & K &= (\mathrm{Sp}(3) \times T^1)/Z_2, \\
G &= E_6/Z_3, & K &= (\mathrm{SO}(10) \times \mathrm{SO}(2))/Z_2, \\
G &= E_6/Z_2, & K &= (S(U(5) \times U(1)) \times \mathrm{SU}(2))/Z_2, \\
G &= E_6/Z_2, & K &= (\mathrm{SU}(6)/Z_3 \times T^1)/Z_2, \\
G &= E_6/Z_2, & K &= [(\mathrm{SO}(8) \times \mathrm{SO}(2)) \times \mathrm{SO}(2)]/Z_2, \\
G &= E_7/Z_2, & K &= E_6 \times T^1/Z_3, \\
G &= E_7/Z_2, & K &= (\mathrm{SU}(2) \times (\mathrm{SO}(10) \times \mathrm{SO}(2)))/Z_2, \\
G &= E_7/Z_2, & K &= S(U(7) \times U(1))/Z_1, \\
G &= E_8, & K &= \mathrm{SO}(14) \times \mathrm{SO}(2),
\end{aligned} \tag{4.66}$$

$$\begin{aligned}
G = E8, & & K = (E7 \times T^1)/Z_2, \\
G = G2, & & K = \text{SU}(3), \\
G = F4, & & K = (\text{SU}(3) \times \text{SU}(3))/Z_3, \\
G = E6/Z_3, & & K = (\text{SU}(3) \times \text{SU}(3) \times \text{SU}(3))/(Z_2 \times Z_2), \\
G = E7/Z_3, & & K = (\text{SU}(3) \times (\text{SU}(6)/Z_2))/Z_3, \\
G = E8, & & K = (\text{SU}(3) \times E6)/Z_3, \\
G = E8, & & K = \text{SU}(9)/Z_3, \\
G = \text{Spin}(8), & & K = \text{SU}(3)/Z_3, \\
G = \text{Spin}(8), & & K = G2.
\end{aligned} \tag{4.66 cont.}$$

One can find all the details concerning a construction of an almost complex structure on the above homogeneous spaces in the fourth position of Ref. [86]. Due to the almost complex structure on G/K we can construct a left-invariant skewsymmetric form on G . Let us remind to the reader that in this case G/K is not a symmetric space.

In this way we can introduce some nonsymmetric metrics on:

$$\begin{aligned}
& \text{SO}(2n) - n, n \geq 3, \\
& \text{SO}(8) - 6, \\
& \text{Sp}(n) - n, \\
& \text{SO}(2n+1) - n,
\end{aligned}$$

$\text{SU}(n) - a$ number of partitions of n in such a way that $n = r_1 + r_2 + r_3$, $r_1 \leq r_2 \leq r_3$, $r_2 > 0$, $G2 - 2$, $F4 - 3$, $E6 - 5$, $E7 - 4$, $E8 - 4$ different Einstein–Kaufmann structures coming from almost Kählerian structures on (4.65–66).

All details concerning constructions presented here can be found in Refs. [7], [8], [9], [10], [16], [21]–[27], [51], [52], [64], [70], [71]–[77].

5. The Nonsymmetric Jordan–Thiry Theory over $V = E \times G/G_0$

Let P be the principal fibre bundle with the structural group H , over $V = E \times G/G_0$ with a projection π and let us define on this bundle a connection ω (Fig. 2A). Let us suppose that H is semisimple. On the base $V = E \times G/G_0$ we define a nonsymmetric metric tensor such that:

$$\begin{aligned}\gamma_{AB} &= \gamma_{(AB)} + \gamma_{[AB]}, \\ \gamma_{AB}\gamma^{CB} &= \gamma_{BA}\gamma^{BC} = \delta_A^C,\end{aligned}\tag{5.1}$$

where the order of the indices is important. We also define on V the connection $\bar{\omega}^A_B$,

$$\bar{\omega}^A_B = \bar{\Gamma}^A_{BC}\bar{\theta}^C\tag{5.2}$$

such that:

$$\bar{D}\gamma_{A+B-} = \bar{D}\gamma_{AB} - \gamma_{AD}\bar{Q}^D_{BC}(\bar{\Gamma})\theta^C = 0,\tag{5.3}$$

where $\bar{D}$ is the exterior covariant derivative with respect to $\bar{\omega}^A_B$ and $\bar{Q}^A_{BC}(\bar{\Gamma})$ is the torsion of $\bar{\omega}^A_B$. Using (4.39) one easily finds that the connection (5.2) has the following shape:

$$\bar{\omega}^A_B = \left(\begin{array}{c|c} \tilde{\omega}^\alpha_\beta & 0 \\ \hline 0 & \hat{\omega}^{\tilde{a}}_{\tilde{b}} \end{array} \right),\tag{5.4}$$

where $\tilde{\omega}^\alpha_\beta$ is the connection on the space-time E and $\hat{\omega}^{\tilde{a}}_{\tilde{b}}$ is the connection on the manifold $M = G/G_0$ with the following properties:

$$\begin{aligned}\tilde{D}g_{\alpha+\beta-} &= \tilde{D}g_{\alpha\beta} - g_{\alpha\delta}\tilde{Q}^\delta_{\beta\gamma}(\tilde{\Gamma})\theta^\gamma = 0, \\ \tilde{Q}^\alpha_{\beta\alpha}(\tilde{\Gamma}) &= 0,\end{aligned}\tag{5.5}$$

$$\hat{D}g_{\tilde{a}+\tilde{b}-} = \hat{D}g_{\tilde{a}\tilde{b}} - g_{\tilde{a}\tilde{d}}\hat{Q}^{\tilde{d}}_{\tilde{b}\tilde{c}}(\hat{\Gamma})\bar{\theta}^{\tilde{c}} = 0,\tag{5.6}$$

$\bar{D}$ is the exterior covariant derivative with respect to the connection $\bar{\omega}^\alpha_\beta$ on E , $\bar{Q}^\alpha_{\beta\gamma}(\bar{\omega})$ is the tensor of torsion for $\bar{\omega}^\alpha_\beta$. $\hat{D}$ means the exterior covariant derivative with respect to the connection $\hat{\omega}^{\tilde{a}}_{\tilde{b}}$ and $\hat{Q}^{\tilde{a}}_{\tilde{b}\tilde{c}}(\hat{\Gamma})$ is the tensor of torsion for $\hat{\omega}^{\tilde{a}}_{\tilde{b}}$. The second condition of Eq. (5.5) is not necessary to define Einstein connection on E . Moreover we suppose

it to be in line with some classical results. On the space-time E we also define the second affine connection $\overline{W}^\alpha{}_\beta$ such that:

$$\overline{W}^\alpha{}_\beta = \overline{\omega}^\alpha{}_\beta - \frac{2}{3}\delta^\alpha{}_\beta \overline{W}, \quad (5.7)$$

where

$$\overline{W} = \overline{W}_\gamma \overline{\theta}^\gamma = \frac{1}{2}(\overline{W}^\sigma{}_{\gamma\sigma} - \overline{W}^\sigma{}_{\sigma\gamma})\overline{\theta}^\gamma.$$

Thus on the space-time E we have all the geometrical quantities from N.G.T.: two connections $\overline{\omega}^\alpha{}_\beta$ and $\overline{W}^\alpha{}_\beta$ and the nonsymmetric metric $g_{\alpha\beta}$ (see [18], [34], [47], [65], [66]). Now let us turn to the nonsymmetric metrization of the bundle P . According to section 4 we have:

$$\varkappa_{\tilde{A}\tilde{B}} = \left(\begin{array}{c|c} \gamma_{AB} & 0 \\ \hline 0 & \rho^2 \ell_{ab} \end{array} \right), \quad (5.8)$$

where $\rho = \rho(x, y)$ is a scalar field on $E \times M$ and

$$\gamma_{AB} = \left(\begin{array}{c|c} g_{\alpha\beta} & 0 \\ \hline 0 & r^2 g_{\tilde{a}\tilde{b}} \end{array} \right) \quad \text{and} \\ \ell_{ab} = h_{ab} + \xi k_{ab}, \quad \det(\ell_{ab}) \neq 0$$

or

$$\begin{aligned} \varkappa_{(\tilde{A}\tilde{B})} \theta^{\tilde{A}} \otimes \theta^{\tilde{B}} &= \gamma_{(AB)} \theta^A \otimes \theta^B + \rho^2 h_{ab} \theta^a \otimes \theta^b, \\ \varkappa_{[\tilde{A}\tilde{B}]} \theta^{\tilde{A}} \wedge \theta^{\tilde{B}} &= \gamma_{[AB]} \theta^A \wedge \theta^B + \rho^2 \xi k_{ab} \theta^a \wedge \theta^b, \end{aligned} \quad (5.9)$$

where $\theta^a = \lambda \omega^a$. From the Kaluza–Klein Theory and Jordan–Thiry we know that λ is proportional to $\sqrt{G_N}$ (see [1], [2], [7], [8], [9], [10], [18]), $\lambda \sim \sqrt{G_N}$. We work with such a system of units that $\lambda = 2$.

Now we define on P , a connection $\omega^{\tilde{A}}{}_{\tilde{B}}$ right-invariant (Ad-covariant) with respect to the action of group H on P such that:

$$D\gamma_{\tilde{A}+\tilde{B}-} = D\gamma_{\tilde{A}\tilde{B}} - \gamma_{\tilde{A}\tilde{D}} Q^{\tilde{D}}{}_{\tilde{B}\tilde{C}}(\Gamma) \theta^{\tilde{C}} = 0, \quad (5.10)$$

where $\omega^{\tilde{A}}{}_{\tilde{B}} = \Gamma^{\tilde{A}}{}_{\tilde{B}\tilde{C}} \theta^{\tilde{C}}$. D is the exterior covariant derivative with respect to the connection $\omega^{\tilde{A}}{}_{\tilde{B}}$ and $Q^{\tilde{A}}{}_{\tilde{B}\tilde{C}}(\Gamma)$ is the tensor of torsion for the connection $\omega^{\tilde{A}}{}_{\tilde{B}}$. After some calculations one finds:

$$\omega^{\tilde{A}}{}_{\tilde{B}} = \left(\begin{array}{c|c} \pi^*(\overline{\omega}^A{}_B) - \rho^2 \ell_{db} \gamma^{MA} L^d{}_{MB} \theta^b & L^A{}_{BC} \theta^C - \frac{1}{\rho} \gamma_{BD} \tilde{\gamma}^{(DC)} \rho_{,C} \theta^a \\ \hline \rho^2 \ell_{bd} \gamma^{AB} (2H^d{}_{CB} - L^d{}_{CB}) \theta^C - \rho \tilde{\gamma}^{(AB)} \rho_{,B} \ell_{bc} \theta^C & \frac{1}{\rho} \gamma_{BD} \tilde{\gamma}^{(DC)} \rho_{,C} \delta_b^a \theta^B + \tilde{\omega}^a{}_b \end{array} \right), \quad (5.11)$$

where $\tilde{\gamma}^{(AB)}$ is the inverse tensor for $\gamma_{(AB)}$

$$\begin{aligned}\gamma_{(AB)}\tilde{\gamma}^{(AC)} &= \delta^C_B, \\ L^d_{MB} &= -L^d_{BM},\end{aligned}\tag{5.12}$$

is an Ad-type tensor (Ad-covariant) on P such that

$$\begin{aligned}\ell_{dc}\gamma_{MB}\gamma^{CM}L^d_{CA} + \ell_{cd}\gamma_{AM}\gamma^{MC}L^d_{BC} &= 2\ell_{cd}\gamma_{AM}\gamma^{MC}H^d_{BC}, \\ \tilde{\omega}^a_b &= \tilde{\Gamma}^a_{bc}\theta^c,\end{aligned}\tag{5.13}$$

$$\begin{aligned}\ell_{db}\tilde{\Gamma}^d_{ac} + \ell_{ad}\tilde{\Gamma}^d_{cb} &= -\ell_{db}C^d_{ac}, \\ \tilde{\Gamma}^d_{ac} &= -\tilde{\Gamma}^d_{ca}, \quad \tilde{\Gamma}^d_{ad} = 0.\end{aligned}\tag{5.14}$$

We define on P a second connection:

$$W^{\tilde{A}}_{\tilde{B}} = \omega^{\tilde{A}}_{\tilde{B}} - \frac{4}{3(m+2)}\delta^{\tilde{A}}_{\tilde{B}}\overline{W}.\tag{5.15}$$

Thus we have on P all $(m+4)$ -dimensional analogues of geometrical quantities from N.G.T., i.e.:

$$W^{\tilde{A}}_{\tilde{B}}, \omega^{\tilde{A}}_{\tilde{B}} \quad \text{and} \quad \varkappa_{\tilde{A}\tilde{B}}.$$

Formulae (5.11)–(5.13) are similar to the analogous formulae from Refs. [23], [24] (see also [18]). The one difference is that now we have $E \times M$ in place of E . The connection (5.15) is analogous to the connection W from Refs. [19], [23] and from Part 4. The form $\overline{W}$ is horizontal one, $\text{hor } \overline{W} = \overline{W}$ (in the sense of the connection ω on the bundle P).

Let us calculate the Moffat-Ricci curvature scalar for the connection $W^{\tilde{A}}_{\tilde{B}}$.

$$R(W) = \varkappa^{\tilde{A}\tilde{B}} \left(R^{\tilde{C}}_{\tilde{A}\tilde{B}\tilde{C}}(W) + \frac{1}{2}R^{\tilde{C}}_{\tilde{C}\tilde{A}\tilde{B}}(W) \right),\tag{5.16}$$

where $R^{\tilde{A}}_{\tilde{B}\tilde{C}\tilde{D}}(W)$ is the curvature tensor for the connection $W^{\tilde{A}}_{\tilde{B}}$. Using results from Refs. [19], [23], [24] one gets:

$$\begin{aligned}R(W) &= \overline{R}(\overline{W}) + \frac{1}{r^2}R(\widehat{\Gamma}) + \frac{1}{\lambda^2\rho^2}\tilde{R}(\tilde{\Gamma}) + \\ &\quad - \frac{\lambda^2\rho^2}{4}\ell_{ab}(2H^aH^b - L^{aMN}H^b_{MN}) - \frac{2\lambda^2\tilde{M}}{\rho^2}\tilde{\gamma}^{(BN)}\rho_{,B}\rho_{,N} + \underline{P}(\rho),\end{aligned}\tag{5.17}$$

where:

$$\begin{aligned} \underline{P}(\rho) = & \frac{\lambda^2 n}{8\rho} \gamma_{\text{DB}} \tilde{\gamma}^{(\text{DC})} \rho_{,\text{C}} \gamma^{\text{MN}} \overline{Q}^{\text{B}}_{\text{MN}}(\overline{\omega}) + \\ & + \frac{\lambda^2 n}{4\rho^2} \overline{\nabla}_{\text{A}}(\rho \tilde{\gamma}^{(\text{AB})} \rho_{,\text{B}}) + \frac{\lambda^2}{4} n \gamma^{\text{BC}} \overline{\nabla}_{\text{C}} \left(\frac{1}{\rho} \gamma_{\text{BD}} \tilde{\gamma}^{(\text{DE})} \rho_{,\text{E}} \right) + \\ & + \frac{\lambda^2 \rho}{8} \gamma^{\text{MN}} \left\{ \overline{\nabla}_{\text{M}} \left(\frac{1}{\rho} \gamma_{\text{DN}} \tilde{\gamma}^{(\text{DC})} \rho_{,\text{C}} \right) - \overline{\nabla}_{\text{N}} \left(\frac{1}{\rho} \gamma_{\text{DM}} \tilde{\gamma}^{(\text{DC})} \rho_{,\text{C}} \right) \right\}, \end{aligned} \quad (5.18)$$

$$\tilde{\text{M}} = \ell^{[\text{dc}]} \ell_{[\text{dc}]} - n(n-1), \quad (5.19)$$

$\overline{Q}^{\text{B}}_{\text{MN}}(\overline{\omega})$ means the torsion of the connection $\overline{\omega}^{\text{A}}_{\text{B}}$ on $V = E \times M = E \times G/G_0$ and $\overline{\nabla}_{\text{A}}$ is the covariant derivative with respect to this connection. $\overline{R}(\overline{W})$ is the Moffat–Ricci curvature scalar on the space-time E for the connection $\overline{W}^{\alpha}_{\beta}$. $R(\widehat{\Gamma})$ is the Moffat–Ricci curvature scalar for the connection $\widehat{\omega}^{\alpha}_{\beta}$ on the homogeneous space $M = G/G_0$, $\widetilde{R}(\widetilde{\Gamma})$ is the Moffat–Ricci curvature scalar for the connection $\widetilde{\omega}^a_b$. Notice that in the curvature scalar we pass from $\lambda = 2$ to the arbitrary value of this constant.

$$H^a = \gamma^{[\text{AB}]} H^a_{[\text{AB}]} = g^{[\alpha\beta]} H^a_{\alpha\beta} + \frac{1}{r^2} g^{[\tilde{a}\tilde{b}]} H^a_{\tilde{a}\tilde{b}}, \quad (5.20)$$

$$\begin{aligned} L^{a\text{MN}} = & \gamma^{\text{AM}} \gamma^{\text{BN}} L^a_{\text{AB}} = \delta^{\text{M}}_{\mu} \delta^{\text{N}}_{\gamma} g^{\alpha\mu} g^{\beta\gamma} L^a_{\alpha\beta} + \\ & + \frac{1}{r^2} (g^{\alpha\mu} g^{\tilde{b}\tilde{n}} L^a_{\alpha\tilde{b}} + g^{\tilde{a}\tilde{n}} g^{\beta\gamma} L^a_{\tilde{a}\beta}) \delta^{\text{M}}_{\mu} \delta^{\text{N}}_{\tilde{n}} + \\ & + \frac{1}{r^4} g^{\tilde{a}\tilde{m}} g^{\tilde{b}\tilde{n}} L^a_{\tilde{a}\tilde{b}} \delta^{\text{M}}_{\tilde{m}} \delta^{\text{N}}_{\tilde{n}}. \end{aligned} \quad (5.21)$$

Let us consider the condition (5.13). One gets:

$$\ell_{\text{dc}} g_{\mu\beta} g^{\gamma\mu} L^d_{\gamma\alpha} + \ell_{\text{cd}} g_{\alpha\mu} g^{\mu\gamma} L^d_{\beta\gamma} = 2\ell_{\text{cd}} g_{\alpha\mu} g^{\mu\gamma} H^d_{\beta\gamma}, \quad (5.22)$$

$$\ell_{\text{dc}} g_{\tilde{m}\tilde{b}} g^{\tilde{c}\tilde{m}} L^d_{\tilde{c}\tilde{a}} + \ell_{\text{cd}} g_{\tilde{a}\tilde{m}} g^{\tilde{m}\tilde{c}} L^d_{\tilde{b}\tilde{c}} = 2\ell_{\text{cd}} g_{\tilde{a}\tilde{m}} g^{\tilde{m}\tilde{c}} H^d_{\tilde{b}\tilde{c}}, \quad (5.23)$$

$$\ell_{\text{dc}} g_{\mu\beta} g^{\gamma\mu} L^d_{\gamma\tilde{a}} + \ell_{\text{cd}} g_{\tilde{a}\tilde{m}} g^{\tilde{m}\tilde{c}} L^d_{\beta\tilde{c}} = 2\ell_{\text{cd}} g_{\tilde{a}\tilde{m}} g^{\tilde{m}\tilde{c}} H^d_{\beta\tilde{c}}. \quad (5.24)$$

One finds that:

$$\begin{aligned} -\ell_{ab} L^{a\text{MN}} H^b_{\text{MN}} = & -\ell_{ab} \left(g^{\alpha\mu} g^{\beta\nu} L^a_{\alpha\beta} H^b_{\mu\nu} + \right. \\ & + \frac{2}{r^2} g^{\alpha\mu} g^{\tilde{b}\tilde{n}} L^a_{\alpha\tilde{b}} H^b_{\mu\tilde{n}} + \frac{1}{r^4} g^{\tilde{a}\tilde{m}} g^{\tilde{b}\tilde{n}} L^a_{\tilde{a}\tilde{b}} H^b_{\tilde{m}\tilde{n}} \Big) = \\ = & -\ell_{ab} \left(L^{a\mu\nu} H^b_{\mu\nu} + \frac{2}{r^2} g^{\tilde{b}\tilde{n}} L^{\alpha\mu}_{\tilde{b}} H^b_{\mu\tilde{n}} + \right. \\ & + \frac{1}{r^4} g^{\tilde{a}\tilde{m}} g^{\tilde{b}\tilde{n}} L^a_{\tilde{a}\tilde{b}} H^b_{\tilde{m}\tilde{n}} \Big), \end{aligned} \quad (5.25)$$

where:

$$\begin{aligned} L^{\alpha\mu\nu} &= g^{\alpha\mu} g^{\beta\nu} L^a_{\alpha\beta}, \\ L^{\alpha\mu}_{\tilde{b}} &= g^{\alpha\mu} L^a_{\alpha\tilde{b}}. \end{aligned}$$

For $\ell_{ab}H^aH^b = h_{ab}H^aH^b$, we have the following:

$$h_{ab}H^aH^b = h_{ab}H^a_0H^b_0 + \frac{2}{r^2}h_{ab}H^a_0H^b_1 + \frac{1}{r^4}h_{ab}H^a_1H^b_1, \quad (5.26)$$

where

$$H^a_0 = g^{[\alpha\beta]}H^a_{\alpha\beta} \quad (5.27)$$

and

$$H^a_1 = g^{[\tilde{a}\tilde{b}]}H^a_{\tilde{a}\tilde{b}}. \quad (5.28)$$

And finally we get for $R(W)$:

$$\begin{aligned} R(W) &= \overline{R}(\overline{W}) + \frac{1}{r^2}R(\widehat{\Gamma}) + \frac{4}{\lambda^2 r^2}\widetilde{R}(\widetilde{\Gamma}) - \frac{\lambda^2 \rho^2}{4}\ell_{ab}(2H^a_0H^b_0 - L^{a\mu\nu}H^b_{\mu\nu}) + \\ &\quad - \frac{\lambda^2 \rho^2}{4r^2}\ell_{ab}(4H^{(a}_0H^{b)}_1 - 2g^{\tilde{b}\tilde{n}}L^a_{\tilde{b}}H^b_{\mu\tilde{n}}) + \\ &\quad - \frac{\lambda^2 \rho^2}{4r^4}\ell_{ab}(2H^a_1H^b_1 - g^{\tilde{a}\tilde{m}}g^{\tilde{b}\tilde{n}}L^a_{\tilde{a}\tilde{b}}H^b_{\tilde{m}\tilde{n}} + \\ &\quad - \frac{\widetilde{M}}{\rho^2}g_{\gamma\delta}\widetilde{g}^{(\delta\nu)}\rho_{,\nu}\widetilde{g}^{(\gamma\beta)}\rho_{,\beta} + \underline{P}(\rho), \end{aligned} \quad (5.29)$$

where $\widetilde{g}^{(\delta\nu)}$ is the inverse tensor of $g_{(\alpha\beta)}$ such that:

$$\widetilde{g}^{(\delta\nu)}g_{(\delta\mu)} = \delta^\nu_\mu. \quad (5.30)$$

Let us calculate the density for the Moffat–Ricci curvature scalar for the connection $W^{\tilde{A}}_{\tilde{B}}$. We have:

$$\sqrt{|\varkappa|}R(W) = \sqrt{-g}r^{n_1}\sqrt{|\widetilde{g}|}\sqrt{|\ell|}\rho^n R(W), \quad (5.31)$$

where

$$g = \det(g_{\alpha\beta}), \quad \widetilde{g} = \det(g_{\tilde{a}\tilde{b}}), \quad \ell = \det(\ell_{ab}) \quad (5.32)$$

and

$$\varkappa = \det(\varkappa_{\tilde{A}\tilde{B}}) = g \cdot r^{n_1} \cdot \widetilde{g} \cdot \rho^n \ell. \quad (5.33)$$

After some calculation one gets:

$$\begin{aligned}
\sqrt{|\varkappa|}R(W) = & \sqrt{-gr^{n_1}}\sqrt{|\tilde{g}|}\ell\left\{\rho^n\overline{R}(\overline{W}) + \frac{\tilde{R}(\tilde{\Gamma})}{\lambda^2\rho^{2-n}} + \frac{\rho^n}{r^2}R(\widehat{\Gamma}) + \right. \\
& + \frac{\lambda^2}{4}\rho^{n+2}\ell_{ab}(2H^a{}_0H^b{}_0 - L^{a\mu\nu}H^b{}_{\mu\nu}) + \\
& + \frac{\lambda^2}{4r^2}\rho^{n+2}\ell_{ab}(4H_0^{(a}H_1^{b)} - 2g^{\tilde{b}\tilde{n}}L^{a\mu}{}_{\tilde{b}}H^b{}_{\mu\tilde{n}}) + \\
& + \frac{\lambda^2}{4r^4}\rho^{n+2}\ell_{ab}(2H_1^aH_1^b - g^{\tilde{a}\tilde{m}}g^{\tilde{b}\tilde{n}}L^a{}_{\tilde{a}\tilde{b}}H^b{}_{\tilde{m}\tilde{n}}) + \\
& + \frac{\lambda^2}{4}\rho^{n-2}(\overline{M}\tilde{g}^{(\gamma\mu)}\rho_{,\gamma}\rho_{,\mu} + \\
& \left. + n^2g^{[\mu\nu]}g_{\delta\mu}\tilde{g}^{(\delta\gamma)}\rho_{,\nu}\rho_{,\gamma})\right\} + \partial_M K^M, \tag{5.34}
\end{aligned}$$

where

$$\overline{M} = (\ell^{[dc]}\ell_{[dc]} - 3n(n-1)) \tag{5.35}$$

and

$$K^M = \frac{n}{2}\rho^{n-1}\sqrt{|\varkappa|}(5\tilde{\gamma}^{(MC)} - \gamma^{NM}\gamma_{DN}\tilde{\gamma}^{(DC)}) \cdot \rho_{,C}. \tag{5.35'}$$

If we define an integral of action:

$$S \sim \int_U \sqrt{|\varkappa|}R(W)d^{m+4}x, \tag{5.36}$$

(where $d^{m+4}x = d^4x d\mu_H(h) \cdot dm(y)$, $d\mu_H(h)$ is a biinvariant measure on a group H and $dm(y)$ is a measure on M induced by a biinvariant measure on a group G , $x \in E$, $h \in H$, $y \in M$), and the variation principle for $R(W)$ then the full divergence $\partial_M K^M$ does not play any role. It could play a certain role in topological problems.

One can write

$$K^\mu = \frac{n}{2}\rho^{n-1}\sqrt{g\tilde{g}}(5\tilde{g}^{(\mu\gamma)} - g^{\nu\mu}g_{\delta\nu}\tilde{g}^{(\delta\gamma)})\rho_{,\gamma}$$

and

$$K^{\tilde{m}} = \frac{n}{2r^2}\rho^{n-1}\sqrt{g\tilde{g}}(5g^{(\tilde{m}\tilde{c})} - g^{\tilde{n}\tilde{m}}g_{\tilde{d}\tilde{n}}\tilde{g}^{(\tilde{d}\tilde{c})})\rho_{,\tilde{c}}.$$

Thus we really only have to deal with $B(W)$:

$$B(W) = \sqrt{\varkappa}R(W) - \partial_M K^M. \tag{5.37}$$

In the calculations we use results from Refs. [23], [24], [26], [27] and we suppose that $\rho = \rho(x)$ is really the scalar field on the manifold E (space-time). The latest means that:

$$\rho_{,\tilde{a}} = 0, \tag{5.38}$$

for every $\tilde{a} = 5, 6, 7, \dots, n_1 + 4$.

Moreover it is interesting to consider a more general case with ρ depending on $x \in E$ and $y \in M$ i.e.

$$\rho = \rho(x, y). \quad (5.39)$$

In this case we expand ρ into a complete set of real functions defined on M .

$$\rho = \sum_{k=0}^{\infty} \rho_k(x) \chi_k(y) \quad (5.40)$$

and consider such a ρ that:

$$\|\tilde{\rho}(x)\|^2 = \sum_{k=0}^{\infty} \rho_k^2(x) < \infty. \quad (5.41)$$

The condition (5.41) means that $\tilde{\rho}(x) \in \ell^2$ (a Hilbert space) $x \in E$ i.e. ℓ^2 -valued function defined on E . If functions $\chi_k(y)$ defined on M form an orthogonal basis in $L^2(M, dm, \sqrt{|\tilde{g}|})$ i.e.

$$\|\chi_k\|^2 = \frac{1}{V_2} \int_M \sqrt{|\tilde{g}|} \chi_k^2(y) dm(y) < \infty \quad (5.42)$$

and

$$(\chi_k, \chi_l) = \frac{1}{V_2} \int_M \sqrt{|\tilde{g}|} \chi_k(y) \chi_l(y) dm(y) = \|\chi_k\|^2 \delta_{kl} \quad k, l = 0, 1, 2 \dots \quad (5.43)$$

and for every real $f(y)$ such that

$$\|f\|^2 = \frac{1}{V_2} \int_M \sqrt{|\tilde{g}|} f^2(y) dm(y) < \infty, \quad (5.44)$$

$$f(y) = \sum_{k=0}^{\infty} f_k \chi_k(y), \quad (5.45)$$

$$\sum_{k=0}^{\infty} f_k^2 < \infty, \quad (5.46)$$

$$f_k = \frac{1}{V_2} \int_M \sqrt{|\tilde{g}|} f(y) \chi_k(y) dm(y). \quad (5.47)$$

The convergence of (5.40), (5.45) is understood in the sense of L^2 norm. Thus in this more general case we have to do with a tower of scalar fields $\rho_k(x)$, $k = 0, 1, 2, 3 \dots$

We can arrange a basis $\chi_k(y)$ in such way that:

$$\chi_0(y) = 1. \quad (5.48)$$

It is easy to see that this function is normalized to one:

$$\|\chi_0\| = 1. \quad (5.49)$$

Using a Schmidt orthogonalization procedure one gets remaining χ_k , $k = 1, 2 \dots$ from generalized harmonics on M such that:

$$(\chi_k, \chi_l) = \delta_{kl}. \quad (5.50)$$

In this way the condition (5.38) means a truncation condition for ρ i.e.

$$\rho(x) = \rho_0(x) + \sum_{k=1}^{\infty} \rho_k(x) \chi_k(y) \quad (5.51)$$

and we consider only the first term ρ_0

$$\rho(x) = \rho_0(x). \quad (5.52)$$

The general treatment of ρ is given in section 16.

The generalized harmonics on (M, h^0) are eigenfunctions of the usual Beltrami–Laplace operator on (M, h^0) (which is an elliptic operator for M is compact and without boundaries).

$$\Delta \eta_k = a_k \eta_k, \quad (5.53)$$

where η_k is an eigenfunction of Δ and a_k is an eigenvalue, $k = 0, 1, 2 \dots$

$$u\Delta = \frac{1}{2}(d * d * + * d * d) \quad (5.54)$$

is the Beltrami–Laplace operator on M , $*$ —means a Hodge’s star and d is an exterior derivative on M , u means a volume form on M (n_1 -form). Sometimes we define $\delta = *d*$. One can write Δ in a more convenient form:

$$\Delta f = \text{div}(\text{grad}(f)) = -\frac{1}{\sqrt{|h^0|}}(h^{0\tilde{a}\tilde{b}}\sqrt{|h^0|}f_{,\tilde{b}})_{,\tilde{a}}, \quad (5.54a)$$

where $h^0 = \det(h_{\tilde{a}\tilde{b}}^0)$.

It is easy to see that $a_k = 0$ corresponds to a constant function (a manifold M is compact).

Functions η_k form a basis in a different Hilbert space of L^2 -type i.e. $\eta_k \in L^2(M, dm)$. Moreover $L^2(M, dm)$ and $L^2(M, dm, \sqrt{|\tilde{g}|})$ are unitary equivalent since measures $d\mu_1 = dm$ and $d\mu_2 = \sqrt{|\tilde{g}|} dm$ are such that $\mu_2 \ll \mu_1$ and $\mu_1 \ll \mu_2$. The isomorphism

$$\begin{aligned} \mathbf{U} : L^2(M, dm, \sqrt{|\tilde{g}|}) &\rightarrow L^2(M, dm), \\ f_2 = \mathbf{U}(f_1) &= |\tilde{g}|^{1/4} f_1 \end{aligned}$$

establishes this equivalence.

The truncation procedure can be obtained in a different way taking

$$\rho(x, y) = \rho_0(x) \rho_1(y), \quad x \in E, \quad y \in M.$$

In this way after averaging over the manifold M we get basically the same shape of the lagrangian with exactly the same factors depending on the scalar field $\rho_0(x)$.

It is worth to notice that the formulae (5.21)–(5.23) do not change under the general conformal transformation—the redefinition of the tensor $g_{\mu\nu}$. This is a general property of these formulae i.e.

$$g_{\mu\nu} \rightarrow C \cdot g_{\mu\nu}, \quad (5.55)$$

where $C = C(x)$ is the conformal factor.

$R(W)$ is invariant with respect to the right action of the group H on $\underline{P}$. Thus an integration over the group H is trivial. Let us consider the following two Ad_H -type 2-forms with values in the Lie algebra of H , $(\mathfrak{h})$.

$$\bar{L} = \frac{1}{2} L^d_{\text{MB}} \theta^{\text{M}} \wedge \theta^{\text{B}} X_d \quad (5.56)$$

and

$$\bar{Q} = \frac{1}{2} Q^d_{\text{MB}} \theta^{\text{M}} \wedge \theta^{\text{B}} X_d. \quad (5.57)$$

One gets

$$\bar{L} = \Omega - \frac{1}{2} \bar{Q}, \quad (5.58)$$

where Ω is the curvature of the connection ω on P (over $E \times G/G_0$) and Q^d_{MB} is torsion in additional gauge dimensions. This generalizes our considerations in Nonabelian Kaluza-Klein Theory to higher dimensional space-time.

6. Dimensional Reduction Procedure

Let us consider equation (5.21), (5.23) and (5.24) *modulo* equation (4.37), (4.38) and (4.39). One finds:

$$\ell_{ij}g_{\mu\beta}g^{\gamma\mu}\tilde{L}^i_{\gamma\alpha} + \ell_{ji}g_{\alpha\mu}g^{\mu\gamma}\tilde{L}^i_{\beta\gamma} = 2\ell_{ji}g_{\alpha\mu}g^{\mu\gamma}\tilde{H}^i_{\beta\gamma}, \quad (6.1)$$

where $\ell_{ij} = \ell_{cd}\alpha^c_i\alpha^d_j$ is a right-invariant nonsymmetric metric on the group G and

$$L^c_{\mu\nu} = \alpha^c_i\tilde{L}^i_{\mu\nu}, \quad (6.2)$$

$\tilde{L}^i_{\mu\nu}$ plays the role of the second tensor of strength (an induction tensor) for the Yang–Mills' field with the gauge group G . $\tilde{H}^i_{\mu\nu}$ is of course the first tensor of strength of this field. The polarization tensor is defined as usual (see [18], [19], [23], [24], [26], [27]).

$$\tilde{L}^i_{\mu\nu} = \tilde{H}^i_{\mu\nu} - 4\pi\tilde{M}^i_{\mu\nu}. \quad (6.3)$$

We introduce two Ad_G -type 2-forms with values in the Lie algebra $\mathfrak{g}$ (of G)

$$\begin{aligned} \tilde{L} &= \frac{1}{2}\tilde{L}^i_{\mu\nu}\theta^\mu \wedge \theta^\nu Y_i, \\ \tilde{M} &= \frac{1}{2}\tilde{M}^i_{\mu\nu}\theta^\mu \wedge \theta^\nu Y_i \end{aligned}$$

and one easily writes

$$\tilde{L} = \tilde{\Omega}_E - 4\pi\tilde{M} = \tilde{\Omega}_E - \frac{1}{2}Q,$$

where $\tilde{Q} = \frac{1}{2}\tilde{Q}^i_{\mu\nu}\theta^\mu \wedge \theta^\nu Y_i$.

In this way we get a geometrical interpretation of a Yang–Mills' induction tensor in terms of the curvature and torsion in additional dimensions (see Ref. [18]).

From (5.24) and (4.38) one gets:

$$\ell_{dc}g_{\mu\beta}g^{\gamma\mu}L^d_{\gamma\tilde{a}} + \ell_{cd}g_{\tilde{a}\tilde{m}}g^{\tilde{m}\tilde{c}}L^d_{\beta\tilde{c}} = 2\ell_{cd}g_{\tilde{a}\tilde{m}}g^{\tilde{m}\tilde{c}}\nabla^{\text{gauge}}_{\beta}(\Phi^d_{\tilde{c}}). \quad (6.4)$$

From (5.22) and (4.39) one gets:

$$\ell_{dc}g_{\tilde{m}\tilde{b}}g^{\tilde{c}\tilde{m}}L^d_{\tilde{c}\tilde{a}} + \ell_{cd}g_{\tilde{a}\tilde{m}}g^{\tilde{m}\tilde{c}}L^d_{\tilde{b}\tilde{c}} = 2\ell_{cd}g_{\tilde{a}\tilde{m}}g^{\tilde{m}\tilde{c}}(C^d_{ab}\Phi^a_{\tilde{b}}\Phi^b_{\tilde{c}} - \mu^d_i f^i_{\tilde{b}\tilde{c}} - \Phi^d_{\tilde{a}}f^d_{\tilde{b}\tilde{c}}), \quad (6.5)$$

One can rewrite the formula (6.5) in the following shape

$$\ell_{dc}g_{\tilde{m}\tilde{b}}g^{\tilde{c}\tilde{m}}L^d_{\tilde{c}\tilde{a}}(1) + \ell_{cd}g_{\tilde{a}\tilde{m}}g^{\tilde{m}\tilde{c}}L^d_{\tilde{b}\tilde{c}}(1) = 2\ell_{cd}g_{\tilde{a}\tilde{m}}g^{\tilde{m}\tilde{c}}C^d_{ab}\Phi^a_{\tilde{b}}\Phi^b_{\tilde{c}}, \quad (6.5a)$$

$$\ell_{dc}g_{\tilde{m}\tilde{b}}g^{\tilde{c}\tilde{m}}L^d_{\tilde{c}\tilde{a}}(2) + \ell_{cd}g_{\tilde{a}\tilde{m}}g^{\tilde{m}\tilde{c}}L^d_{\tilde{b}\tilde{c}}(2) = 2\ell_{cd}g_{\tilde{a}\tilde{m}}g^{\tilde{m}\tilde{c}}\mu^d_{\hat{i}}f^{\hat{i}}_{\tilde{b}\tilde{c}}, \quad (6.5b)$$

$$\ell_{dc}g_{\tilde{m}\tilde{b}}g^{\tilde{c}\tilde{m}}L^d_{\tilde{c}\tilde{a}}(3) + \ell_{cd}g_{\tilde{a}\tilde{m}}g^{\tilde{m}\tilde{c}}L^d_{\tilde{b}\tilde{c}}(3) = 2\ell_{cd}g_{\tilde{a}\tilde{m}}g^{\tilde{m}\tilde{c}}f^{\tilde{d}}_{\tilde{b}\tilde{c}}\Phi^d_{\tilde{d}} \quad (6.5c)$$

and

$$L^d_{\tilde{a}\tilde{b}} = L^d_{\tilde{a}\tilde{b}}(1) - L^d_{\tilde{a}\tilde{b}}(2) - L^d_{\tilde{a}\tilde{b}}(3). \quad (6.5*)$$

Moreover we can write $L(i)$, $i = 1, 2, 3$ in the following forms

$$L^d_{\tilde{a}\tilde{b}}(1) = C^d_{be}\Psi^b_{\tilde{a}}\Psi^e_{\tilde{b}}, \quad (6.5a*)$$

$$L^d_{\tilde{a}\tilde{b}}(2) = \mu^d_{\hat{i}} \cdot m^{\hat{i}}_{\tilde{b}\tilde{c}}, \quad (6.5b*)$$

$$L^d_{\tilde{a}\tilde{b}}(3) = r^{\tilde{d}}_{\tilde{a}\tilde{b}}\Phi^d_{\tilde{d}}, \quad (6.5c*)$$

where $\Psi^b_{\tilde{a}}(x)$ is a multiple of scalar fields transforming with respect to groups G and H as $\Phi^b_{\tilde{a}}(x)$, $m^{\hat{i}}_{\tilde{b}\tilde{c}}$ are left-invariant skewsymmetric 2-forms on M with values in the Lie algebra $\mathfrak{g}_0$, $r^{\tilde{d}}_{\tilde{a}\tilde{b}}$ is a left-invariant tensor field on M . One gets

$$\ell_{dc}g_{\tilde{m}\tilde{b}}g^{\tilde{c}\tilde{m}}\Psi^b_{\tilde{c}}\Psi^e_{\tilde{a}} + \ell_{cd}g_{\tilde{a}\tilde{m}}g^{\tilde{m}\tilde{c}}\Psi^b_{\tilde{b}}\Psi^e_{\tilde{c}} = 2\ell_{cd}g_{\tilde{a}\tilde{m}}g^{\tilde{m}\tilde{c}}\Phi^b_{\tilde{b}}\Phi^e_{\tilde{c}}, \quad (6.5a**)$$

$$\ell_{dc}g_{\tilde{m}\tilde{b}}g^{\tilde{c}\tilde{m}}m^{\hat{i}}_{\tilde{c}\tilde{a}} + \ell_{cd}g_{\tilde{a}\tilde{m}}g^{\tilde{m}\tilde{c}}m^{\hat{i}}_{\tilde{b}\tilde{c}} = 2\ell_{cd}g_{\tilde{a}\tilde{m}}g^{\tilde{m}\tilde{c}}f^{\hat{i}}_{\tilde{b}\tilde{c}}, \quad (6.5b**)$$

$$\ell_{dc}g_{\tilde{m}\tilde{b}}g^{\tilde{c}\tilde{m}}r^{\tilde{d}}_{\tilde{c}\tilde{a}} + \ell_{cd}g_{\tilde{a}\tilde{m}}g^{\tilde{m}\tilde{c}}r^{\tilde{d}}_{\tilde{b}\tilde{c}} = 2\ell_{cd}g_{\tilde{a}\tilde{m}}g^{\tilde{m}\tilde{c}}f^{\tilde{d}}_{\tilde{a}\tilde{b}} \quad (6.5c**)$$

and

$$L^d_{\tilde{a}\tilde{b}} = C^d_{be}\Psi^b_{\tilde{a}}\Psi^e_{\tilde{b}} - \mu^d_{\hat{i}} \cdot m^{\hat{i}}_{\tilde{a}\tilde{b}} + r^{\tilde{d}}_{\tilde{a}\tilde{b}}\Phi^d_{\tilde{d}}. \quad (6.5***)$$

It seems that sometimes the last formula will be easier to apply in order to find an explicit form of the Higgs' potential.

Now we have the following formula for $B(W)$:

$$\begin{aligned} B(W) = & \sqrt{-g|\tilde{g}||\ell|} \left\{ r^{n_1} \rho^n \overline{R}(\overline{W}) + \frac{\rho^{n-2}}{\lambda^2} r^{n_1} \tilde{R}(\tilde{W}) + \right. \\ & + r^{n_1-2} \rho^n \widehat{R}(\widehat{W}) - r^{n_1} \rho^{n+2} \frac{\lambda^2}{4} \tilde{\ell}_{ij} (2\tilde{H}^i \tilde{H}^j - \tilde{L}^{i\mu\nu} \tilde{H}^j_{\mu\nu}) + \\ & + \frac{\lambda^2 \rho^{n+2} r^{n_1-2}}{2} \ell_{ab} g^{\tilde{b}\tilde{n}} L^{a\mu}_{\tilde{b}} \nabla_{\mu}^{\text{gauge}} \Phi^b_{\tilde{n}} + \\ & - \frac{\lambda^2 \rho^{n+2} r^{n_1-4}}{4} \ell_{ab} [2g^{[\tilde{a}\tilde{b}]} (C^a_{cd} \Phi^c_{\tilde{a}} \Phi^d_{\tilde{b}} - \mu^a_{\hat{i}} f^{\hat{i}}_{\tilde{a}\tilde{b}} - \Phi^a_{\tilde{d}} f^{\tilde{d}}_{\tilde{a}\tilde{b}}) \times \\ & \times g^{[\tilde{n}\tilde{m}]} (C^b_{ef} \Phi^e_{\tilde{n}} \Phi^f_{\tilde{m}} - \mu^b_{\hat{i}} f^{\hat{i}}_{\tilde{n}\tilde{m}} - \Phi^b_{\tilde{e}} f^{\tilde{e}}_{\tilde{n}\tilde{m}}) + \end{aligned} \quad (6.6)$$

$$\begin{aligned}
& -g^{\tilde{a}\tilde{m}}g^{\tilde{b}\tilde{n}}L^a_{\tilde{a}\tilde{b}} \cdot (C^b_{cd}\Phi^c_{\tilde{m}}\Phi^d_{\tilde{n}} - \mu^b_{\hat{i}}f^{\hat{i}}_{\tilde{m}\tilde{n}} - \Phi^b_{\tilde{e}}f^{\tilde{e}}_{\tilde{m}\tilde{n}})] + \\
& -\frac{\lambda^2\rho^{n+2}r^{n_1-2}}{2}h_{ab}\alpha^a_i\tilde{H}^i g^{[\tilde{a}\tilde{b}]}(C^b_{cd}\Phi^c_{\tilde{a}}\Phi^d_{\tilde{b}} - \mu^b_{\hat{i}}f^{\hat{i}}_{\tilde{a}\tilde{b}} - \Phi^b_{\tilde{d}}f^{\tilde{d}}_{\tilde{a}\tilde{b}}) + \\
& -r^{n_1}\rho^{n-2}(\overline{M}\tilde{g}^{(\gamma\mu)}\rho_{,\gamma}\rho_{,\mu} + n^2g^{[\mu\nu]}g_{\delta\mu}\tilde{g}^{(\delta\gamma)}\rho_{,\nu}\rho_{,\gamma}) \Big\}. \tag{6.6 cont.}
\end{aligned}$$

Where

$$\tilde{L}^{i\mu\nu} = g^{\alpha\mu}g^{\beta\nu}\tilde{L}^i_{\alpha\beta} \tag{6.7}$$

and $\tilde{L}^i_{\alpha\beta}$ obeys (6.1).

One can define the following two Ad_H -type 2-forms with value in the Lie algebra of H , ($\mathfrak{h}$)

$$\overset{\vee}{L} = \frac{1}{2}L^d_{\tilde{a}\tilde{b}}\theta^{\tilde{a}} \wedge \theta^{\tilde{b}}X_d, \quad \overset{\vee}{Q} = \frac{1}{2}Q^d_{\tilde{a}\tilde{b}}\theta^{\tilde{a}} \wedge \theta^{\tilde{b}}X_d,$$

in such a way that

$$\overset{\vee}{L} = \overset{\vee}{\Omega} - \frac{1}{2}\overset{\vee}{Q},$$

where $\overset{\vee}{\Omega} = \frac{1}{2}H^d_{\tilde{a}\tilde{b}}\theta^{\tilde{a}} \wedge \theta^{\tilde{b}}X_d$. These forms are defined on P (over G/G_0). In this way we get similar formulae for the Higgs' field and for Yang–Mills' field.

Let us define the following quantity $M^d_{\tilde{a}\tilde{b}}$ such that

$$L^d_{\tilde{a}\tilde{b}} = H^d_{\tilde{a}\tilde{b}} - 4\pi M^d_{\tilde{a}\tilde{b}}. \tag{6.8}$$

This quantity is an analogue of the polarization tensor for the Higgs' field.

In this way we have

$$\overset{\vee}{Q} = 8\pi \overset{\vee}{M} = 4\pi M^d_{\tilde{a}\tilde{b}}\theta^{\tilde{a}} \wedge \theta^{\tilde{b}}X_d. \tag{6.8a}$$

The form $\overset{\vee}{M}$ is an Ad_H -type 2-form.

$$\tilde{H}^i = g^{[\alpha\beta]}\tilde{H}^i_{\alpha\beta}, \tag{6.9}$$

$L^a_{\alpha\beta}$ obeys (6.5) and scalar field $\Phi^a_{\tilde{b}}$ satisfies the following constraints:

$$\Phi^c_{\tilde{b}}f^{\tilde{b}}_{\hat{i}\tilde{a}} = \mu^a_{\hat{i}}\Phi^b_{\tilde{a}}C^c_{ab}. \tag{6.10}$$

Let us define the integral of action for $B(W)$

$$S = -\frac{1}{V_1V_2r^{n_1}} \int_{\mathbf{U}} ((B(W) d^n x) d^{n_1} x) d^4 x, \tag{6.11}$$

where $\mathbf{U} = V \times M \times H$, $V \subset E$.

$$V_1 = \int_H \sqrt{-\ell} d^n x, \tag{6.12}$$

where $d^n x = d\mu_H(h)$ is a biinvariant measure on a group H and

$$V_2 = \int_M \sqrt{|\tilde{g}|} d^{n_1} x, \quad (6.13)$$

$d^{n_1} x = dm(y)$ is a measure on M which is quasi-invariant with respect to the action of G on M and $d^4 x$ means integration over space-time coordinates. After some calculations one gets:

$$S = - \int_V \sqrt{-g} B(\overline{W}, g, \tilde{A}, \rho, \Phi) d^4 x, \quad V \subset E, \quad (6.14)$$

where

$$\begin{aligned} B(\overline{W}, g, \tilde{A}, \rho, \Phi) = & \rho^n \cdot \overline{R}(\overline{W}) + \frac{\lambda^2}{4} \left[8\pi \rho^{n+2} \mathcal{L}_{\text{YM}}(\tilde{A}) + \rho^{n+2} \frac{2}{r^2} \mathcal{L}_{\text{kin}}(\overset{\text{gauge}}{\nabla} \Phi) + \right. \\ & \left. + \rho^{n+2} \frac{1}{r^4} V(\Phi) - \rho^{n+2} \frac{4}{r^2} \mathcal{L}_{\text{int}}(\Phi, \tilde{A}) + \rho^{n-2} \mathcal{L}_{\text{scal}}(\rho) \right] + \lambda_c, \end{aligned} \quad (6.15)$$

$\overline{R}(\overline{W})$ is the Moffat–Ricci curvature scalar on the space-time E and plays the role of the gravitational lagrangian

$$\mathcal{L}_{\text{YM}}(\tilde{A}) = -\frac{1}{8\pi} \ell_{ij} (2\tilde{H}^i \tilde{H}^j - \tilde{L}^{i\mu\nu} \tilde{H}^j_{\mu\nu}) \quad (6.16)$$

is the lagrangian for the Yang–Mills' field with the gauge group G in the Nonsymmetric-Nonabelian Kaluza–Klein Theory. $\tilde{H}^i$ is defined in (6.9) and $\tilde{L}^{i\mu\nu}$ by (6.1).

$$\begin{aligned} \mathcal{L}_{\text{kin}}(\overset{\text{gauge}}{\nabla} \Phi) = & \frac{1}{V_2} \int_M \sqrt{|\tilde{g}|} d^{n_1} x (\ell_{ab} g^{\tilde{b}\tilde{n}} L^{a\mu}_{\tilde{b}} \overset{\text{gauge}}{\nabla}_{\mu} \Phi^b_{\tilde{n}}) = \\ = & \ell_{ab} g^{\alpha\mu} \frac{1}{V_2} \int_M \sqrt{|\tilde{g}|} d^{n_1} x (g^{\tilde{b}\tilde{n}} L^a_{\alpha\tilde{b}} \overset{\text{gauge}}{\nabla}_{\mu} \Phi^b_{\tilde{n}}) \end{aligned} \quad (6.17)$$

is the kinetic part of the lagrangian for the scalar field $\Phi^b_{\tilde{n}}$. It is easy to see that $\mathcal{L}_{\text{kin}}$ is a quadratic form with respect to $\overset{\text{gauge}}{\nabla} \Phi$ (gauge derivative with respect to the connection $\tilde{\omega}_E$) and is invariant with respect to the action of the groups H and G .

$$\begin{aligned} V(\Phi) = & \frac{\ell_{ab}}{V_2} \int_M \sqrt{|\tilde{g}|} d^{n_1} x [2g^{[\tilde{m}\tilde{n}]} (C^a_{cd} \Phi^c_{\tilde{m}} \Phi^d_{\tilde{n}} - \mu^a_{\hat{i}} f^{\hat{i}}_{\tilde{m}\tilde{n}} - \Phi^a_{\tilde{e}} f^{\tilde{e}}_{\tilde{m}\tilde{n}}) g^{[\tilde{a}\tilde{b}]} \cdot \\ & \cdot (C^b_{ef} \Phi^e_{\tilde{a}} \Phi^f_{\tilde{b}} - \mu^b_{\hat{j}} f^{\hat{j}}_{\tilde{a}\tilde{b}} - \Phi^b_{\tilde{a}} f^{\tilde{d}}_{\tilde{a}\tilde{b}}) + \\ & - g^{\tilde{a}\tilde{m}} g^{\tilde{b}\tilde{n}} L^a_{\tilde{a}\tilde{b}} (C^b_{cd} \Phi^c_{\tilde{m}} \Phi^d_{\tilde{n}} - \mu^b_{\hat{i}} f^{\hat{i}}_{\tilde{m}\tilde{n}} - \Phi^b_{\tilde{e}} f^{\tilde{e}}_{\tilde{m}\tilde{n}})] \end{aligned} \quad (6.18)$$

is the self-interacting term for the field Φ . It is invariant with respect to the action of the groups H and G . This term is a polynomial of 4th order in Φ 's (a Higgs' field

potential term).

$$\mathcal{L}_{\text{int}}(\Phi, \tilde{A}) = h_{ab}\mu_i^a \tilde{H}^i \underline{g}^{[\tilde{a}\tilde{b}]} \cdot (C_{cd}^b \Phi_{\tilde{a}}^c \Phi_{\tilde{b}}^d - \mu_{\hat{i}}^b f_{\tilde{a}\tilde{b}}^{\hat{i}} - \Phi_{\tilde{d}}^b f_{\tilde{a}\tilde{b}}^{\tilde{d}}) \quad (6.19)$$

is the term describing nonminimal coupling between the scalar field Φ and the Yang–Mills’ field where

$$\underline{g}^{[\tilde{a}\tilde{b}]} = \frac{1}{V_2} \int_M \sqrt{|\tilde{g}|} d^{n_1} x g^{[\tilde{a}\tilde{b}]}, \quad (6.20)$$

$$\lambda_c = \frac{\rho^n}{\lambda^2} \tilde{R}(\tilde{\Gamma}) + \frac{\rho^n}{r^2 V_2} \int_M \sqrt{|\tilde{g}|} \hat{R}(\hat{\Gamma}) d^{n_1} x = \frac{\rho^n}{\lambda^2} \tilde{R}(\tilde{\Gamma}) + \frac{\rho^n}{r^2} \tilde{P}. \quad (6.21)$$

This term is also invariant with respect to the action of groups H and G . One can write (6.17) and (6.18) in a different form.

$$\begin{aligned} \mathcal{L}_{\text{kin}}(\overset{\text{gauge}}{\nabla} \Phi) &= \ell_{ab}(g^{\beta\nu} L^a{}_{\tilde{b}} \overset{\text{gauge}}{\nabla}_{\mu} \Phi^b{}_{\tilde{n}})_{av} \\ &= \ell_{ab} g^{\alpha\mu} (g^{\tilde{b}\tilde{n}} L^a{}_{\alpha\tilde{b}} \overset{\text{gauge}}{\nabla}_{\mu} \Phi^b{}_{\tilde{n}})_{av}, \\ V(\Phi) &= 2\ell_{ab} ([g^{([\tilde{m}\tilde{n}][\tilde{a}\tilde{b}])} (C_{cd}^a \Phi_{\tilde{m}}^c \Phi_{\tilde{n}}^d - \mu_{\hat{i}}^a f_{\tilde{m}\tilde{n}}^{\hat{i}} - \Phi_{\tilde{e}}^a f_{\tilde{m}\tilde{n}}^{\tilde{e}}) \times \\ &\quad \times (C_{ef}^b \Phi_{\tilde{a}}^e \Phi_{\tilde{b}}^f - \mu_{\hat{j}}^b f_{\tilde{a}\tilde{b}}^{\hat{j}} - \Phi_{\tilde{d}}^b f_{\tilde{a}\tilde{b}}^{\tilde{d}}) + \\ &\quad + \underline{g}^{\tilde{a}\tilde{m}\tilde{b}\tilde{n}} L^a{}_{\tilde{a}\tilde{b}} (C_{cd}^b \Phi_{\tilde{m}}^c \Phi_{\tilde{n}}^d - \mu_{\hat{i}}^b f_{\tilde{m}\tilde{n}}^{\hat{i}} - \Phi_{\tilde{e}}^b f_{\tilde{m}\tilde{n}}^{\tilde{e}}))]_{av}, \end{aligned} \quad (6.22)$$

where

$$\begin{aligned} (\dots)_{av} &= \frac{1}{V_2} \int_M \sqrt{|\tilde{g}|} dx^{n_1} (\dots), \\ \underline{g}^{\tilde{a}\tilde{m}\tilde{b}\tilde{n}} &= g^{\tilde{a}\tilde{m}} g^{\tilde{b}\tilde{n}}, \\ \underline{g}^{([\tilde{m}\tilde{n}][\tilde{a}\tilde{b}])} &= g^{[\tilde{m}\tilde{n}]} \cdot g^{[\tilde{a}\tilde{b}]}. \end{aligned} \quad (6.24)$$

The connection $\hat{\Gamma}_{\tilde{b}\tilde{c}}^{\tilde{a}}$ on M satisfies the following compatibility conditions:

$$g_{\tilde{a}\tilde{b}} \hat{\Gamma}_{\tilde{a}\tilde{c}}^{\tilde{d}} + g_{\tilde{a}\tilde{d}} \hat{\Gamma}_{\tilde{c}\tilde{b}}^{\tilde{d}} = g_{\tilde{a}\tilde{d}} \mathcal{K}_{\tilde{b}\tilde{c}}^{\tilde{d}} + g_{\tilde{a}\tilde{b},\tilde{c}}, \quad (6.25)$$

where $\mathcal{K}_{\tilde{b}\tilde{c}}^{\tilde{d}}$ are nonholonomicity coefficients, “,” means the action of a vector field defined on M and induced by a left invariant vector field on G . For $g_{\tilde{a}\tilde{b},\tilde{c}}$ one easily finds

$$g_{\tilde{a}\tilde{b},\tilde{c}}(x) = \zeta k_{\tilde{a}\tilde{b},\tilde{c}}^0 + h_{\tilde{a}\tilde{b},\tilde{c}}^0. \quad (6.26)$$

For $\mathcal{L}_{\text{scal}}(\rho)$ we have the following:

$$\mathcal{L}_{\text{scal}}(\rho) = (\overline{M} \tilde{g}^{(\gamma\mu)} \rho_{,\gamma} \rho_{,\mu} + n^2 g^{[\mu\nu]} g_{\delta\mu} \tilde{g}^{(\delta\gamma)} \rho_{,\nu} \rho_{,\gamma}), \quad (6.27)$$

where

$$\overline{M} = (\ell_{[dc]} \ell^{[dc]} - 3n(n-1)) \geq 0. \quad (6.27a)$$

According to section 3 $B(W, \tilde{A}, \Phi, \Psi)$ is invariant with respect to the right action of the group G on the bundle $Q(E, G)$. Thus we do not see any additional dimensions. They can be easily dropped due to the integration over the group G .

Let us consider $M = S^2 \cong \text{SO}(3)/\text{SO}(2)$ (see sec. 4 and in particular Eq. (4.23)). One gets for $h^0 = h^0_{\tilde{a}\tilde{b}} \tilde{V}^{\tilde{a}} \otimes \tilde{V}^{\tilde{b}}$

$$h^0_{\tilde{a}\tilde{b}} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad (6.28)$$

where

$$\tilde{V}^1 = d\theta, \quad \tilde{V}^2 = \sin \theta d\varphi \quad (6.29)$$

and

$$d\tilde{V}^1 = 0, \quad d\tilde{V}^2 = \cot \theta \tilde{V}^1 \wedge \tilde{V}^2. \quad (6.30)$$

On $S^2 = \text{SO}(3)/\text{SO}(2)$ we introduce a coordinate system in a known way, i.e.

$$(\theta, \varphi) = \tilde{\varphi}(g \exp(\theta e_1 + \varphi e_2)), \quad g \in \text{SO}(3), \quad \text{SO}(3) \xrightarrow{\tilde{\varphi}} S^2 \quad (*)$$

and getting it in a neighbourhood of (θ_0, φ_0) . In particular for $(0, 0)$. In this way we have a local trivialization of the bundle G , (θ, φ, ψ) . Moreover we have a local section of G , σ : in a neighbourhood of $\tilde{\varphi}(g)$ due to $(*)$. The forms $\tilde{V}_1, \tilde{V}_2$ are pullbacks of V_1, V_2 .

$$\tilde{V}_i = \sigma^* V_i, \quad i = 1, 2.$$

They are dual to left-invariant vector fields $\tilde{e}_1, \tilde{e}_2$ on M , $\tilde{e}_i(\theta, \varphi) = L'_{g^{-1}} e_i(0, 0)$, $i = 1, 2$. We have

$$\begin{pmatrix} \tilde{e}_1(\theta, \varphi) \\ \tilde{e}_2(\theta, \varphi) \end{pmatrix} = \begin{pmatrix} \frac{\partial}{\partial \theta} & \\ 1 & \frac{\partial}{\partial \varphi} \end{pmatrix} \quad \text{and} \quad [\tilde{e}_1, \tilde{e}_2] = -\cot \theta \tilde{e}_2.$$

Thus we have

$$g_{\tilde{a}\tilde{b}} = \begin{pmatrix} 1 & \zeta \cos \theta \\ -\zeta \cos \theta & 1 \end{pmatrix}. \quad (6.31)$$

Substituting Eqs. (6.30–31) into (6.25) one finds

$$\widehat{\bar{F}}^1_{11} = \widehat{\bar{F}}^2_{22} = 0 \quad (6.32)$$

and

$$\begin{aligned}
\widehat{T}_{22}^1 + \widehat{T}_{11}^2 - \zeta \cos \theta \widehat{T}_{21}^2 &= \zeta \cos \theta, \\
\widehat{T}_{11}^2 + \widehat{T}_{21}^1 + \zeta \cos \theta \widehat{T}_{21}^2 &= -3\zeta \sin \theta, \\
\widehat{T}_{12}^1 + \widehat{T}_{21}^1 - \zeta \cos \theta \widehat{T}_{12}^2 + \zeta \cos \theta \widehat{T}_{21}^2 &= 2\zeta \sin \theta, \\
\widehat{T}_{12}^2 + \widehat{T}_{22}^1 + \zeta \cos \theta \widehat{T}_{12}^1 &= 0, \\
\widehat{T}_{22}^1 + \widehat{T}_{21}^2 + \zeta \cos \theta \widehat{T}_{21}^2 &= 2 \cot \theta, \\
\widehat{T}_{21}^2 + \widehat{T}_{12}^2 + \zeta \cos \theta \widehat{T}_{21}^1 - \zeta \cos \theta \widehat{T}_{12}^1 &= -2 \cot \theta.
\end{aligned} \tag{6.33}$$

The existence of the connection $\widehat{T}$ (a nonsymmetric) is guaranteed by the theorem from the second position of Ref. [3]. In order to apply it let us transform h^0 and g to holonomic system of coordinates.

One gets

$$g_{\tilde{a}'\tilde{b}'} = \begin{pmatrix} 1 & \zeta \sin \theta \cos \theta \\ -\zeta \sin \theta \cos \theta & \sin^2 \theta \end{pmatrix}, \tag{6.34a}$$

$$h^0_{\tilde{a}'\tilde{b}'} = \begin{pmatrix} 1 & 0 \\ 0 & \sin^2 \theta \end{pmatrix}, \tag{6.34b}$$

$$g = \det(g_{\tilde{a}'\tilde{b}'}) = \sin^2 \theta (1 + \zeta^2 \cos^2 \theta) \neq 0, \tag{6.35a}$$

$$h^0 = \det(h^0_{\tilde{a}'\tilde{b}'}) = 4 \sin^2 \theta \neq 0, \tag{6.35b}$$

$$\sigma = \frac{(2g_{[12]})^2}{g} = \frac{4 \cos^2 \theta}{(1 + \zeta^2 \cos^2 \theta)}, \tag{6.35c}$$

$$W = \left(\frac{\partial \sigma}{\partial \theta}, \frac{\partial \sigma}{\partial \varphi} \right) = (\sigma_{,\theta}, 0) \neq 0, \tag{6.35d}$$

Thus all the conditions are satisfied except $\theta = 0, \pi, 0 \leq \varphi \leq 2\pi$ and because of this the nonsymmetric connection $\widehat{\omega}$ exist (except singular line $\theta = 0, \pi, 0 \leq \varphi \leq 2\pi$, where it is symmetric i.e. Riemannian one). The system of linear equations (6.33) can be solved because its matrix is invertible. The determinant of this matrix is equal to

$$2(2\zeta^2 \cos^2 \theta - \zeta^3 \cos^3 \theta + \zeta \cos \theta - 1)$$

and nonzero.

Let us write the remaining connection coefficients. One gets

$$\begin{aligned}
\widehat{\bar{\Gamma}}_{22}^1 &= \frac{(3\zeta^2 \cos^2 \theta - 10\zeta^4 \cos^4 \theta - 6\zeta^4 \cos^3 \theta \sin \theta - 4\zeta^3 \frac{\cos^4 \theta}{\sin \theta} - 3\zeta^2 \cos \theta \sin \theta - 2 \cot \theta)}{2(\zeta^3 \cos^3 \theta - 2\zeta \cos^2 \theta - \zeta \cos \theta + 1)}, \\
\widehat{\bar{\Gamma}}_{11}^2 &= \frac{5\zeta \sin \theta (\zeta^2 \cos^2 \theta - 2\zeta^2 \cos \theta \sin \theta + 1)}{2(2\zeta^2 \cos^2 \theta - \zeta^3 \cos \theta + \zeta \cos \theta - 1)}, \\
\widehat{\bar{\Gamma}}_{21}^2 &= \frac{2\zeta \cos \theta (\zeta^2 \cos \theta \sin \theta + 2 \cot \theta - \zeta \cos \theta - 2\zeta \sin \theta)}{(2\zeta^2 \cos^2 \theta - \zeta^3 \cos^3 \theta + \zeta \cos \theta - 1)}, \\
\widehat{\bar{\Gamma}}_{21}^1 &= \frac{(2\zeta^3 \cos^3 \theta + 4\zeta \cos \theta \cot \theta + 2\zeta^3 \cos^2 \theta \sin \theta + 3\zeta \sin \theta - 2\zeta^2 \cos \theta \sin \theta - \zeta \cos \theta - 4\zeta^2 \cos \theta \cot \theta)}{2(\zeta^3 \cos^3 \theta - 2\zeta \cos^2 \theta - \zeta \cos \theta + 1)}, \\
\widehat{\bar{\Gamma}}_{12}^1 &= \frac{(4\zeta^3 \cos \theta \sin \theta - 5\zeta \sin \theta + \zeta^2 \cos^2 \theta + 3\zeta^2 \cos \theta \sin \theta + 2\zeta \cos \theta \cot \theta - \zeta \cos \theta - 2 \cot \theta)}{(2\zeta^2 \cos^2 \theta - \zeta^3 \cos^3 \theta - \zeta \cos \theta + 1)}, \\
\widehat{\bar{\Gamma}}_{12}^2 &= \frac{(7\zeta^4 \cos^3 \theta \sin \theta + 12\zeta^2 \cos^2 \theta \cot \theta - 5\zeta^2 \cos \theta \sin \theta - 4 \cot \theta + \zeta^4 \cos^4 \theta - 4\zeta^4 \cos^4 \theta \cot \theta - \zeta^2 \cos^2 \theta)}{2(2\zeta^2 \cos^2 \theta - \zeta^3 \cos^3 \theta + \zeta \cos \theta - 1)},
\end{aligned} \tag{6.36}$$

Eqs. (6.32–3) and (6.36) give us an example of Einstein–Kaufmann connection on a homogeneous space $-S^2 \cong \text{SO}(3)/\text{SO}(2)$. Let us calculate the Moffat–Ricci scalar of the curvature for the connection $\widehat{\bar{\Gamma}}_{\tilde{b}\tilde{c}}^{\tilde{a}}$, $(\widehat{\bar{\omega}}_{\tilde{b}}^{\tilde{a}})$. In order to do this let us find $g^{\tilde{a}\tilde{b}}$. One finds

$$g^{\tilde{a}\tilde{b}} = \frac{1}{(1 + \zeta^2 \cos^2 \theta)} \begin{pmatrix} 1 & -\zeta \cos \theta \\ \zeta \cos \theta & 1 \end{pmatrix}. \tag{6.37}$$

Moreover it is easier to calculate the scalar of curvature using results from the second position of Ref. [4]. In this paper one has found the general solution of compatibility conditions for Einstein–Kaufmann connection in 2-dimensional case. In particular

$$\widehat{\bar{\omega}}_{\tilde{b}}^{\tilde{n}} = \tilde{\omega}_{\tilde{b}}^{\tilde{n}} + \frac{1}{\bar{g}} h^{\tilde{n}\tilde{a}} \tilde{\nabla}_{\tilde{a}} k_{\tilde{b}\tilde{m}}^0 V^{\tilde{m}} - \frac{2}{\bar{g}} [(\tilde{\nabla}_{\alpha} k_{(\tilde{b}}^0{}_{\tilde{n}}) k_{\tilde{m}}^0{}_{\tilde{a}})] V^{\tilde{m}}, \tag{6.38}$$

where $\tilde{\omega}_{\tilde{b}}^{\tilde{n}}$ is the Riemannian connection for $h^0_{\tilde{n}\tilde{b}} = g_{(\tilde{n}\tilde{b})}$.

$$\bar{g} = \frac{\det(g_{\tilde{a}\tilde{b}})}{\det(h^o_{\tilde{a}\tilde{b}})}$$

is a scalar, and $\tilde{\nabla}$ means a covariant derivative with respect to $\tilde{\omega}_{\tilde{b}}^{\tilde{a}}$.

One gets

$$\begin{aligned}\widehat{R}(\widehat{\Gamma}) &= g^{\tilde{b}\tilde{m}} \tilde{R}_{\tilde{b}\tilde{m}}(\tilde{\Gamma}) + \frac{1}{4} \zeta g^{\tilde{b}\tilde{m}} \left(\tilde{\nabla}_{\tilde{b}} \left(\frac{1}{\tilde{g}} \nabla_{\tilde{a}} k^{0\tilde{a}}{}_{\tilde{m}} \right) - \tilde{\nabla}_{\tilde{m}} \left(\frac{1}{\tilde{g}} \tilde{\nabla}_{\tilde{a}} k^{0\tilde{a}}{}_{\tilde{b}} \right) \right) + \\ &\quad - \zeta^2 g^{\tilde{b}\tilde{m}} \left(\tilde{\nabla}_{\tilde{q}} \left(\frac{1}{\tilde{g}} (\tilde{\nabla} k^0_{(\tilde{b}}{}^{\tilde{q}}) k^0_{\tilde{m}}) \right) - \tilde{\nabla}_{(\tilde{m}} \left(\frac{1}{\tilde{g}} (\tilde{\nabla}_{\tilde{a}} k^0_{\tilde{b}}{}^{\tilde{q}}) k^0_{\tilde{q}}{}^{\tilde{a}} \right) \right),\end{aligned}\quad (6.39)$$

where $\tilde{R}_{\tilde{b}\tilde{m}}(\tilde{\Gamma})$ is the Ricci tensor for the Riemannian connection $\tilde{\Gamma}(\tilde{\omega}^{\tilde{a}}_{\tilde{b}})$. One gets after some algebra

$$\begin{aligned}\widehat{R}(\widehat{\Gamma}) &= R(\theta, \zeta) = \frac{2}{(1 + \zeta^2 \cos^2 \theta)} + \frac{\zeta^2 \cos^2 \theta (1 + 9\zeta^2 - 8\zeta^4 + \zeta^2 (16\zeta^2 - 1) \cos^2 \theta)}{2(1 + \zeta^2 \cos^2 \theta)^3} + \\ &\quad - \frac{4\zeta^2 \sin \theta (1 - 4 \cos \theta - 4\zeta^2 \cos^3 \theta)}{2(1 + \zeta^2 \cos^2 \theta)^3}.\end{aligned}\quad (6.39a)$$

In a general case i.e. for $\dim M \geq 3$ we should transform Eq. (6.25) to the holonomic system of coordinates and to use recurrence relations for the connection in order to find it (see Ref. [3] for more details).

In this case we can find a general solution (Ref. [3]).

$$\overline{\omega}^{\tilde{n}}_{\tilde{z}} = \tilde{\omega}^{\tilde{n}}_{\tilde{z}} + u^{\tilde{n}}_{\tilde{z}}, \quad (6.40)$$

where $\tilde{\omega}^{\tilde{n}}_{\tilde{z}}$ is a Riemannian connection for $h^0_{\tilde{a}\tilde{b}}$ and

$$u^{\tilde{n}}_{\tilde{z}} = u^{\tilde{n}}_{\tilde{z}\tilde{m}} \theta^{\tilde{m}}, \quad (6.41)$$

$$\begin{aligned}u^{\tilde{n}}_{\tilde{z}\tilde{m}} &= \frac{1}{2} (L^{\tilde{n}}_{\tilde{z}\tilde{m}} - 2\zeta^2 k^0_{[\tilde{m}}{}^{\tilde{a}} L_{\tilde{z}]\tilde{a}\tilde{b}} k^{0\tilde{n}\tilde{b}}) + \zeta h^{0\tilde{n}\tilde{e}} \{ L^{\tilde{a}}_{\tilde{e}(\tilde{z}} k^0_{\tilde{m})\tilde{a}} + \\ &\quad + \zeta k^0_{\tilde{e}}{}^{\tilde{b}} (k^0_{(\tilde{m}}{}^{\tilde{c}} L_{\tilde{z})\tilde{a}\tilde{b}} k^0_{\tilde{e}}{}^{\tilde{a}} - L_{\tilde{e}\tilde{a}\tilde{b}} k^0_{(\tilde{z}} k^0_{\tilde{m})}{}^{\tilde{c}}) \},\end{aligned}\quad (6.42)$$

for $L_{\tilde{a}\tilde{b}\tilde{c}}$ one has

$$L_{\tilde{a}\tilde{b}\tilde{c}} = -\zeta (\tilde{\nabla}_{\tilde{a}} k^0_{\tilde{b}\tilde{c}} + \tilde{\nabla}_{\tilde{b}} k^0_{\tilde{a}\tilde{c}} - \tilde{\nabla}_{\tilde{c}} k^0_{\tilde{a}\tilde{b}}), \quad (6.43)$$

$\tilde{\nabla}$ means a covariant derivative with respect to the connection $\tilde{\omega}^{\tilde{a}}_{\tilde{b}}$.

For a scalar of curvature one gets:

$$\widehat{R}(\widehat{\Gamma}) = g^{\tilde{b}\tilde{d}} \tilde{R}_{\tilde{b}\tilde{d}} + g^{\tilde{b}\tilde{d}} (\tilde{\nabla}_{\tilde{a}} u^{\tilde{a}}_{\tilde{b}\tilde{d}} - \tilde{\nabla}_{\tilde{a}} u^{\tilde{a}}_{\tilde{b}\tilde{a}}) + \frac{1}{2} g^{\tilde{b}\tilde{d}} (\tilde{\nabla}_{\tilde{b}} u^{\tilde{a}}_{\tilde{a}\tilde{d}} - \tilde{\nabla}_{\tilde{a}} u^{\tilde{a}}_{\tilde{a}\tilde{b}}). \quad (6.44)$$

Let us notice that if k^0 is induced by the Kählerian structure on $M = G/G_0$ (see sec. 4) we have

$$L_{\tilde{a}\tilde{b}\tilde{c}} = 0 \quad (6.45)$$

and

$$\widehat{\omega}^{\tilde{n}}_{\tilde{z}} = \tilde{\omega}^{\tilde{n}}_{\tilde{z}}. \quad (6.46)$$

In this case one simply gets

$$\widehat{\widehat{R}}(\widehat{\Gamma}) = g^{\tilde{b}\tilde{d}} \tilde{R}_{\tilde{b}\tilde{d}}(\omega). \quad (6.47)$$

Moreover the Riemannian Ricci tensor can be easily calculated in terms of structure constants of G and G_0 . One gets

$$\tilde{R}_{\tilde{b}\tilde{c}}(\omega) = -\frac{1}{4}h_{\tilde{b}\tilde{c}} - \frac{3}{4}h_{\hat{e}\hat{f}}f^{\tilde{e}\tilde{a}}_{\tilde{c}}f^{\tilde{f}}_{\tilde{b}\tilde{a}}. \quad (6.48)$$

Thus we have finally from (6.47)

$$\widehat{\widehat{R}}(\widehat{\Gamma}) = -\frac{1}{4}g^{\tilde{b}\tilde{c}}h_{\tilde{b}\tilde{c}} - \frac{3}{4}h_{\hat{e}\hat{f}}g^{\tilde{b}\tilde{c}}f^{\hat{e}\hat{a}}_{\tilde{c}}f^{\hat{f}}_{\tilde{b}\tilde{a}}, \quad (6.49)$$

i.e. in the case of $\widehat{\omega}^{\tilde{n}}_{\tilde{z}} = \tilde{\omega}^{\tilde{n}}_{\tilde{z}}$. In general one gets Eq. (6.44) where the first term is given by Eq. (6.49).

Let us reconsider S^2 in a different context i.e introducing on S^2 Riemannian and symplectic structures coming from $\mathbb{C}$ via stereographic projection.

One gets

$$\xi = \frac{x}{1+|z|^2}, \quad \eta = \frac{y}{1+|z|^2}, \quad \chi = \frac{|z|^2}{1+|z|^2}, \quad (6.50)$$

where $(\xi, \eta, \chi) \in S^2$ and $z = x + iy \in \mathbb{C}$ and

$$x = \frac{\xi}{1-\chi}, \quad y = \frac{\eta}{1-\chi}. \quad (6.50a)$$

Using spherical coordinates on S^2 one easily obtains

$$z = x + iy = \text{ctg} \frac{\theta}{2} e^{-i\varphi}, \quad (6.51)$$

where $-\frac{\pi}{2} < \theta < \frac{\pi}{2}$, $0 \leq \varphi < 2\pi$ and

$$\begin{aligned} \xi &= \sin \theta \cos \varphi, \\ \eta &= \sin \theta \sin \varphi, \\ \chi &= \cos \theta. \end{aligned} \quad (6.51a)$$

On the complex plane $\mathbb{C}$ we have a Kählerian structure induced by a Hermitian metric tensor

$$p = dz d\bar{z} = (dx^2 + dy^2). \quad (6.52)$$

The symplectic form is given by

$$\bar{k} = \frac{1}{2i} dz \wedge d\bar{z} = dx \wedge dy. \quad (6.53)$$

After some calculation one gets using Eq. (6.51)

$$p = \frac{1}{4 \sin^4 \frac{\theta}{2}} (d\theta^2 + \sin^2 \theta d\varphi^2) \quad (6.52a)$$

and

$$\bar{k} = \frac{1}{4 \sin^4 \frac{\theta}{2}} \left(-\frac{1}{2} \sin \theta d\theta \wedge d\varphi^2 \right). \quad (6.53a)$$

The tensor

$$ds^2 = d\theta^2 + \sin^2 \theta d\varphi^2 \quad (6.54)$$

is a Riemannian metric tensor on S^2 induced by the euclidean metric in E^3 . It is conformally equivalent to the flat geometry on $\mathbb{C}$ (locally). The symplectic form

$$-\frac{1}{2} \sin \theta d\theta \wedge d\varphi \quad (6.55)$$

corresponds to the symplectic form on $\mathbb{C}$ with the same conformal factor $(4 \sin^4 \frac{\theta}{2})^{-1}$. The metric $ds^2 = d\theta^2 + \sin^2 \theta d\varphi^2$ and the 2-form $k^0 = \sin \theta d\theta \wedge d\varphi$ from Riemannian and symplectic structure on S^2 (really Kählerian). Thus we have on S^2 a nonsymmetric tensor

$$g_{\tilde{a}\tilde{b}} = h_{\tilde{a}\tilde{b}}^0 + \zeta k_{\tilde{a}\tilde{b}}^0, \quad (6.56)$$

such that

$$\tilde{\nabla}_{\tilde{a}} k_{\tilde{b}\tilde{c}}^0 = 0, \quad (6.57)$$

$\tilde{\nabla}$ means covariant derivative with respect to Riemannian metric induced by (6.54) on S^2

$$g_{\tilde{a}\tilde{b}} = \begin{pmatrix} 1 & \zeta \sin \theta \\ -\zeta \sin \theta & \sin^2 \theta \end{pmatrix}, \quad (6.58)$$

$$g^{\tilde{a}\tilde{b}} = \frac{1}{(1 + \zeta^2) \sin^2 \theta} \begin{pmatrix} \sin^2 \theta & -\zeta \sin \theta \\ \zeta \sin \theta & 1 \end{pmatrix}. \quad (6.59)$$

One gets

$$\widehat{R}(\widehat{\Gamma}) = g^{\tilde{a}\tilde{b}} \tilde{R}(\tilde{\omega}) = \frac{4}{(1 + \zeta^2)}. \quad (6.60)$$

The form k^0 introduced here can be lifted to $\text{SO}(3)$ to get a left-invariant 2-form on $\text{SO}(3)$.

7. Conformal Transformation of $g_{\mu\nu}$. Transformation of the Scalar Field ρ

In section 6 we obtained the lagrangian density $B(\overline{W}, g, \tilde{A}, \rho, \Phi)$ such that:

$$\begin{aligned}
B(\overline{W}, g, \tilde{A}, \rho, \Phi) = \sqrt{-g} = & \left\{ \rho^n \cdot \overline{R}(\overline{W}) + \frac{\lambda^2}{4} \left[8\pi \rho^{n+2} \mathcal{L}_{\text{YM}}(\tilde{A}) + \right. \right. \\
& + \rho^{n+2} \frac{2}{r^2} \mathcal{L}_{\text{kin}}(\overset{\text{gauge}}{\nabla} \Phi) - \rho^{n+2} \frac{1}{r^4} V(\Phi) - \rho^{n+2} \frac{4}{r^2} \mathcal{L}_{\text{int}}(\Phi, \tilde{A}) \Big] + \\
& \left. - \frac{\lambda^2}{4} \rho^{n-2} \mathcal{L}_{\text{scal}}(\rho) + \frac{4}{\lambda^2 \rho^{2-n}} \tilde{R}(\tilde{\Gamma}) + \rho^n \frac{1}{r^2} \tilde{P} \right\}, \tag{7.1}
\end{aligned}$$

where $\mathcal{L}_{\text{YM}}$, $\mathcal{L}_{\text{kin}}$, V , $\mathcal{L}_{\text{int}}$, $\mathcal{L}_{\text{scal}}$ and $\tilde{P}$ are defined in section 6 by formulae (6.16), (6.17), (6.18), (6.19) and (6.21). This lagrangian density is a generalization of the lagrangian from Bergmann's paper (see Ref. [50]). P. G. Bergmann considers the lagrangian for the tensor-scalar theory of gravitation including Jordan–Thiry theory and Brans–Dicke theory. Our lagrangian is more general for two reasons. We have non-symmetric metric tensor $g_{\mu\nu}$ as a metric ($\overline{R}(\overline{W})$ is the lagrangian of gravitational field from the Nonsymmetric Theory of Gravitation). The lagrangian (7.1) possesses also apart from lagrangian for gauge field (in the Bergmann's paper it is an electromagnetic field) lagrangian for Higgs' field coupled to Yang–Mills' field. We also get two terms which play the role of the cosmological terms. In the Bergmann's paper there are four arbitrary functions of scalar field ρ , f_1 , f_2 , f_3 , f_4 . Here we have more such functions:

$$\begin{aligned}
f_1(\rho) &= \rho^n, & f_2(\rho) &= 8\pi \rho^{n+2}, \\
f_{2'}(\rho) &= -\frac{2}{r^2} \rho^{n+2}, & f_{2''}(\rho) &= \frac{1}{r^4} \rho^{n+2}, \\
f_{2'''}(\rho) &= \frac{4}{r^2} \rho^{n+2}, & f_3(\rho) &= \rho^{n+2}, \\
f_{4'}(\rho) &= \frac{4}{\lambda^2 \rho^{2-n}}, & f_{4''}(\rho) &= \frac{1}{r^2} \rho^{n-2}.
\end{aligned} \tag{7.2}$$

Now we proceed with the conformal transformation for the metric $g_{\mu\nu}$ and the transformation of the scalar field ρ . This is only the redefinition of $g_{\mu\nu}$ and ρ .

$$\rho = e^{-\Psi}, \quad (7.3)$$

$$g_{\mu\nu} \rightarrow e^{n\Psi} \cdot g_{\mu\nu} = \frac{1}{\rho^n} g_{\mu\nu}. \quad (7.4)$$

This procedure comes of course from Ref. [50]. The only difference is that $g_{\mu\nu}$ is now nonsymmetric. After transformations (7.3) and (7.4) we get the following:

$$B(\overline{W}, g, \tilde{A}, \Psi, \Phi) = \sqrt{-g} \left\{ \overline{R}(\overline{W}) + \frac{\lambda^2}{4} \left(8\pi e^{-(n+2)\Psi} \mathcal{L}_{\text{YM}}(\tilde{A}) + \frac{2e^{-2\Psi}}{r^2} \mathcal{L}_{\text{kin}}(\overset{\text{gauge}}{\nabla} \Phi) + \right. \right. \\ \left. \left. - \frac{e^{(n-2)\Psi}}{r^4} V(\Phi) - \frac{4e^{(n-2)\Psi}}{r^2} \mathcal{L}_{\text{int}}(\Phi, \tilde{A}) \right) - \frac{\lambda^2}{4} \mathcal{L}_{\text{scal}}(\Psi) + \frac{1}{\lambda^2} e^{(n+2)\Psi} \tilde{R}(\tilde{\Gamma}) + \frac{e^{n\Psi}}{r^2} \tilde{P} \right\}, \quad (7.5)$$

where

$$\mathcal{L}_{\text{scal}}(\Psi) = (\overline{M} \tilde{g}^{(\gamma\nu)} + n^2 g^{[\mu\nu]} g_{\delta\mu} \tilde{g}^{(\delta\gamma)}) \Psi_{,\nu} \Psi_{,\gamma}, \quad (7.6) \\ \overline{M} = (\ell^{[dc]} \ell_{[dc]} - 3n(n-1)).$$

It is easy to see that the scalar field Ψ is chargeless (it has no colour charges). However, it couples the gauge (Yang–Mills') field and the Higgs' field due to the terms:

$$8\pi e^{-(n+2)\Psi} \mathcal{L}_{\text{YM}}(\tilde{A}), \quad (7.7a)$$

$$+ \frac{2e^{-2\Psi}}{r^2} \mathcal{L}_{\text{kin}}(\overset{\text{gauge}}{\nabla} \Phi), \quad (7.7b)$$

$$- \frac{e^{(n-2)\Psi}}{r^4} V(\Phi), \quad (7.7c)$$

$$- \frac{4e^{(n-2)\Psi}}{r^2} \mathcal{L}_{\text{int}}(\Phi, \tilde{A}), \quad (7.7d)$$

It also couples the cosmological terms:

$$\frac{4}{\lambda^2} e^{(n+2)\Psi} \tilde{R}(\tilde{\Gamma}), \quad (7.8a)$$

$$\frac{e^{n\Psi}}{r^2} \tilde{P}. \quad (7.8b)$$

These six terms (7.7a–d) and (7.8a–b) suggest that the scalar field is massive. This is different than in Brans–Dicke theory, where the scalar field couples the trace of the energy-momentum tensor for matter. Our results from this section generalize results from Refs. [23], [24].

We can consider a more general case for the lagrangian in the theory i.e.

$$(R(W) + \beta) \sqrt{|\mathcal{K}|}. \quad (7.9)$$

In this case we get an additional cosmological term $\beta\rho^n$ or in terms of the field Ψ and after a redefinition of the nonsymmetric metric $g_{\mu\nu}$

$$\left(\beta \int dm(y) \sqrt{|\widetilde{g}|}\right) e^{n\Psi}. \quad (7.10)$$

This term can be added to remaining cosmological terms in the theory. Moreover we do not consider this term anymore because it is not in the real spirit of the Kaluza–Klein (Jordan–Thiry) Theory.

8. Symmetry Breaking. Higgs' Mechanism. Mass Generation. Cosmological Constant

Before passing to symmetry breaking in our theory we do some cosmetic manipulations, connecting constants. The connection ω on the fibre bundle P has no correct physical dimensions. It is necessary to pass in all formulae from ω to $\alpha_S \frac{1}{\sqrt{\hbar c}} \omega$

$$\omega \rightarrow \alpha_S \frac{1}{\sqrt{\hbar c}} \omega, \quad (8.1)$$

where $\hbar$ is the Planck's constant, c is the velocity of light in the vacuum and α_S is dimensionless coupling constant for the Yang–Mills' field if this field couples matter. For example in the electromagnetic case $\alpha_S = \frac{1}{\sqrt{137}}$. We use $\alpha_g = \alpha_S^2 = \frac{g^2}{\hbar c}$ where g is a coupling constant for a gauge field. The redefinition of ω is equivalent to a usual treatment in a local section $e : V \supset U \rightarrow P$, $e^* \omega = \frac{g}{\hbar c} A$. Now our quantities have correct physical dimensions.

Using (8.1) one easily writes the integral of action (6.11):

$$\begin{aligned} S = & -\frac{1}{r^2} \int_{\mathbf{U}} \sqrt{-g} d^4 x \left[\overline{R}(\overline{W}) + \frac{8\pi\lambda^2\alpha_S^2}{4c\hbar} \cdot \left(e^{-(n+2)\Psi} \cdot \mathcal{L}_{\text{YM}} + \right. \right. \\ & \left. \left. + \frac{e^{-2\Psi}}{4\pi r^2} \mathcal{L}_{\text{kin}} - \frac{e^{(n-2)\Psi}}{8\pi r^4} V(\Phi) - \frac{e^{(n-2)\Psi}}{2\pi r^4} \mathcal{L}_{\text{int}}(\Phi, \tilde{A}) - \mathcal{L}_{\text{scal}}(\Psi) \right) + \lambda_c \right]. \end{aligned} \quad (8.2)$$

Thus we get the integral of action for the matter described by the Yang–Mills' field and scalar field coupled to gravity. If we want to be in line with the ordinary coupling between gravity and matter we should put:

$$\frac{8\pi\lambda^2\alpha_S^2}{4c\hbar} = \frac{8\pi G_N}{c^4}. \quad (8.3)$$

One gets:

$$\lambda = \frac{2}{\alpha_S} \ell_{pl} = \frac{2}{\sqrt{\alpha_g}} \ell_{pl}, \quad (8.4)$$

where ℓ_{pl} is the Planck's length $\ell_{pl} = \sqrt{\frac{G_N \hbar}{c^3}} \simeq 10^{-33}$ cm. In this case we have:

$$\lambda_c = \left(\frac{e^{2(n+2)\Psi} \alpha_S^2}{\ell_{pl}^2} \tilde{R}(\tilde{\Gamma}) + \frac{e^{2(n+1)\Psi}}{r^2} \tilde{P} \right). \quad (8.5)$$

For $V(\Phi)$ one gets:

$$\begin{aligned}
V(\Phi) = & 2\ell_{ab} \left(\underline{g}^{([\tilde{m}\tilde{n}], [\tilde{a}\tilde{b}])} \times \left(\alpha_S \frac{1}{\sqrt{\hbar c}} C_{cd}^a \Phi_{\tilde{m}}^c \Phi_{\tilde{n}}^d - \frac{1}{\alpha_S} \sqrt{\hbar c} \mu_{\hat{i}}^a f_{\tilde{m}\tilde{n}}^{\hat{i}} - \Phi_{\tilde{e}}^a f_{\tilde{m}\tilde{n}}^{\tilde{e}} \right) \times \right. \\
& \times \left(\alpha_S \frac{1}{\sqrt{\hbar c}} C_{ef}^b \Phi_{\tilde{a}}^e \Phi_{\tilde{b}}^f - \frac{1}{\alpha_S} \sqrt{\hbar c} \mu_{\hat{j}}^b f_{\tilde{a}\tilde{b}}^{\hat{j}} - \Phi_{\tilde{d}}^b f_{\tilde{a}\tilde{b}}^{\tilde{d}} \right) \Bigg)_{av} + \\
& - \frac{\ell_{ab}}{V_2} \int_M \sqrt{|\tilde{g}|} d^{n_1} x \left[g^{\tilde{a}\tilde{n}} g^{\tilde{b}\tilde{m}} L^a_{\tilde{a}\tilde{b}} \cdot \right. \\
& \times \left. \left(\alpha_S \frac{1}{\sqrt{\hbar c}} C_{cd}^b \Phi_{\tilde{m}}^c \Phi_{\tilde{n}}^d - \frac{1}{\alpha_S} \sqrt{\hbar c} \mu_{\hat{i}}^b f^{\text{hati}}_{\tilde{m}\tilde{n}} - \Phi_{\tilde{e}}^b f_{\tilde{m}\tilde{n}}^{\tilde{e}} \right) \right], \tag{8.6}
\end{aligned}$$

where

$$\underline{g}^{([\tilde{m}\tilde{n}], [\tilde{a}\tilde{b}])} = g^{[\tilde{m}\tilde{n}]} g^{[\tilde{a}\tilde{b}]}. \tag{8.7}$$

Let us pass to spontaneous symmetry breaking and the Higgs' mechanism in our theory. In order to do this we look for the critical points (the minima) of the potential $V(\Phi)$. The scalar factor before $V(\Phi)$ is positive and has no influence on these considerations. However, our field Φ satisfies the constraints

$$\Phi_{\tilde{b}}^c f_{\hat{i}\tilde{a}}^{\tilde{b}} - \mu_{\hat{i}}^a \Phi_{\tilde{a}}^b C_{ab}^c = 0. \tag{8.8}$$

Thus we must look for the critical points of

$$V' = V + \psi^{\hat{i}\tilde{d}}_c (\Phi_{\tilde{b}}^c f_{\hat{i}\tilde{d}}^{\tilde{b}} - \mu_{\hat{i}}^a \Phi_{\tilde{d}}^b C_{ab}^c), \tag{8.9}$$

where $\psi^{\hat{i}\tilde{d}}_c$ is a Lagrange multiplier. Let us calculate $\frac{\delta V'}{\delta \Phi}$. One finds

$$\begin{aligned}
\frac{\delta V'}{\delta \Phi^w_{\tilde{v}}} = & \ell_{ab} \left(\left\{ \left[4 \underline{g}^{([\tilde{m}\tilde{n}], [\tilde{a}\tilde{b}])} \left(\alpha_S \frac{1}{\sqrt{\hbar c}} C_{cd}^a \Phi_{\tilde{m}}^c \Phi_{\tilde{n}}^d - \frac{1}{\alpha_S} \sqrt{\hbar c} \mu_{\hat{i}}^a f_{\tilde{m}\tilde{n}}^{\hat{i}} - \Phi_{\tilde{e}}^a f_{\tilde{m}\tilde{n}}^{\tilde{e}} \right) \right] + \right. \right. \\
& - \underline{g}^{\tilde{n}\tilde{a}, \tilde{m}\tilde{b}} \cdot L^a_{\tilde{n}\tilde{m}} - \underline{g}^{\tilde{m}\tilde{n}, \tilde{r}\tilde{p}} \cdot \left(\frac{\delta L^a_{\tilde{m}\tilde{n}}}{\delta H^b_{\tilde{a}\tilde{b}}} \right) \times \\
& \times \left(\alpha_S \frac{1}{\sqrt{\hbar c}} C_{ce}^d \Phi_{\tilde{r}}^c \Phi_{\tilde{p}}^e - \frac{1}{\alpha_S} \sqrt{\hbar c} \mu_{\hat{i}}^a f_{\tilde{r}\tilde{p}}^{\hat{i}} - \Phi_{\tilde{e}}^d f_{\tilde{r}\tilde{p}}^{\tilde{e}} \right) \times \\
& \times \left. \left(\alpha_S \frac{1}{\sqrt{\hbar c}} C_{ce}^d \Phi_{\tilde{r}}^c \delta^f_w \delta^{\tilde{v}}_{\tilde{b}} - \delta^b_w \delta^{\tilde{d}}_{\tilde{v}} f_{\tilde{a}\tilde{b}}^{\tilde{d}} \right) \right\} \Bigg)_{av} + \\
& + \psi^{\hat{i}\tilde{d}}_c (\delta^c_w f_{\hat{i}\tilde{d}}^{\tilde{v}} - \mu_{\hat{i}}^a C_{aw}^c \delta^{\tilde{v}}_{\tilde{d}}), \tag{8.10}
\end{aligned}$$

where

$$H^b_{\alpha\beta} = \left(\alpha_S \frac{1}{\sqrt{\hbar c}} C_{cd}^d \Phi_{\tilde{a}}^c \Phi_{\tilde{b}}^d - \frac{1}{\alpha_S} \sqrt{\hbar c} \mu_{\hat{i}}^b f_{\tilde{a}\tilde{b}}^{\hat{i}} - \Phi_{\tilde{e}}^b f_{\tilde{a}\tilde{b}}^{\tilde{e}} \right) \tag{8.11}$$

and $\frac{\delta L^a_{\tilde{m}\tilde{n}}}{\delta H^b_{\tilde{a}\tilde{b}}}$ satisfies the following equation:

$$\ell_{dc}g_{\tilde{m}\tilde{b}}g^{\tilde{c}\tilde{m}}\frac{\delta L^d_{\tilde{e}\tilde{a}}}{\delta H^w_{\tilde{p}\tilde{q}}} + \ell_{cd}g_{\tilde{a}\tilde{m}}g^{\tilde{m}\tilde{c}}\frac{\delta L^d_{\tilde{b}\tilde{e}}}{\delta H^w_{\tilde{p}\tilde{q}}} = 2\ell_{cd}g_{\tilde{a}\tilde{m}}g^{\tilde{m}\tilde{c}}\delta^d_w\delta^{\tilde{p}}_{\tilde{b}}\delta^{\tilde{q}}_{\tilde{c}}. \quad (8.12)$$

It is easy to see that, if

$$H^a_{\tilde{m}\tilde{n}} = 0, \quad (8.13)$$

then

$$\frac{\delta V'}{\delta \Phi} = 0, \quad (8.14)$$

if (8.8) is satisfied. This was noticed in Refs. [73], [78] and it is known in the symmetric theory. $H^a_{\tilde{m}\tilde{n}}$ is the part of the curvature of ω over the manifold M . Thus this means that Φ^0_{crt} , satisfying (8.13), is a “pure gauge”. If potential $V(\Phi)$ is positively defined, then we have the absolute minimum of V .

$$V(\Phi^0_{\text{crt}}) = 0. \quad (8.15)$$

But apart from this solution there are some others. Potential V (see (8.6)) is the result of the subtraction of two positively defined terms. This means that there exist some local minima of V and the shape of the potential is shown on Fig. 3. Values of V at these local minima are not zero. If the potential is not positively defined, the critical point Φ^0_{crt} is not a point of the minimum and the shape of the potential is as on Fig. 4. In this case we have no minimum. All of these properties depend on values of the constants ξ and ζ and on groups G , G_0 , H . The case with no minimum is not interesting from the physical point of view. Radiative corrections may change the situation as shown on Fig. 5 (see [81], [88]). Let us examine the situation described in Fig. 3. At the point of local minimum we have:

$$V(\Phi^1_{\text{crt}}) \neq 0 \quad (8.16)$$

and

$$H^a_{\tilde{m}\tilde{n}}(\Phi^1_{\text{crt}}) \neq 0. \quad (8.17)$$

This means that Φ^1_{crt} is not a “pure gauge” and the gauge configuration connected to Φ^1_{crt} is not trivial. This indicates that the local minimum is not a vacuum state. It is a “false vacuum” in contradistinction to “true vacuum” for the absolute minimum Φ^0_{crt} . It is possible that the “tunnel effect” from point A to B (see Fig. 3) results in a second or first order phase transition (see [81–85]). Such a possibility was considered by Weinberg and a second (local) minimum occurred due to radiative corrections. Here we have this from the very beginning on the classical level.

Now we answer the question of what is symmetry breaking in our theory if we choose one of the critical values of Φ^0_{crt} (We choose one of the degenerated vacuum states and the spontaneous breaking of symmetry takes place). In Ref. [73] it was shown that if




Fig. 3. Two shapes of the potential $V(\Phi)$ in the case that there exist two minima, absolute and local (the “true” and “false” vacua)




Fig. 4. The shape of potential $V(\Phi)$ in the case without minima

$H^a_{\tilde{m}\tilde{n}} = 0$ and (8.8) is satisfied then the symmetry is reduced to G_0 . In the case of the second minimum (local minimum—false vacuum), the unbroken symmetry will be in general different. Let us call it G'_0 and its Lie algebra $\mathfrak{g}'_0$. This will be the symmetry which preserves Φ_{crt}^1 and the constraint (8.8) imposed on Φ_{crt}^1 . It is easy to see that the Lie algebra of this unbroken group preserves Eq. (8.10) and constraint (8.8) under

Ad-action. Thus it is necessary to find all symmetries of these equations. The other possibility is to find symmetries of the hypersurface, defined by $V(\Phi) = \text{const.}$ and the constraints (8.8). This strongly depends on the details concerning ξ , ζ , G , G_0 , H in contradistinction to the absolute minimum Φ_{crt}^0 —true vacuum. Maybe it would be possible to find a similar geometrical interpretation as in the case of Φ_{crt}^0 . But in general this group will be completely different.




Fig. 5. The possible influence of radiative corrections on the shape of the potential $V(\Phi)$ in the case without minima

For the symmetry group of V is larger than G (it is H) we expect some scalars which remains massless after the symmetry breaking in both cases (i.e. $k = 0$, $k = 1$, “true” and “false” vacuum cases). They became massive only through radiative corrections. They are often referred as the pseudo-Goldstone bosons.

Let us consider a problem of the Higgs’ multiplet $\Phi_{\tilde{a}}^c$ on E . One can find a representation space (N) of $\Phi_{\tilde{a}}^c$ in the following way (see Ref. [80]). Let

$$\text{Ad}_G \rightarrow \sum_i \oplus \underline{n}_i \oplus \text{Ad}_{G_0} \quad (8.18)$$

be the decomposition of the adjoint representation of G , where $\underline{n}_i$ are irreducible representations of G_0 and let us consider the branching rule of Ad_H

$$\text{Ad}_H \rightarrow \sum_j \oplus (\underline{n}'_j \otimes \underline{m}_j), \quad (8.19)$$

where $\underline{n}'_j$ are irreducible representations of G_0 and $\underline{m}_j$ are irreducible representations of G . The latest formula comes from the known fact that

$$G_0 \otimes G \subset H. \quad (8.20)$$

Thus for each pair $(\underline{n}_i, \underline{n}'_j)$ where $\underline{n}_i$ and $\underline{n}'_j$ are identical irreducible representations of G_0 there is an $\underline{m}_j$ multiplet of Higgs' field on E . In this way we can decompose Φ into a sum:

$$\Phi = \sum_{(\underline{n}_i, \underline{n}'_j)} \oplus \Phi_{\underline{m}_j}^{(\underline{n}_i, \underline{n}'_j)}, \quad \text{or} \quad N = \sum_{(\underline{n}_i, \underline{n}'_j)} \oplus \underline{m}_j.$$

Thus the multiplet of Higgs' field is quite complicated in contradistinction to the usual case where the Higgs' field belongs to the adjoint representation of a chosen group. Moreover in our case we have a smaller number of parameters in the theory. The theory (it means a symmetry breaking) is established by a coupling constant α_S , a radius r , parameters coming from the nonsymmetry of the theory ξ, ζ a homomorphism μ , an embedding of G in $H(\alpha_i^c)$ and an embedding of G_0 in G i.e. the manifold M .

In the classical approach we need much more parameters in order to get a desired symmetry breaking pattern.

Let us notice that in the branching rule for Ad_H we have a term $\text{Ad } G_0 \otimes \mathbb{I}$ and in the decomposition for $\text{Ad } G$, the $\text{Ad } G_0$. Thus the space N contains the representation space for $\mathbb{I}$.

Let us pass to the integral of action (8.2) in the two vacuum cases: Φ_{crt}^0 and Φ_{crt}^1 . Let us expand the Higgs' field $\Phi_{\tilde{b}}^{\tilde{a}}$ in the neighbourhood of $(\Phi_{\text{crt}}^k)^{a_{\tilde{b}}}$, $k = 0, 1$.

$$\Phi_{\tilde{b}}^{a_{\tilde{b}}} = (\Phi_{\text{crt}}^k)^{a_{\tilde{b}}} + (\phi^k)^{a_{\tilde{b}}} \quad (8.21)$$

and apply this formula for $e^* \overset{\text{gauge}}{\nabla}_\mu \Phi_{\tilde{a}}^{b_{\tilde{a}}}$ and $V(\Phi)$. We get

$$e^* \overset{\text{gauge}}{\nabla}_\mu \Phi_{\tilde{a}}^{b_{\tilde{a}}} = e^* \overset{\text{gauge}}{\nabla}_\mu (\phi^k)^{b_{\tilde{a}}} + \alpha_S \frac{1}{\sqrt{\hbar c}} ((\Phi_{\text{crt}}^k)^{a_{\tilde{a}}} C_{ac}^b \alpha_j^c \tilde{A}_\mu^j + (\Phi_{\text{crt}}^k)^{a_{\tilde{b}}} f_{\tilde{a}j}^{\tilde{b}} \tilde{A}_\mu^j), \quad (8.22)$$

$$V(\Phi) = V(\Phi_{\text{crt}}^k) + \tilde{V}^k(\phi^k), \quad k = 0, 1, \quad (8.23)$$

where $V(\Phi_{\text{crt}}^k)$ is the value of V for the critical value of Φ and $\tilde{V}^k$ is the polynomial of fourth order in ϕ^k . If we put (8.19) to (6.4) and to (6.5) we can solve these equations with respect to $L^a_{\tilde{m}\tilde{n}}$ and $L^b_{\mu\alpha}$ in a certain order of approximation (there are some procedures in N.G.T. (see Refs. [79], [80] and sec. 17). Thus we can expand the kinetic

term $\mathcal{L}_{\text{kin}}(\overset{\text{gauge}}{\nabla}_\mu \Phi)$ with respect to the decomposition (8.19) and we get a mass-matrix for vector bosons $\tilde{A}_\mu^j$ which strongly depends on Φ_{crt}^k .

$$-e^{-2\Psi} N^{\mu\nu} M_{ij}^2(\Phi_{\text{crt}}^k) \tilde{A}_\mu^i \tilde{A}_\nu^j. \quad (8.24)$$

Let us consider the expression

$$(\Phi_{\text{crt}}^k)^m {}_{\tilde{n}} C_{ms}^b \alpha_i^s + (\Phi_{\text{crt}}^k)^b {}_{\tilde{m}} f^{\tilde{m}}{}_{\tilde{n}i} \quad (*)$$

in order to find its interpretation. One easily notices that it equals to

$$([\text{Ad}'_H(Y_i) + \text{Ad}'_G(Y_i)] \cdot [\Phi_{\text{crt}}^k])^b {}_{\tilde{m}}$$

(Ad_H and Ad_G mean the adjoint representation of the group H and G respectively.)

Thus if $k = 0$ $(*)$ equals zero for $Y_i \in \mathfrak{g}_0$ and if $k = 1$ $(*)$ equals zero for $Y_i \in \mathfrak{g}'_0$. The latest statements come from the invariancy of the vacuum state with respect to the action of the group G_0 for $k = 0$ (G'_0 for $k = 1$). Generators of $\mathfrak{g}_0$ ($\mathfrak{g}'_0$) should annihilate vacuum state. Thus the matrix elements $M_{ij}^2(\Phi_{\text{crt}}^k)$ are zero for i, j corresponding to $\mathfrak{g}_0$ ($\mathfrak{g}'_0$).

From the invariancy of the potential V with respect to the action of the group G one gets

$$\left. \frac{\partial^2 V}{\partial \Phi^b {}_{\tilde{n}} \partial \Phi^d {}_{\tilde{d}}} \right|_{\Phi = \Phi_{\text{crt}}^k} (T^b {}_{\tilde{n}} {}^c {}_{\tilde{c}})_i [\Phi_{\text{crt}}^k]^{\tilde{c}}{}_c = 0, \quad (**)$$

where

$$(T^b {}_{\tilde{n}} {}^c {}_{\tilde{c}})_i [\Phi_{\text{crt}}^k]^{\tilde{c}}{}_c = [\Phi_{\text{crt}}^k]^m {}_{\tilde{n}} C_{ms}^b \alpha_i^s + [\Phi_{\text{crt}}^k]^b {}_{\tilde{m}} f^{\tilde{m}}{}_{\tilde{n}i}.$$

Matrix $N^{\mu\nu}$ depends on the tensor $g_{\alpha\beta}$ and for $g_{\alpha\beta} = g_{\beta\alpha}$ we have $N^{\mu\nu} = g^{\mu\nu}$. In the zeroth order of approximation for $g_{\alpha\beta}$ we get:

$$-e^{-2\Psi} \eta^{\mu\nu} M_{ij}^2(\Phi_{\text{crt}}^k) \tilde{A}_\mu^i \tilde{A}_\nu^j, \quad (8.24a)$$

where $\eta^{\mu\nu}$ is the Minkowski tensor. Eigenvalues of $M_{ij}^2(\Phi_{\text{crt}}^k)$ are the squares of the masses of the gauge bosons. The secular equation $\det(M^2 - m^2 J) = 0$ gives us a mass spectrum of massive vector bosons.

Thus there is an orthogonal matrix $(A^i{}_j) = A$ such that $A^T = A^{-1}$ and

$$A^{-1} M^2(\Phi_{\text{crt}}^k) A = \begin{bmatrix} m_1^2(\Phi_{\text{crt}}^k) & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & m_{\ell_k}^2(\Phi_{\text{crt}}^k) \end{bmatrix}, \quad k = 0, 1. \quad (8.25)$$

$\ell_0 = n_1$, $\ell_1 = \dim G - \dim G'_0$.

In this way we transform the broken vector fields into massive vector fields

$$\tilde{B}^{i'}{}_\mu = \sum_{j=1}^{\ell_k} A^{i'}{}_j \tilde{A}^j{}_\mu, \quad (8.26)$$

such that

$$\eta^{\mu\nu} \sum_{j=1}^{l_k} m_j^2(\Phi_{\text{crt}}^k) \tilde{B}_{\mu}^j \tilde{B}_{\nu}^j = \eta^{\mu\nu} M_{ij}^2(\Phi_{\text{crt}}^k) \tilde{A}_{\mu}^i \tilde{A}_{\nu}^j. \quad (8.27)$$

The matrix A depends of course, on the critical point of the Higgs' potential.

Moreover we should remember the formula (4.51) which is a constraint for Higgs field. In this way we should differentiate with respect to independent components of Φ . In all the formulae concerning Higgs field we suppose (removing spurious components) that we have to do with independent components. Solving the equation (4.51) we get independent components of Higgs field. Let the components be $\hat{\Phi}^{a_1}_{\tilde{a}_1}$ and

$$\Phi^a_{\tilde{a}} = \Phi^a_{\tilde{a}}(\hat{\Phi}^{a_1}_{\tilde{a}_1}).$$

In this case we get

$$\frac{\delta V}{\delta \hat{\Phi}^{a_1}_{\tilde{a}_1}} = \frac{\delta V}{\delta \Phi^a_{\tilde{a}_1}} \frac{\delta \hat{\Phi}^{a_1}_{\tilde{a}_1}}{\delta \hat{\Phi}^{a_1}_{\tilde{a}_1}}$$

and

$$\frac{\delta^2 V}{\delta \hat{\Phi}^{b_1}_{\tilde{b}_1} \delta \hat{\Phi}^{a_1}_{\tilde{a}_1}} = \frac{\delta^2 V}{\delta \Phi^b_{\tilde{b}} \delta \Phi^a_{\tilde{a}}} \cdot \frac{\delta \Phi^b_{\tilde{b}}}{\delta \hat{\Phi}^{b_1}_{\tilde{b}_1}} \frac{\delta \Phi^a_{\tilde{a}}}{\delta \hat{\Phi}^{a_1}_{\tilde{a}_1}}.$$

In all the formulae containing the derivatives of V we suppose to pass from $\frac{\delta}{\delta \Phi}$ to $\frac{\delta}{\delta \hat{\Phi}}$ using above formulae (the same in the matrix below).

The matrix of the masses for Higgs' bosons can be obtained in a similar way i.e.

$$V(\Phi) = V(\Phi_{\text{crt}}^k) + \frac{1}{2} \frac{\delta^2 V}{\delta \Phi^a_{\tilde{a}} \delta \Phi^b_{\tilde{b}}} \Big|_{\Phi=\Phi_{\text{crt}}^k} \phi_{\tilde{a}}^a \phi_{\tilde{b}}^b + \dots \quad (8.28)$$

The matrix

$$m^2(\Phi_{\text{crt}}^k)_{\tilde{a} \tilde{b}} = - \frac{\delta^2 V}{\delta \Phi^a_{\tilde{a}} \delta \Phi^b_{\tilde{b}}} \Big|_{\Phi=\Phi_{\text{crt}}^k} \quad (8.29)$$

is positively defined and symmetric. Thus we can diagonalize it and we get

$$\frac{\delta^2 V}{\delta \Phi^a_{\tilde{a}} \delta \Phi^b_{\tilde{b}}} \Big|_{\Phi=\Phi_{\text{crt}}^k} \phi_{\tilde{a}}^a \phi_{\tilde{b}}^b = - \sum_{j=1}^{\ell_k} m_j^2(\Phi_{\text{crt}}^k)_{\tilde{a} \tilde{b}} \Psi_{\tilde{a}}^a \Psi_{\tilde{b}}^a, \quad (8.30)$$

where

$$\Psi_{\tilde{a}}^a = \sum_{b, \tilde{b}} A_{\tilde{b}}^{\tilde{a}} A_{\tilde{a}}^b \phi_{\tilde{b}}^b. \quad (8.31)$$

The matrix $A_{\tilde{a}}^{\tilde{b}}$ depends on the critical point of the Higgs' potential. For the mass matrix one has:

$$(A^{-1})^{\tilde{c} \tilde{a}} m^2(\Phi_{\text{crt}}^k)_{\tilde{a} \tilde{b}} A_{\tilde{d}}^{\tilde{b}} A_{\tilde{c}}^{\tilde{d}} = (m^2(\Phi_{\text{crt}}^k)_{\tilde{c} \tilde{d}}) \delta^{\tilde{d} \tilde{c}} \delta^{\tilde{c} \tilde{d}}. \quad (8.32)$$

The eigenvalue problem for $m^2(\Phi_{\text{crt}}^k)$ can be posed as follows

$$m^2(\Phi_{\text{crt}}^k)^{\tilde{a}\tilde{b}} X^a_{\tilde{a}} = m^2(\Phi_{\text{crt}}^k) X^{\tilde{b}}_{\tilde{b}}. \quad (8.33)$$

The mass spectrum of Higgs' particles one gets from the secular equation

$$\det([m^2(\Phi_{\text{crt}}^k)] - m^2(\Phi_{\text{crt}}^k)J) = 0. \quad (8.34)$$

One easily writes the shape of mass matrix:

$$\begin{aligned} m^2(\Phi_{\text{crt}}^k)^{\tilde{a}}_{\tilde{a}}{}^{\tilde{r}}_{\tilde{r}} = & -\frac{1}{r^4} \cdot \frac{\delta^2 V}{\delta H^c_{\tilde{n}\tilde{m}} \delta H^d_{\tilde{p}\tilde{q}}} \left[\alpha_s \sqrt{\frac{\hbar}{c}} C^c_{av} (\delta^{\tilde{a}}_{\tilde{n}} (\Phi_{\text{crt}}^k)^v_{\tilde{m}} - (\Phi_{\text{crt}}^k)^v_{\tilde{n}} \delta^{\tilde{a}}_{\tilde{m}}) - f^{\tilde{a}}_{\tilde{n}\tilde{m}} \delta^c_{\tilde{a}} \right] \cdot \\ & \cdot \left[\alpha_s \sqrt{\frac{\hbar}{c}} C^d_{ru} (\delta^{\tilde{r}}_{\tilde{p}} (\Phi_{\text{crt}}^k)^u_{\tilde{q}} - (\Phi_{\text{crt}}^k)^u_{\tilde{p}} \delta^{\tilde{r}}_{\tilde{q}} - f^{\tilde{r}}_{\tilde{p}\tilde{q}} \right] + \\ & + \frac{\alpha_s}{r^4} \sqrt{\frac{\hbar}{c}} \left(\frac{\delta V}{\delta H^c_{\tilde{n}\tilde{m}}} \right) C^c_{ar} (\delta^{\tilde{a}}_{\tilde{n}} \delta^{\tilde{r}}_{\tilde{m}} - \delta^{\tilde{a}}_{\tilde{m}} \delta^{\tilde{r}}_{\tilde{n}}). \end{aligned} \quad (8.35)$$

The diagonalization procedure of the matrix $m^2(\Phi_{\text{crt}}^k)^{\tilde{a}}_{\tilde{a}}{}^{\tilde{b}}_{\tilde{b}}$ can be achieved in the two following ways. The first way more theoretical is as follows.

The matrix defines a quadratic form on the representation space— N for Higgs' field. Moreover the space N can be decomposed into Higgs' multiplets $\underline{m}_j$ and according this decomposition the matrix can be written in a block diagonal form

$$[m^2(\Phi_{\text{crt}}^k)] = \sum_j \oplus m_j^2(\Phi_{\text{crt}}^k). \quad (8.36)$$

We can diagonalize every matrix $m_j^2(\Phi_{\text{crt}}^k)$. Corresponding to the multiplet $\underline{m}_j$.

The second way is based on the following observation.

The matrix $m^2(\Phi_{\text{crt}}^k)^{\tilde{a}}_{\tilde{a}}{}^{\tilde{b}}_{\tilde{b}}$ can be transformed into a different matrix $(n_1 n) \times (n_1 n)$ forming an index from two indices $\tilde{a}$ and a

$$\bar{a} = \alpha a + \beta \tilde{a} + \gamma, \quad (8.37)$$

where α, β, γ are integers. The new index $\bar{a}$ should be unambiguous. Thus we must choose α, β, γ in such a way that for every $\bar{a} \in N_1^{n_1 n}$ the equation (8.37) has only one solution for $a \in N_1^n, \tilde{a} \in N_1^{n_1}$.

After this we can diagonalize $m^2(\Phi_{\text{crt}}^k)$ as an ordinary matrix. In sec. 17 we consider some expansion procedures for $m^2(\Phi_{\text{crt}}^k)$ and the second way is more convenient in this case.

What is the scale of these masses? It is easy to see that:

$$m_{\bar{A}} \cong \frac{\alpha_S}{r} \left(\frac{\hbar}{c} \right), \quad (8.38)$$

where $m_{\bar{A}}$ is the mass of a typical vector boson which obtained mass due to the Higgs' mechanism. The parameter, radius r , defines the scale of these masses as in Refs. [75], [80]. The spectrum of masses of all massive vector bosons depends, of course, on the

details of the theory. One can say the same about masses of scalar particles (Higgs' particles). Due to decomposition (8.18), we get mass terms from the expansion (8.20) and the scale of mass for massive scalars will be the same as for massive vector bosons. We can also define a so called deconfinement parameter $\Lambda(G)$ of order of a mass of typical Higgs' particle. $\Lambda(G)$ in our theory is of the same order as $m_{\tilde{A}}$. An interesting point will arise here if some of these particles remain without a rest mass or not. This has some importance in the hierarchy of mass scales and the hierarchy of physical interactions (see Ref. [89]). Moreover we expect massless scalars, the so-called pseudo-Goldstone bosons. It is interesting to notice that the scale of masses is the same in both cases $k = 0, 1$, however there is a different spectrum of masses for the vector and scalar particles and with different unbroken groups. It is easy to notice that due to a factor $e^{-2\Psi}$ in Eq. (8.22) we will have to do with an effective scale of masses depending on Ψ .

Let us consider the following decomposition of the connection ω_E defined on the principal fibre bundle $Q(E, G)$

$$\omega_E = \omega_E^0 + \sigma_E, \quad (8.39)$$

$$\omega_E^0 \in \mathfrak{g}, \quad \sigma_E \in \mathfrak{m}, \quad (8.40)$$

corresponding to the decomposition of the Lie algebra $\mathfrak{g}$

$$\mathfrak{g} = \mathfrak{g}_0 \dot{+} \mathfrak{m}.$$

In this way we consider a reduction of $Q(E, G)$ to $Q_0(E, G)$ induced by an embedding of G_0 into G . The form ω_E^0 is a connection defined on $Q_0(E, G)$ and σ_E is a tensorial form defined on $Q(E, G)$. We suppose that the reduction of the bundle Q to Q_0 is possible. The condition for this is quoted in sec. 12. In this way the transport of the connection is possible, because G_0 is a closed subgroup of G . The form ω_E^0 corresponds to the Yang–Mills' field (massless vector bosons) which remains after symmetry breaking. The tensorial form σ_E corresponds to massive vector bosons.

One gets for the curvature form

$$\Omega_E = \Omega_E^0 + D^0 \sigma_E + [\sigma_E, \sigma_E], \quad (8.41)$$

where Ω_E^0 is a curvature form for ω_E^0 and D^0 means a covariant derivative with respect to ω_E^0 . Thus

$$\Omega_E^0 = \frac{1}{2} \tilde{H}^i_{\mu\nu} \theta^\mu \wedge \theta^\nu \beta^i_{\hat{i}} Y_i, \quad (8.42a)$$

$$\sigma_E = \sigma^{\tilde{a}} Y_{\tilde{a}}, \quad (8.42b)$$

$$e^* D^0 \sigma_E = \overset{\text{gauge}}{\nabla}_{[\mu}^0 \bar{\sigma}_{\nu]}^{\tilde{a}} \bar{\theta}^\mu \wedge \bar{\theta}^\nu Y_{\tilde{a}}, \quad (8.42c)$$

$$\overset{\text{gauge}}{\nabla}_{[\mu}^0 \bar{\sigma}_{\nu]}^{\tilde{a}} = \partial_{[\mu} \bar{\sigma}_{\nu]}^{\tilde{a}} + f^{\tilde{a}}_{\tilde{b}i} \beta^i_{\hat{i}} \bar{\sigma}_{[\nu}^{\tilde{b}} \bar{A}_{\mu]}^{\hat{i}}, \quad (8.42d)$$

$$e^* \sigma_E = \bar{\sigma}_\nu^{\tilde{a}} \bar{\theta}^\nu Y_{\tilde{a}} = \bar{\sigma}_\nu^i \bar{\theta}^\nu \beta_i^{\tilde{a}} Y_{\tilde{a}}, \quad (8.42e)$$

$$e^* \omega_E^0 = \tilde{A}_\mu^i \beta_i^{\tilde{a}} \bar{\theta}^\mu Y_{\tilde{a}}, \quad (8.42f)$$

The matrices $\beta_i^{\tilde{a}}, \beta_{\tilde{a}}^i$ define an embedding of $\mathfrak{g}_0$ into $\mathfrak{g}$ (G_0 into G), ∇_μ^0 means a gauge derivative with respect to the connection ω_E^0 and e is a local section.

The above decomposition of ω_E corresponds to a case, where the symmetry group is broken from G to G_0 , i.e. to the first type of the critical points of the Higgs' potential. Moreover in the case of the second type of the critical point we can repeat the decomposition of ω_E according to the decomposition

$$\mathfrak{g} = \mathfrak{g}'_0 \dot{+} \mathfrak{m}', \quad (8.43)$$

where $\mathfrak{g}'_0$ corresponds to $G]_0$ —a group which remains unbroken. One gets

$$\omega_E = \omega_E^{'o} + \sigma_E'$$

and $Q(E, G)$ is reduced to $Q(E, G'_0)$.

One easily connects σ_E and σ_E' with $\tilde{B}$ (i.e. fields with defined non-zero rest mass, because $\tilde{A}$ have not defined masses) fields

$$e^* \sigma_E = \frac{g}{\hbar c} \tilde{B}_\mu^i \bar{\theta}^\mu Y_{i'}, \quad (8.44)$$

where

$$Y_{i'} = (A^{-1})_{i'}^i Y_i = (A^{-1})_{i'}^i \beta_i^{\tilde{a}} Y_{\tilde{a}}. \quad (8.45)$$

The matrix A is defined by the Eq. (8.25*). The same holds for σ_E' .

Let us consider the following gauge transformation i.e. a change of a local section of $Q(E, G)$ from e to f

$$e(x) = U^{(k)}{}^{-1}(x) f(x), \quad (***)$$

where

$$U^{(k)}(x) = \exp\left(\sum_{\tilde{a}} \eta_{\tilde{a}}^\nu(x) Y_{\tilde{a}}^\nu\right),$$

$$\text{for } k = 0, Y_{\tilde{a}}^\nu \in \mathfrak{m} \text{ i.e. } \tilde{a} = \tilde{a}, \quad \text{for } k = 1, Y_{\tilde{a}}^\nu \in \mathfrak{m}'.$$

and $\eta_{\tilde{a}}^\nu(x)$ is a multiplet of scalar fields on E transforming according to the $\text{Ad } G_0$ ($\text{Ad } G'_0$). $Y_{\tilde{a}}^\nu$ span $\mathfrak{m}$ or $\mathfrak{m}'$ ($k = 0$ or $k = 1$). Such a gauge condition is called a “unitary gauge”.

Let us consider the following parametrization of the Higgs' field

$$\Phi_{\tilde{a}}^c = \text{Ad}_G(U(x))_{\tilde{a}}^{\tilde{a}'} \cdot \text{Ad}_H(U(x))_{c'}^c \cdot ((\Phi_{\text{crt}}^k)_{\tilde{a}'}^{c'} + \phi_{\tilde{a}'}^{(k)c'}(x)). \quad (****)$$

For $k = 0$ we suppose that $\phi_{\tilde{a}}^{(0)c_0} = 0$, where c_0 is an established index. For $k = 1$ it is more complex and strongly depends on the details of the theory. Moreover this is always possible to give similar condition. The nonzero $\phi_{\tilde{a}}^{(k)c}$ fields are usually called surviving Higgs' fields.

Thus we can transform Higgs' and gauge fields i.e.

$$\begin{aligned} (\Phi^U)^c_{\tilde{a}} &= \text{Ad}_H(U^{-1})^c_{c'} \cdot \text{Ad}_G(U^{-1})^{\tilde{a}'}_{\tilde{a}} \Phi^c_{\tilde{a}'} = (\Phi^k_{\text{crt}})^a_{\tilde{a}} + \varphi^c_{\tilde{a}}(x), \\ \tilde{B}'_i(x) Y_i &= \text{Ad}_H(U(x))^i_j \tilde{B}^j_\mu(x) Y_i - \frac{\hbar c}{g} \partial_\mu U(x) U^{-1}(x). \end{aligned}$$

One easily notice that

$$[\phi^k_{\text{crt}}]_{\tilde{a}}^c = [\eta^k_{\text{crt}}]_{\tilde{a}} = 0. \quad (8.46)$$

Moreover simple counting of the degrees of freedom on both sides of the Eq. (****) reveals that some of $\phi_{\tilde{a}}^{(k)c}$ fields are not independent (neglecting the fact that $\Phi^c_{\tilde{a}}$ satisfies a constraint).

Thus we should constrain $\phi_{\tilde{a}}^{(k)c}$ in such a way that $(\dim G - \dim G_0) \dim H = \dim G^{(k)} + \text{rank}(\phi^{(k)})$ where $G = G_0$ and $G = G'_0$. One easily gets

$$\nabla_\mu^0 \Phi^c_{\tilde{a}} = \text{Ad}_H(U(x))^c_{c'} \cdot \text{Ad}_G(U^{-1}(x))^{\tilde{a}'}_{\tilde{a}} \cdot \nabla_\mu^0 (\Phi^u)^{c'}_{\tilde{a}'}.$$

On the level of the tensorial form σ_E one establishes that

$$f^* \sigma_E^{(k)} = \text{Ad}_G(U^{-1}(x))(e^* \sigma_E^{(k)}) - d U(x) \cdot U^{-1}(x)$$

and

$$\mathcal{L}_{\text{YM}}(\tilde{A}) = \mathcal{L}_{\text{YM}}(\tilde{B}) = \mathcal{L}_{\text{YM}}(\tilde{B}').$$

It is important to notice that we should consider a local section for ℓ_{ij} i.e $a_{ij} = e^* \ell_{ij}$ in the lagrangian for the Yang–Mills' field.

The fields $\tilde{B}'$ are massive with the same masses as $\tilde{B}$. The important point to notice is that the full lagrangian is still G —gauge invariant. Moreover a choice of a particular value of Φ^k_{crt} (which is G_0 invariant) reduces symmetry from G to G_0 (spontaneously). The fields $\eta_a^\nu(x)$ disappear. They are “eaten” by the gauge transformation and due to this the massive vector fields have 3 polarization degrees of freedom. Sometimes $\eta_a^\nu(x)$ are called “would-be Goldstone bosons”. In the matrix of masses they corresponds to zero modes. It means the mass matrix $m^2(\Phi^k_{\text{crt}})^{\tilde{a}}_a \tilde{b}_b$ should be transformed into $m^2(\Phi^0_{\text{crt}})^{\tilde{a}}_a \tilde{b}_b$ where $a, b \neq c_0$ for $k = 0$ and for $k = 1$ into $m^2(\Phi^1_{\text{crt}})^{\tilde{a}'}_{a'} \tilde{b}'_{b'} \Lambda^{\tilde{a}'}_a \tilde{a}_{\tilde{a}'} \Lambda^{b'}_b \tilde{b}_{\tilde{b}'}$, where $\Lambda^{b'}_b \tilde{b}_{\tilde{b}'} = 0$ for those $\tilde{a}, a$ for which $\phi_{\tilde{a}}^{(1)a}$ is constrained to zero. Moreover we can check it by a direct calculations of the eigenvalues of $m^2(\Phi^k_{\text{crt}})^{\tilde{a}}_a \tilde{b}_b$ that it has

$(\dim G - \dim G_0)$ $((\dim G - \dim G'_0))$ zero eigenvalues corresponding to the “eaten” Goldstone bosons. This result can be obtained directly using invariancy of the classical vacuum with respect to the action of the group G_0 or G'_0 .

One of simplest condition to choose independent parametrization of Φ_a^a is to take such $\phi_a^{(k)}$ that

$$\sum_{\tilde{a}} \phi_{\tilde{a}}^{(k) c_0} Y_{\tilde{a}} = \phi^{(k) c_0}$$

and

$$\sum_{\tilde{a}} \eta_{\nu}^{(k) c_0} Y_{\tilde{a}} = \eta^{(k) c_0}$$

are orthogonal in $\mathfrak{g}$ i.e. $\tilde{h}(\phi^{(k) c_0}, \eta^{(k)}) = 0$, where c_0 is an established index.

For both cases if the vector boson $\tilde{B}^i$ is massive after spontaneous symmetry breaking and the Higgs' mechanism, the strength of the Yang–Mills' field which corresponds to $\tilde{B}^i$ has Yukawa-type behaviour with short range

$$\tilde{H}_{\mu\nu}^i \sim \frac{1}{R} e^{-(R/r)}.$$

Due to equation (6.1) we have similarly for

$$\tilde{L}_{\mu\nu}^i \sim \frac{1}{R} e^{-(R/r)}$$

and for $\tilde{M}_{\mu\nu}^i$ (see (6.3))

$$\tilde{M}_{\mu\nu}^i \sim \frac{1}{R} e^{-(R/r)}$$

Thus we get that for the Yang–Mills polarization induced by the skew-symmetric part of the metric for “broken”, massive vector bosons is of short range. Let us pass to the cosmological terms in both of these cases.

$$\lambda_{ck} = e^{(n+2)\Psi} \frac{\alpha_S^2}{\ell_{pl}^2} \tilde{R}(\tilde{\Gamma}) + \frac{e^{n\Psi} \tilde{P}}{r^2} + e^{(n-2)\Psi} \frac{4}{\hbar c r^2} \left(\frac{\ell_{pl}^2}{r^2} \right) V(\Phi_{\text{crt}}^k), \quad k = 0, 1, \quad (8.47)$$

where $\tilde{P}$ is defined by (6.21). These two terms are different and both depend on the scalar field Ψ . Using (8.20) one gets:

$$\lambda_{ck} = e^{(n+2)\Psi} \frac{\alpha_S^2}{\ell_{pl}^2} \tilde{R}(\tilde{\Gamma}) + e^{n\Psi} \frac{m_A^2}{\alpha_S^2} \left(\frac{c}{\hbar} \right)^2 \tilde{P} + 4e^{(n-2)\Psi} \frac{m_A^4}{\alpha_S^4} \left(\frac{\hbar^3}{c^5} \right) \ell_{pl}^2 V(\Phi_{\text{crt}}^k). \quad (8.48)$$

We get that:

$$\tilde{R}(\tilde{\Gamma}) = \frac{Q_s(\xi)}{P_{s+1}(\xi)}, \quad \text{or} \quad \frac{Q_s(\xi)}{P_s(\xi)}, \quad (8.49)$$

where Q_s and P_s are polynomials of the s^{th} order and P_{s+1} is the polynomial of $(s+1)^{\text{th}}$ order with respect to ξ . Q_s and $P_{s+1}(P_s)$ have not common divisors.

$$\ell_{ab} = h_{ab} + \xi k_{ab}. \quad (8.50)$$

In a similar way we can prove that:

$$\widehat{\widehat{R}}(\widehat{\widehat{T}}) = \frac{W_k(x, \zeta)}{V_{k+1}(x, \zeta)}, \quad \text{or} \quad \frac{W_k(x, \zeta)}{V_k(x, \zeta)}, \quad x \in G/G_0, \quad (8.51)$$

where $W_k(x, \zeta)$, $V_k(x, \zeta)$ are polynomials of the k^{th} order with respect to ζ with coefficients depending on $x \in G/G_0$ and $V_{k+1}(x, \zeta)$ is the polynomial of the $(k+1)$ order with respect to ζ with coefficients depending on $x \in G/G_0$. W_k and $V_{k+1}(V_k)$ have not common divisors.

$$\widetilde{g}_{\widetilde{a}\widetilde{b}} = h_{\widetilde{a}\widetilde{b}}^0 + \zeta k_{\widetilde{a}\widetilde{b}}^0. \quad (8.52)$$

However:

$$\begin{aligned} \widetilde{\underline{P}} &= \frac{\int_M \sqrt{|\widetilde{g}|} d^{n_1} x \widehat{\widehat{R}}(\widehat{\widehat{T}})}{\int_M \sqrt{|\widetilde{g}|} d^{n_1} x} = \frac{1}{V_1} \int_M \sqrt{|\widetilde{g}|} d^{n_1} x \frac{W_k(x, \zeta)}{V_{k+1}(x, \zeta)} = \\ &= \frac{R_r(\zeta)}{S_{r+1}(\zeta)} \phi(\zeta), \quad \text{or} \quad \frac{R_r(\zeta)}{S_r(\zeta)} \phi(\zeta) \end{aligned} \quad (8.53)$$

and $\phi(\zeta)$ is a function of ζ where R_r , S_r are polynomials of the r^{th} order with respect to ζ and S_{r+1} is the polynomial of $(r+1)^{\text{st}}$ order with respect to ζ . In both cases we have similar asymptotic behaviour with respect to ξ and ζ if the function ϕ is bounded. R_r and $S_{r+1}(S_r)$ have not common divisors.

$$\widetilde{R}(\widetilde{\Gamma}) \sim \frac{C_1}{\xi} \quad \text{or} \quad \sim C_1, \quad (8.54)$$

$$\widetilde{\underline{P}} \sim \frac{C_2}{\zeta} \quad \text{or} \quad \sim C_2, \quad (8.55)$$

where C_1 and C_2 are constants. If the polynomials Q_s and R_r have real roots ξ_0 and ζ_0 such that:

$$Q_s(\xi_0) = 0 \quad (8.56)$$

and

$$R_r(\zeta_0) = 0 \quad (8.57)$$

we get:

$$\lambda_{ck}(\xi_0, \zeta_0) = 4e^{(n-2)\Psi} \frac{m_{\widetilde{A}}^4}{\alpha_S^4} \left(\frac{\hbar^3}{c^5} \right) \ell_{pl}^2 V(\Phi_{\text{crt}}^k), \quad k = 0, 1. \quad (8.58)$$

In the case for sufficiently large ξ , ζ one gets:

$$\lambda_{ck}(\xi, \zeta) = \left(e^{(n+2)\Psi} \frac{C'_1}{\xi} + e^{n\Psi} \frac{C'_2}{\zeta} \right) + 4 \frac{m_{\tilde{A}}^4}{\alpha_S^4} \left(\frac{\hbar^3}{c^5} \right) \cdot e^{(n-2)\Psi} \ell_{pl}^2 V(\Phi_{\text{crt}}^k)$$

or

$$\lambda_{ck}(\xi, \zeta) = (e^{(n+2)\Psi} C'_1 + e^{n\Psi} C'_2) + 4 \frac{m_{\tilde{A}}^4}{\alpha_S^4} \left(\frac{\hbar^3}{c^5} \right) \cdot e^{(n-2)\Psi} \ell_{pl}^2 V(\Phi_{\text{crt}}^k), \quad (8.59)$$

$k = 0, 1$; C'_1, C'_2 are constants.

Thus in some cases we are able to make the first part of λ_{ck} as small as we want. From the observational data point of view we know that the cosmological constant is almost zero. Thus it occurs in the first or second case (real roots of polynomials Q_s and R_r or in the limit of large ξ and ζ). One gets:

$$\lambda_0 = 0, \quad (8.60)$$

$$\lambda_1 = 4 \frac{m_{\tilde{A}}^4}{\alpha_S^4} \left(\frac{\hbar^3}{c^5} \right) \ell_{pl}^2 V(\Phi_{\text{crt}}^1). \quad (8.61)$$

For the “false vacuum” the cosmological constant is not zero. It is zero only for the “true vacuum”—absolute minimum of the potential V . Thus we get that worlds corresponding to two critical points of the potential V are completely different with different unbroken groups, different mass spectrum for intermediate bosons and different cosmological constants. Thus the “tunnel effect” from point A to B (see Fig. 3) results in a transition between completely different worlds.

In some cosmological models (see [81–85]) one introduces in a phenomenological way the Higgs potential of this type with two critical points: first (true vacuum) with zero cosmological constant and second (false vacuum) with nonzero cosmological constant. If the model of the universe is divided into some number of clusters with the “false” and the “true” vacuum then evolution of clusters with different vacua will be completely different. There are different laws of expansion for cosmological models in General Relativity (in Moffat’s theory too) for two cases: with zero and nonzero cosmological constants (radiation and de Sitter universes). Transition from point A to B results in the first order phase transition in models of the universe. Clusters with “true vacuum” devour clusters with “false vacuum” (see [81]). Thus we can reproduce the inflationary scenario of the Universe (old scenario). If the point A on Fig. 3 is identical with the maximum of V we can reproduce the new inflationary scenario of the Universe (without the “tunnel effect”) as well (the so-called “rolling penny” scenario). We can also work with an extended inflationary scenario, because our theory is equipped with a scalar field Ψ , which plays the role of dilatation scalar field.

In Moffat’s theory of gravitation there are some cosmological models with the phase transition of the second order in the early stages of the universe (see [90–91]). This occurs due to the skewsymmetric part of the metric $g_{\mu\nu}$ on the space-time. It is easy

to see that if the metric on the homogeneous space $M = G/G_0$, $g_{\tilde{a}\tilde{b}}$ is symmetric

$$g_{\tilde{a}\tilde{b}} = h^0_{\tilde{a}\tilde{b}}, \quad (8.62)$$

we never obtain a second local minimum for the potential V .

Let us calculate $\tilde{\underline{P}}$ for $M = S^2 = \text{SO}(3)/\text{SO}(2)$ (a cosmological constant). One gets

$$\tilde{\underline{P}} = \frac{1}{V_2} \int_{S^2} \widehat{\underline{R}}(\widehat{\Gamma}) \sqrt{|\widehat{g}|} dm = \frac{4\pi}{V_2} \int_0^\pi R(\theta, \zeta) \sqrt{1 + \zeta^2 \cos^2 \theta} \sin \theta d\theta. \quad (8.63)$$

One gets after some algebra (for $\zeta \neq 0$)

$$\begin{aligned} \tilde{\underline{P}}(\zeta) = & \left\{ \frac{16|\zeta|^3(\zeta^2 + 1)}{3(2\zeta^2 + 1)(1 + \zeta^2)^{5/2}} \left(\zeta^2 E\left(\frac{|\zeta|}{\sqrt{\zeta^2 + 1}}\right) - (2\zeta^2 + 1)K\left(\frac{|\zeta|}{\sqrt{\zeta^2 + 1}}\right) \right) + \right. \\ & \left. + 8 \ln(|\zeta| \sqrt{\zeta^2 + 1}) + \frac{4(1 + 9\zeta^2 - 8\zeta^4)|\zeta|^3}{3(1 + \zeta^2)^{3/2}} \right\} / (\ln(|\zeta| + \sqrt{\zeta^2 + 1}) + 2\zeta^2 + 1). \end{aligned} \quad (8.64)$$

For $V_2 = \int_{S^2} \sqrt{|\widehat{g}|} dm$ we get

$$V_2 = \frac{2\pi}{|\zeta|} (\ln(|\zeta| + \sqrt{\zeta^2 + 1}) + 2\zeta^2 + 1) \neq 0, \quad (8.65)$$

where

$$K(k) = \int_0^{\pi/2} \frac{d\theta}{\sqrt{1 - k^2 \sin^2 \theta}}, \quad (8.65a)$$

$$E(k) = \int_0^{\pi/2} \sqrt{1 - k^2 \sin^2 \theta} d\theta, \quad 0 \leq k^2 \leq 1 \quad (8.65b)$$

are first and second order elliptic integrals.

It is easy to see that for $\zeta = 0$, $\tilde{\underline{P}}(0) = 0$. The function $\tilde{\underline{P}}$ is plotted on Fig. 6. Let us look for zeros of $\tilde{\underline{P}}$.

$\tilde{\underline{P}}(\zeta_0) = 0$, $\zeta_0 \neq 0$ is a transcendental equation and we should find a solution using numerical methods. One gets $\zeta_0 = \pm 1.36 \dots$ and $\tilde{\underline{P}}(\zeta) \rightarrow -\infty$ if $|\zeta| \rightarrow +\infty$. Let us notice that the integral (8.69) has been calculated under the assumption $\zeta \neq 0$. For $\zeta = 0$ we should do it in a different way. It is easier of course and corresponds to the symmetric connection on S^2 .

Let us consider a more general case given by Eq. (6.49).

One gets in the case of the Kählerian structure.

$$\tilde{\underline{P}} = \frac{1}{V_2} \int_M \widehat{\underline{R}}(\widehat{\Gamma}) \sqrt{|\widehat{g}|} dm(g) = \frac{1}{V_2} \int_M \sqrt{|\widehat{g}|} g^{\tilde{a}\tilde{b}} \tilde{\underline{R}}_{\tilde{a}\tilde{b}}(\tilde{\omega}) dm(g). \quad (8.66)$$




Fig. 6. The function $\tilde{P}$ plotted versus ζ

Moreover $\widehat{\widehat{R}}(\widehat{\widehat{T}})$ is left-invariant with respect to the action of the group G on M . Thus we have

$$\widehat{\widehat{R}}(\widehat{\widehat{T}}(y)) = \widehat{\widehat{R}}(\widehat{\widehat{T}}(gy)), \quad g \in G, \quad y \in M. \quad (8.67)$$

In particular

$$\widehat{\widehat{R}}(\widehat{\widehat{T}}(\{\varepsilon G_0\})) = \widehat{\widehat{R}}(\widehat{\widehat{T}}(y)). \quad (8.68)$$

Thus one gets

$$\tilde{P} = g^{\tilde{a}\tilde{b}} \tilde{\widehat{R}}_{\tilde{a}\tilde{b}}(\tilde{\omega}). \quad (8.69)$$

Let us notice that in the case of S^2 equipped with the nonsymmetric tensor coming from Kählerian structure on S^2 one simply gets (see Eq. (6.60)).

$$P = \frac{4}{(1 + \zeta^2)}. \quad (8.70)$$

Thus for a large ζ we have

$$P \sim \frac{4}{\zeta^2}. \quad (8.71)$$

The interesting point in our theory is a place of electromagnetic field in the theory. Let us suppose that $\mathbf{U}(1)_{el} \subset G$ is a subgroup of the gauge group and let us consider the case when this group is broken to the $\mathbf{U}(1)_{el}$ only. The Higgs field $\Phi_a^a(x)$ in this case takes the special value $(\Phi_{\text{crt}}^0)_a^a$ or $(\Phi'_{\text{crt}})_a^a$ (depending on the appropriate minimum of the Higgs potential $V(\Phi)$). If the Higgs field takes value from the vacuum manifold

G/G_0 (G/G'_0) we can suppose that

$$\overset{\text{gauge}}{\nabla}_\mu \Phi_a^a = 0. \quad (8.72)$$

We can consider a transformation

$$\Sigma(\Phi) \in G/G_0, \quad \Sigma : \tilde{S} \rightarrow G/G_0 \quad (8.73)$$

such that

$$\overset{\text{gauge}}{\nabla}_\mu \hat{\Phi}_a^c(x) = 0. \quad (8.74)$$

Now we consider $G_0 = \mathbf{U}(1)_{\text{el}}$. This is beyond our considerations, because $\mathbf{U}(1)_{\text{el}}$ is not semisimple. However in that case we can reduce the problem to much more simpler one. The Higgs field just belongs to the adjoint representation of the group G and we really have to do with $\phi(x) = \phi^i(x)Y_i$. Thus we can consider a gauge field

$$\tilde{h}_{ij} \phi^i(x) \tilde{\omega}_E^j = \omega_{el}, \quad \overset{\text{gauge}}{\nabla}_\mu \phi = 0, \quad (8.75)$$

which is a one-form with values in R (a $\mathbf{U}(1)$ -group Lie algebra). This form will be considered as an electromagnetic connection. (ϕ belongs to vacuum manifold $G/\mathbf{U}(1)$).

Let us notice that

$$\pi_2(G/\mathbf{U}(1)) = \pi_1(\mathbf{U}(1)) = Z. \quad (8.77)$$

This means that the topological type of the field is characterized by an integer, the topological charge.

Thus we can define a strength $F_{\mu\nu}(x)$ of an electromagnetic field.

$$F_{\mu\nu}(x) = \tilde{h}_{ij} \tilde{H}_{\mu\nu}^i P^j(\Sigma(\phi)), \quad (8.78)$$

where

$$\Sigma : \mathfrak{g} \rightarrow G/\mathbf{U}(1) \quad (8.79)$$

and takes the field ϕ to a vacuum manifolds. P is a generator for the stabilized $\mathbf{U}(1)_\phi$, $\phi \in G/\mathbf{U}(1)$ satisfying the normalization condition:

$$\tilde{h}_{ij} P^i(\phi) P^j(\phi) = 1. \quad (8.80)$$

Moreover, we can compute the electromagnetic field strength in a more general way finding a gauge-invariant expression that coincide with the previous one.

One consider the following expression

$$F_{\mu\nu}(x) = \tilde{h}_{ij} \tilde{H}_{\mu\nu}^i P^j(\Sigma(\phi)) + \varkappa_{ij}(\phi(x)) \overset{\text{gauge}}{\nabla}_\mu \phi^i(x) \overset{\text{gauge}}{\nabla}_\nu \phi^j(x), \quad (8.81)$$

where

$$\varkappa_{ij} = -\varkappa_{ji}. \quad (8.82)$$

This could be rewritten in a different form.

$$F_{\mu\nu}(x) = \tilde{h}_{\hat{i}\hat{j}} \tilde{H}_{\mu\nu}^{\hat{i}} P^{\hat{j}}(\Sigma(\phi)) + \varkappa_{\phi}(\overset{\text{gauge}}{\nabla}_{\mu} \phi, \overset{\text{gauge}}{\nabla}_{\nu} \phi), \quad (8.83)$$

where

$$\varkappa_{\phi}(X, Z) = \varkappa_{\hat{i}\hat{j}}(\phi) X^{\hat{i}} Z^{\hat{j}}, \quad (8.84)$$

$$X = X^{\hat{i}} Y_{\hat{i}}, \quad Z = Z^{\hat{j}} Y_{\hat{j}} \quad (8.85)$$

This form satisfies the following conditions:

$$\varkappa_{\phi}(X, Z) = \varkappa_{\Sigma(\phi)}(\Sigma' X, \Sigma' Z), \quad (8.86)$$

$$\varkappa_{\phi_0}(\text{Ad}_G(A)\phi_0, \text{Ad}_G(B)\phi_0) = -\tilde{h}_{\hat{i}\hat{j}}[A, B]^{\hat{i}} P^{\hat{j}}(\phi_0), \quad (8.87)$$

$A, B \in \mathfrak{g}$ and ϕ_0 is the critical value of the Higgs field. Σ' means a tangent transformation of Σ at ϕ i.e.

$$\Sigma' = \left(\frac{\partial \Sigma^{\hat{i}}(\phi)}{\partial \phi^{\hat{j}}} \right). \quad (8.88)$$

Let us notice that

$$d\omega_{el} = \frac{1}{2} F_{\mu\nu} \theta^{\mu} \wedge \theta^{\nu}$$

In the case of $G = \text{SU}(2)$ one simply gets $\phi(x) \in A_1$, $\phi = \phi^{\hat{i}} X_{\hat{i}}$, and ϕ is a three-dimensional vector in R^3 i.e. $\phi = \overrightarrow{\phi}$ (see the second position of Ref. [17]).

One writes

$$\Sigma(\overrightarrow{\phi}) = a \frac{\overrightarrow{\phi}}{\|\phi\|}, \quad \|\phi\| = \sqrt{\phi_1^2 + \phi_2^2 + \phi_3^2}. \quad (8.89)$$

We also get

$$\overrightarrow{P}(\phi) = \frac{\overrightarrow{\phi}}{\|\phi\|}, \quad (8.90)$$

$$\Sigma'(\phi)_j^i = \frac{a}{\|\phi\|} \left(\delta_j^i - \frac{1}{\|\phi\|^2} \phi^i \phi_j \right) \quad (8.91)$$

and

$$\varkappa_{\phi}(X, Z) = \frac{1}{\|\phi\|^3} \varepsilon_{\hat{i}\hat{j}\hat{k}} \phi^{\hat{i}} X^{\hat{j}} Z^{\hat{k}}. \quad (8.92)$$

Recently have appeared some papers on Higgs'-bosonless Standard Model (see the last two positions of the Ref. [75]). In this approach the remaining Higgs' boson is eliminated from the theory (after spontaneous symmetry breaking) due to conformal invariance of the theory. This interesting approach wants to cure the problem with difficulty in Higgs' boson hunting (maybe because of its very big mass). In our theory we can make the Higgs boson masses very big and we can eliminate one of the remaining

Higgs' boson connecting it to the scalar field Ψ (but only one). This makes our approach closer to the mentioned one.

This can be achieved in the following way. Let us suppose that one of the surviving Higgs' field is φ and let us connect it to the Ψ field by a transformation $\varphi = g(\Psi)$, where g is a smooth function of one variable. In this way the Higgs' field φ disappears from the theory becoming a Ψ field connected to the variable gravitational „constant” (an effective one). If the field φ is the field H^0 from Weinberg–Salam Model, we can remove H^0 from the theory. Let us remind to the reader that the Weinberg–Salam Model (a bosonic part of the model) is covered by the theory (see Ref. [75]). The form of the function g could be established by a requirement that the full lagrangian of Ψ in a linear approximation looks classical.

Let us consider a more general case of symmetry breaking i.e. G to G_0 or to G'_0 corresponding to two possible critical point of the Higgs' potential. In this case we use a smooth transformation

$$\Sigma : \tilde{S} \rightarrow G/G_0 \quad (\text{or } G/G'_0) \quad (8.93)$$

is such a way that

$$\overset{\text{gauge}}{\nabla}_\nu \Phi_a^c = 0. \quad (8.94)$$

Moreover due to decomposition (a reductive one) of the Lie algebra of the group G

$$\mathfrak{g} = \mathfrak{g}_0 \dot{+} \mathfrak{m} = \mathfrak{g}'_0 \dot{+} \mathfrak{m} \quad (8.95)$$

we have a natural parametrization of the homogeneous space G/G_0 (or G/G'_0) in forms of the complement $\mathfrak{m}$ ($\mathfrak{m}'$) in such a way that:

$$(x_5, x_6, \dots, x_{n_1+4}) \rightarrow \varphi \left(g \exp \left(\sum_{\tilde{a}} x_{\tilde{a}} Y_{\tilde{a}} \right) \right), \quad (8.96)$$

which maps a neighbourhood of zero in $\mathfrak{m}(\mathfrak{m}')$ on a neighbourhood of $\varphi(g)$ in G/G_0 . An inverse mapping is a local coordinate system in a neighbourhood of $\varphi(g) \in G/G_0$ on $M = G/G_0$. φ is a natural (canonical) map $G \xrightarrow{\varphi} M = G/G_0$. Let us call this map ψ . One gets

$$\Sigma^a(\Phi) = \psi(\Phi_{\tilde{a}}^a), \quad (8.97)$$

where

$$\psi^{-1}(x_5^a, x_6^a, \dots, x_{n_1+4}^a) = \varphi \left(g \exp \left(\sum_{\tilde{a}} \Phi_{\tilde{a}}^a(x) Y_{\tilde{a}} \right) \right), \quad x \in E, \quad g \in G. \quad (8.98)$$

It is easy to see that for every a , $\Sigma^a(\Phi) \in G/G_0$ (G/G'_0). In this way we do not suppose

$$\overset{\text{gauge}}{\nabla}_\mu \Phi_{\tilde{a}}^a = 0. \quad (8.99)$$

Now we can construct a gauge covariant form of the gauge strength

$$\tilde{H}_{\mu\nu}^{\hat{i}} = h_{ij} \tilde{H}_{\mu\nu}^i P^j(\Sigma(\Phi) \mu_a^{\hat{i}}) + \varkappa^{\tilde{a}\tilde{b}}(\Phi(x)) [(\overset{\text{gauge}}{\nabla}_\mu \Phi_a^{\tilde{a}})(\overset{\text{gauge}}{\nabla}_\nu \Phi_b^{\tilde{b}}) \mu_a^{\hat{k}} \mu_b^{\hat{j}}] f_{\hat{k}\hat{j}}^{\hat{i}}, \quad (8.100)$$

where $P^j(\Phi^a)$ are generators for the stabilizer of $(G_0)_{\Phi^a}$, $\Phi^a \in G/G_0$ (the same for G'_0 case). Satisfying the normalization conditions

$$\|P^j(\Phi^a)\| = 1. \quad (8.101)$$

This is the gauge field strength which remains massless after the symmetry breaking.

If we take under consideration

$$\pi_2(G/G_0) = \pi_1(G_0), \quad (8.102)$$

(the same for the G'_0 case) we can consider a nontrivial topological configuration of the gauge field i.e. the so called „coloured” magnetic monopoles. The gauge field strength defined above is simply the curvature of the connection ω^0_E (or $\omega^{0'}_E$).

$$\overset{\text{gauge}}{D} \tilde{\omega}^0_E = \frac{1}{2} \tilde{H}_{\mu\nu}^{\hat{i}} \theta_{\wedge}^{\mu} \theta^{\nu} Y_{\hat{i}} \quad \text{or} \quad \overset{\text{gauge}}{D} \tilde{\omega}^{0'}_E = \frac{1}{2} \tilde{H}_{\mu\nu}^{\hat{i}} \theta_{\wedge}^{\mu} \theta^{\nu} Y_{\hat{i}}' \\ \left(\overset{\text{gauge}}{D} \tilde{\omega}^0_E = d\tilde{\omega}^0_E + \frac{1}{2} [\omega^0_E, \omega^0_E] \right).$$

and analogically for $\tilde{\omega}^{0'}_E$.

The form $\varkappa^{\tilde{a}\tilde{b}}(\Phi(x))$ satisfies the following conditions:

$$\varkappa^{\tilde{a}\tilde{b}} = \varkappa^{\tilde{b}\tilde{a}}. \quad (8.103)$$

If we define a form

$$\varkappa(\Phi(x))(X, Z) = \sum_{\tilde{a}, \tilde{b}} \varkappa^{\tilde{a}\tilde{b}}(\Phi(x)) X^{\tilde{a}} Z^{\tilde{b}} \quad (8.104)$$

the formula for the strength of the gauge field can be rewritten as follows:

$$\tilde{H}_{\mu\nu}^{\hat{i}}(x) = \langle \tilde{H}_{\mu\nu}(x), P(\Sigma^a(\Phi) \mu_a^{\hat{i}}) \rangle + \\ + \varkappa(\Phi(x)) (\mu_a^{\hat{k}} \overset{\text{gauge}}{\nabla}_\mu \Phi_a^{\tilde{a}} \mu_b^{\hat{j}} \overset{\text{gauge}}{\nabla}_\nu \Phi_b^{\tilde{b}}) f_{\hat{k}\hat{j}}^{\hat{i}}. \quad (8.105)$$

The form $\varkappa(\Phi)$ can be found from the requirements:

$$\varkappa(\Phi)(X, Z) = \varkappa(\Sigma\Phi)(\Sigma'X, \Sigma'Y), \quad (8.106)$$

where $\Sigma' = (\Sigma^a)' = \left(\frac{\partial \Sigma^a}{\partial \Phi_a^b} \right)$ is tangent map to Σ

$$\varkappa(\Phi_{\text{crt}}^k)(\text{Ad}_G(A)_{\hat{k}}^{\hat{k}'} \mu_a^{\hat{k}} \text{Ad}_G(A)_{\tilde{a}}^{\tilde{a}'} (\Phi_{\text{crt}}^k)_{\tilde{a}'}^a, \text{Ad}_G(B)_{\hat{j}}^{\hat{j}'} \mu_b^{\hat{j}} \text{Ad}_G(B)_{\tilde{b}}^{\tilde{b}'} (\Phi_{\text{crt}}^k)_{\tilde{b}'}^b) = \\ -\langle [A, B], p^i(\Phi_{\text{crt}}^k) \rangle, \quad k = 0, 1, \quad (8.108)$$

where $\langle X, Y \rangle = h_{ij} X^i X^j$ and $A, B, X, Y \in \mathfrak{g}$.

We can consider in our theory topologically nontrivial field configurations labeled by $\pi_2(G/G_0)$ or $\pi_2(G/G'_0)$. They are „coloured” magnetic charges. In the case of $G_0 = \mathbf{U}(1)$ we have to do with ordinary magnetic charges.

Let us define a magnetic and a „coloured” magnetic field

$$\vec{H}(\vec{x}) = (F_{23}(\vec{x}), F_{31}(\vec{x}), F_{12}(\vec{x})), \quad (8.109)$$

$$\vec{H}^{\hat{i}}(\vec{x}) = (F_{23}^{\hat{i}}(\vec{x}), F_{31}^{\hat{i}}(\vec{x}), F_{12}^{\hat{i}}(\vec{x})) \quad (8.110)$$

and magnetic and coloured magnetic charge (monopole):

$$m = \frac{1}{4\pi} \oint \vec{H}(\vec{x}) d\vec{S}, \quad (8.111)$$

$$m^{\hat{i}} = \frac{1}{4\pi} \oint \vec{H}^{\hat{i}}(\vec{x}) d\vec{S}, \quad (8.112)$$

where the integrals are taken over a sphere infinitely far away i.e. for a sphere at infinity. In this case we suppose that

$$\begin{aligned} \Phi_a^a(\vec{x}) &\in G/G_0 \quad \text{if} \quad \|\vec{x}\| \rightarrow \infty \quad \text{and} \\ \nabla_\mu^{\text{gauge}} \Phi_a^a(\vec{x}) &\rightarrow 0, \quad \text{if} \quad \|\vec{x}\| \rightarrow \infty, \end{aligned}$$

where $\|\vec{x}\| = \sqrt{|x_1|^2 + |x_2|^2 + |x_3|^2}$.

For example we can get a nontrivial topological gauge configurations for the following classical Lie groups

$$\begin{aligned} \text{SO}(n), \pi_1(\text{SO}(n)) &= \mathbb{Z}_2, \quad \text{for } n > 1, \\ \mathbf{U}(n), \pi_1(\mathbf{U}(n)) &= \mathbb{Z}, \quad \text{for } n \geq 1. \end{aligned}$$

In the case of $\text{SU}(n)$, $\text{Sp}(n)$, and $\text{Spin}(n)$, $n \geq 1$ we get trivial gauge configurations, became

$$\pi_1(\text{SU}(n)) = \pi_1(\text{Spin}(n)) = \pi_1(\text{Sp}(n)) = 0.$$

Thus we can expect „coloured” magnetic charges (monopoles) for $\text{SO}(n+1)$ and $\mathbf{U}(n)$, $n \geq 1$ (equal to G_0 or G'_0).

Finally let us remind to the reader that in the case of G_0 group (a „true” vacuum case) we have to do with a trivial gauge configuration of the Higgs’ field $\Phi_a^a(x)$ (i.e. Φ_{crt}^0 case). In this case a part of a strength of the multi-dimensional gauge field connected to Higgs’ field **does vanish**. In the case of G_0 -group (a „false” vacuum care) this part of the strength of the multidimensional gauge field **does not vanish**. Thus the „magnetic” coloured charges (monopoles) in both cases have different physical meanings. In some sense in the second care they corresponds to something which could be called „exited” magnetic monopoles (coloured ones) i.e. for Φ_{crt}^1 case.

It is worth to say we suppose now that G_0 or G'_0 is simply connected. The topological (magnetic) charges depend only on the configurations of Higgs’ fields $\Phi_a^a(\vec{x})$

and does not depend on the gauge field $\tilde{\omega}^0_E$ or $\tilde{\omega}^{0'}_E$ ($\tilde{\omega}_{el}$ in the case of electromagnetic field).

We defined the topological type of the field by looking of the asymptotic behaviour of the scalar field Φ . Thus the topological type of the field is an element of $\{S^2, G/G_0\}$ that is a homotopy class of mapping from S^2 into the vacuum manifold. When G_0 is simply connected, the set $\{S^2, G/G_0\}$ can be identified with $\pi_2(G/G_0) = \pi_1(G_0)$, which is a finite group.

Thus there are r topological charges, as many as there are magnetic charges. Clearly, the magnetic charges completely determine the integral values of the topological charges of a field. In addition to integral topological charges, there may be charges taking values in a finite group $\mathbb{Z}_m$ (i.e. $\mathbb{Z}_2$), these do not correspond to magnetic charges. They are coloured magnetic charges (monopoles). In general we can consider topological charges for $\tilde{H}^i_{\mu\nu}$ field

$$\chi(\tilde{\omega}_E) = \frac{1}{16\pi^2} \int_E \langle \tilde{H} \wedge \tilde{H} \rangle, \quad (8.113)$$

where

$$\langle \tilde{H} \wedge \tilde{H} \rangle = -\frac{1}{2} \tilde{h}_{ij} \tilde{H}^i_{\alpha\beta} \tilde{H}^j_{\gamma\delta} \tilde{\theta}^d_{\wedge} \tilde{\theta}^p_{\wedge} \tilde{\theta}^\gamma_{\wedge} \tilde{\theta}^\delta_{\wedge}$$

and similarly for $\tilde{H}^{\hat{i}}_{\mu\nu}$

$$\hat{\chi}(\tilde{\omega}^0_E) = \frac{1}{16\pi^2} \int_E \langle \hat{\tilde{H}} \wedge \hat{\tilde{H}} \rangle, \quad (8.114)$$

where

$$\langle \hat{\tilde{H}} \wedge \hat{\tilde{H}} \rangle = -\frac{1}{2} \tilde{h}_{\hat{i}\hat{j}} \hat{\tilde{H}}^{\hat{i}}_{\alpha\beta} \hat{\tilde{H}}^{\hat{j}}_{\gamma\delta} \tilde{\theta}^\alpha_{\wedge} \tilde{\theta}^\beta_{\wedge} \tilde{\theta}^\gamma_{\wedge} \tilde{\theta}^\delta_{\wedge}.$$

The topological number of a field $\tilde{\omega}_E$ (or $\tilde{\omega}^0_E$) on E can be defined when the strength of the fields $\tilde{H}^{\hat{i}}_{\alpha\beta}$ or $\tilde{H}^i_{\alpha\beta}$ tends to zero sufficiently fast as $\|x\| \rightarrow \infty$, $\|x\|^2 = x_1^2 + x_2^2 + x_3^2 + x_4^2$ (In this case we take $E = R^4$). If the field $\tilde{H}$ decays quickly $\tilde{\omega}_E$ coincides asymptotically with a pure gauge field, that is it has asymptotics of $g^{-1} dg$, where $g(x)$ is a G -valued function. In that case we have

$$\begin{aligned} \chi(\tilde{\omega}_E) &= -\frac{1}{96\pi^2} \int_{S^3} \langle \tilde{\omega}_E \wedge [\tilde{\omega}_E, \tilde{\omega}_E] \rangle \\ &= -\frac{1}{96\pi^2} \int_{S^3} \langle g^{-1} dg \wedge [g^{-1} dg, g^{-1} dg] \rangle, \end{aligned} \quad (8.115)$$

where S^3 is taken as a sphere at infinity. In this case the topological charges are classified by $\pi_3(G)$, $\pi_3(G_0)$ or $\pi_3(G'_0)$.

9. Variational Principle. Equations of Fields. Interpretation of the Scalar Field Ψ

Let us consider Palatini variational principle for the action S .

$$\delta S = 0. \quad (9.1)$$

It is easy to see that (9.1) is equivalent to

$$\delta \int_{\mathbf{U}} L(g, \bar{W}, \tilde{A}, \Psi, \Phi) \sqrt{-g} d^4x = 0, \quad \mathbf{U} \subset E. \quad (9.2)$$

We have the following independent quantities $g_{\mu\nu}$, $\bar{W}^\lambda_{\mu\nu}$, $\tilde{\omega}_E$, Ψ and Φ . We vary with respect to the independent quantities. After some calculations one gets:

$$\bar{R}_{\mu\nu}(\bar{W}) - \frac{1}{2}g_{\mu\nu}\bar{R}(\bar{W}) = \frac{8\pi K}{c^4} (T_{\mu\nu}^{\text{gauge}} + T_{\mu\nu}(\Phi) + T_{\mu\nu}^{\text{scal}}(\Psi) + T_{\mu\nu}^{\text{int}} + g_{\mu\nu}\Lambda), \quad (9.3)$$

$$\tilde{g}^{[\mu\nu]}_{,\nu} = 0, \quad (9.4)$$

$$\bar{\nabla}_\nu g^{[\mu\nu]} = 0, \quad (9.4a)$$

$$g_{\mu\nu,\sigma} - g_{\xi\nu}\bar{\Gamma}^\xi_{\mu\sigma} - g_{\mu\xi}\bar{\Gamma}^\xi_{\sigma\nu} = 0, \quad (9.5)$$

$$\begin{aligned} & ((n^2 + 2M)\tilde{g}^{(\alpha\mu)} - n^2 g^{\nu\mu} g_{\delta\nu} \tilde{g}^{(\alpha\delta)}) \frac{\partial^2 \Psi}{\partial x^\alpha \partial x^\mu} + \\ & + \frac{1}{\sqrt{-g}} \partial_\mu \left\{ \sqrt{-g} \left[n^2 \tilde{g}^{(\alpha\mu)} - \frac{n^2}{2} g_{\delta\nu} (g^{\nu\alpha} \tilde{g}^{(\mu\delta)} + g^{\nu\mu} \tilde{g}^{(\mu\alpha)}) - 2M \tilde{g}^{(\mu\alpha)} \right] \right\} \times \\ & \times \frac{\partial \Psi}{\partial x^\alpha} - 8\pi(n+2)e^{-(n+2)\Psi} \mathcal{L}_{\text{YM}}(\tilde{A}) - \frac{4e^{-2\Psi}}{r^2} \mathcal{L}_{\text{kin}}(\Phi, \tilde{A}) + \\ & + \frac{(n-2)}{r^4} e^{(n-2)\Psi} V(\Phi) + \frac{4(n-2)}{r^2} e^{(n-2)\Psi} \mathcal{L}_{\text{int}}(\Phi, \tilde{A}) + \\ & - \frac{n}{r^2} e^{n\Psi} \tilde{P} - \frac{(n+2)\alpha_S^2}{\ell_{pl}^2} e^{(n+2)\Psi} \tilde{R}(\tilde{\Gamma}) = 0, \end{aligned} \quad (9.6)$$

$$\begin{aligned}
\text{gauge} \nabla_\mu (\tilde{\ell}_{ij} \tilde{L}^{i\alpha\mu}) &= 2 g^{[\alpha\beta]} \text{gauge} \nabla_\beta (\tilde{h}_{ij} g^{[\mu\nu]} \tilde{H}^i_{\mu\nu}) + \\
&+ 2\sqrt{-g} \alpha_S \frac{1}{\sqrt{\hbar c}} \frac{e^{n\Psi}}{r^2} \left[\ell_{ab} g^{\tilde{b}\tilde{n}} g^{\mu\alpha} L^a_{\mu\tilde{b}} (\Phi^d_{\tilde{c}} C^b_{dc} \alpha^c_j + \Phi^b_{\tilde{a}} f^{\tilde{a}}_{\tilde{n}j}) + \right. \\
&+ \left. \left(\frac{\delta L^a_{\beta\tilde{b}}}{\delta \nabla_\alpha \Phi^w_{\tilde{v}}} \right) \ell_{ab} g^{\tilde{b}\tilde{n}} g^{\beta\mu} (\text{gauge} \nabla_\mu \Phi^b_{\tilde{n}}) (\Phi^d_{\tilde{w}} C^w_{dc} \alpha^c_j + \Phi^w_{\tilde{a}} f^{\tilde{a}}_{\tilde{n}j}) \right]_{av} + \\
&+ 4\sqrt{-g} \frac{e^{2n\Psi}}{r^2} h_{ab} \mu^a_k \tilde{\ell}_{ij} \ell^{ki} \tilde{g}^{[\tilde{a}\tilde{b}]} \text{gauge} \nabla_\mu \left\{ g^{[\mu\alpha]} \times \right. \\
&\times \left[\frac{1}{\alpha_S} \sqrt{\hbar c} C^b_{cd} \Phi^c_{\tilde{a}} \Phi^d_{\tilde{b}} - \alpha_S \left(\frac{\hbar}{c} \mu^b_i \hat{f}^i_{\tilde{a}\tilde{b}} - \alpha_S \frac{1}{\sqrt{\hbar c}} \Phi^b_{\tilde{d}} f^{\tilde{d}}_{\tilde{a}\tilde{b}} \right) \right] \Big\} + \\
&+ (n+2) \partial_\beta \Psi [\tilde{\ell}_{ij} \tilde{L}^{i\beta\alpha} - 2 g^{[\mu\nu]} (\tilde{h}_{ij} g^{[\mu\nu]} \tilde{H}^i_{\mu\nu})], \tag{9.7}
\end{aligned}$$

$$\begin{aligned}
\text{gauge} \nabla_\mu (\ell_{ab} \tilde{L}^{a\mu}_{\tilde{b}})_{av} &= -\sqrt{-g} \frac{e^{n\Psi}}{2r^2} \left\{ \left(\frac{\delta V'}{\delta \Phi^b_{\tilde{n}}} \right) g_{\tilde{b}\tilde{n}} + \right. \\
&- 2\sqrt{-g} e^{n\Psi} \mu^e_i (\tilde{H}^i_{\mu\nu} g^{[\mu\nu]}) h_{ed} \left(\frac{2}{\alpha_S} \sqrt{\hbar c} \underline{g}^{[\tilde{a}\tilde{n}]} C^d_{cb} \Phi^c_{\tilde{a}} g_{\tilde{b}\tilde{n}} + \right. \\
&- \left. \left. \alpha_S \frac{1}{\sqrt{\hbar c}} g^{[\tilde{c}\tilde{d}]} f^{\tilde{n}}_{\tilde{c}\tilde{d}} g_{\tilde{b}\tilde{n}} \right) + 2\partial_\mu \Psi \ell_{ab} \tilde{L}^{a\mu}_{\tilde{b}} \right\}_{av}, \tag{9.8}
\end{aligned}$$

where

$$\begin{aligned}
\text{gauge} T_{\alpha\beta} &= -\frac{\tilde{\ell}_{ij}}{4\pi} \left\{ g_{\gamma\beta} g^{\tau\rho} g^{\varepsilon\gamma} \tilde{L}^i_{\rho\alpha} \tilde{L}^j_{\tau\varepsilon} - 2g^{[\mu\nu]} \tilde{H}^{(i}_{\mu\nu} \tilde{H}^{j)}_{\alpha\beta} + \right. \\
&- \left. \frac{1}{4} g_{\alpha\beta} [\tilde{L}^{i\mu\nu} \tilde{H}^j_{\mu\nu} - 2(g^{[\mu\nu]} \tilde{H}^i_{\mu\nu})(g^{[\gamma\sigma]} \tilde{H}^j_{\gamma\sigma})] \right\} \tag{9.9}
\end{aligned}$$

is the energy-momentum tensor for the gauge (Yang–Mills') field with the zero trace.

$$\text{gauge} T_{\alpha\beta} g^{\alpha\beta} = 0, \tag{9.10}$$

$$\begin{aligned}
\text{scal} T_{\alpha\beta}(\Psi) &= -\frac{e^{(n+2)\Psi}}{16\pi} \left\{ (g_{\varkappa\alpha} g_{\omega\beta} + g_{\omega\alpha} g_{\varkappa\beta}) \times \right. \\
&\times \tilde{g}^{(\gamma\varkappa)} \tilde{g}^{(\nu\omega)} \cdot \left[\frac{n^2}{2} (g^{\xi\mu} g_{\nu\xi} - \delta^\mu_\nu) \Psi_{,\mu} + \overline{M} \Psi_{,\nu} \right] \Psi_{,\gamma} + \\
&- \left. g_{\alpha\beta} [\overline{M} \tilde{g}^{(\nu\mu)} + n^2 g^{[\mu\nu]} g_{\delta\mu} \tilde{g}^{(\gamma\delta)} \Psi_{,\nu} \Psi_{,\gamma}] \right\} \tag{9.11}
\end{aligned}$$

is the energy-momentum tensor for the scalar field Ψ with nonzero trace

$$T_{\alpha\beta}^{\text{scal}}(\Psi)g^{\alpha\beta} \neq 0 \quad (9.12)$$

$$K = G_N e^{-(n+2)\Psi} = G_{\text{eff}}(\Psi). \quad (9.13)$$

It plays the role of an effective gravitational constant.

$$T_{\mu\nu}(\Phi) = \frac{e^{(n-4)\Psi}}{4\pi r^2} \ell_{ab} g_{\tilde{b}\tilde{n}} L^a_{\mu\tilde{b}} \overset{\text{gauge}}{\nabla}_\nu \Phi^b_{\tilde{n}} + \\ - \frac{1}{2} g_{\mu\nu} \left(-\frac{e^{2(n-2)\Psi}}{8\pi r^4} V(\Phi) + \frac{e^{(n-4)\Psi}}{4\pi r^2} \ell_{ab} (g^{\tilde{b}\tilde{n}} g^{\alpha\beta} L^a_{\alpha\tilde{b}} \overset{\text{gauge}}{\nabla}_\beta \Phi^b_{\tilde{n}})_{av} \right). \quad (9.14)$$

It is an energy-momentum tensor for the Higgs' field.

$$T_{\mu\nu}^{\text{int}} = -\frac{e^{2(n-2)\Psi}}{2\pi r^2} h_{ab} \mu_i^a \tilde{H}^i_{\mu\nu} \tilde{g}^{[\tilde{a}\tilde{b}]} \left(\frac{1}{\alpha_S} \sqrt{\hbar c} C_{cd}^b \Phi^c_{\tilde{a}} \Phi^d_{\tilde{b}} + \right. \\ \left. - \alpha_S \frac{1}{\sqrt{\hbar c}} \mu_i^b f^{\hat{i}}_{\tilde{a}\tilde{b}} - \alpha_S \frac{1}{\sqrt{\hbar c}} \Phi^b_{\tilde{d}} f^{\tilde{d}}_{\tilde{a}\tilde{b}} \right) + \\ + \frac{e^{2(n-2)\Psi}}{4\pi r^2} g_{\mu\nu} \left[h_{ab} \mu_i^a (\tilde{H}^i_{\alpha\beta} g^{[\alpha\beta]}) \tilde{g}^{[\tilde{a}\tilde{b}]} \times \right. \\ \left. \times \left(\frac{1}{\alpha_S} \sqrt{\hbar c} C_{cd}^b \Phi^c_{\tilde{a}} \Phi^d_{\tilde{b}} - \alpha_S \frac{\hbar}{c} \mu_i^b f^{\hat{i}}_{\tilde{a}\tilde{b}} - \alpha_S \frac{1}{\sqrt{\hbar c}} \Phi^b_{\tilde{d}} f^{\tilde{d}}_{\tilde{a}\tilde{b}} \right) \right]. \quad (9.15)$$

It is an energy-momentum tensor corresponding to the nonminimal interaction term $\mathcal{L}_{\text{int}}(\tilde{A}, \Phi)$.

$$\Lambda = \frac{1}{16\pi G_N} \left(\frac{e^{(2n+4)\Psi} \alpha_S^2}{\ell_{pl}^2} \tilde{R}(\tilde{\Gamma}) + \frac{e^{(n+2)\Psi}}{r^2} \tilde{P} \right) = 16\pi G_N e^{-(n+2)\Psi} \bar{\lambda}_c. \quad (9.16)$$

It plays the role of the “cosmological constant”, which now depends on the scalar field

Ψ . The quantity $\frac{\delta L^a_{\beta\tilde{b}}}{\delta \overset{\text{gauge}}{\nabla}_\alpha \Phi^w_{\tilde{v}}}$ satisfies the following equation:

$$\ell_{dc} g_{\mu\beta} g^{\gamma\mu} \frac{\delta L^d_{\gamma\tilde{a}}}{\delta \overset{\text{gauge}}{\nabla}_\alpha \Phi^w_{\tilde{v}}} + \ell_{cd} g_{\tilde{a}\tilde{m}} g^{\tilde{m}\tilde{c}} \frac{\delta L^d_{\beta\tilde{c}}}{\delta \overset{\text{gauge}}{\nabla}_\alpha \Phi^w_{\tilde{v}}} = 2\ell_{cd} g_{\tilde{a}\tilde{m}} g^{\tilde{m}\tilde{c}} \delta^\alpha_\beta \delta^{\tilde{v}}_{\tilde{c}} \delta^d_w, \quad (9.17)$$

$$\tilde{\tilde{L}}^{i\mu\nu} = \sqrt{-g} g^{\beta\mu} g^{\gamma\nu} \tilde{L}^i_{\beta\gamma}, \quad (9.18)$$

$$\tilde{\tilde{g}}^{[\mu\nu]} = \sqrt{-g} g^{[\mu\nu]}, \quad (9.18a)$$

$\tilde{\tilde{L}}^i_{\beta\gamma}$ satisfies equation (9.1) and $L^a_{\beta\tilde{b}}$ the equation (9.4) $\frac{\delta V'}{\delta \Phi^b_{\tilde{n}}}$ is defined by the equation (8.10). The scalar field $\Phi^c_{\tilde{a}}$ satisfies constraints (8.8). Let us pass to the interpretation of the field equations (9.3–8). Equations (9.3), (9.4) and (9.5) are gravitational equations

from N.G.T. with the following matter sources: Yang–Mills’ field (in the nonsymmetric version), Higgs’ field, scalar field Ψ with a presence of the cosmological term depending on scalar field Ψ . Equation (9.6) is the equation for the scalar field Ψ . This field is of course chargeless, but it interacts with Yang–Mills’ field and Higgs’ field due to some terms in (9.6). It interacts also with cosmological terms, which effectively depend on Ψ . This field due to equation (9.13) has an interpretation as an effective gravitational constant. Equation (9.7) is the equation for Yang–Mills’ field. Now as in the Nonsymmetric Kaluza–Klein Theory we have for this field two tensors of strength $\tilde{H}_{\mu\nu}$ and $\tilde{L}^i_{\mu\nu}$ (ordinary and an induction one) and the nonsymmetric parts of metrics on E and on G induce the polarization $\tilde{M}^i_{\mu\nu}$. In the equation (9.7) we have sources connected to a skewsymmetric part of metric $g_{\mu\nu}$ and to Higgs’ field. Due to the existence of a skewsymmetric part of metric ℓ_{ab} and $g_{\tilde{a}\tilde{b}}$ the current connected to Higgs’ field is more complicated. Equation (9.8) is an equation for Higgs’ field. We write this equation in terms of tensor

$$L^{a\mu}_{\tilde{b}} = g^{\alpha\mu} L^a_{\alpha\tilde{b}}. \quad (9.19)$$

This tensor plays a similar role for $\overset{\text{gauge}}{\nabla}_\alpha \Phi^a_{\tilde{b}}$ as $\tilde{L}^{i\mu}_\alpha$ for $\tilde{H}^i_{\mu\alpha}$. Thus we have in the theory an analogue of the polarization tensor $M^a_{\alpha\tilde{b}}$ for Higgs’ field

$$L^a_{\alpha\tilde{b}} = \overset{\text{gauge}}{\nabla}_\alpha \Phi^a_{\tilde{b}} - \frac{4\pi}{c} M^a_{\alpha\tilde{b}}. \quad (9.20)$$

Let us consider the following two Ad_H - type two-forms $\hat{L} = L^a_{\alpha\tilde{b}} \theta^\alpha \wedge \theta^{\tilde{b}} X_a$ and $\widehat{M} = M^a_{\alpha\tilde{b}} \theta^\alpha \wedge \theta^{\tilde{b}} X_a$. One gets $\hat{L} = \hat{\Omega} - \frac{4\pi}{c} \widehat{M} = \hat{\Omega} - \frac{1}{2} \hat{Q}$, where

$$\hat{Q} = \overset{\text{gauge}}{\nabla}_\alpha \Phi^a_{\tilde{b}} \theta^\alpha \wedge \theta^{\tilde{b}} X_a, \quad \hat{Q} = Q^a_{\alpha\tilde{b}} \theta^\alpha \wedge \theta^{\tilde{b}} X_a.$$

In this way we get a geometrical interpretation of the polarization 2-form of the Higgs’ field as a part of a torsion.

The field Ψ due to (9.13) is connected to the effective gravitational constant. However, it also enters the definition of energy-momentum tensors $T_{\mu\nu}(\Phi)$ and $\overset{\text{int}}{T}_{\mu\nu}$. Thus it plays the role of the universal factor. In the next section we deal with this field in details.

Let us notice that the left-hand side of Eq. (9.7) can be rewritten in the following way

$$\overset{\text{gauge}}{\nabla}_\mu (\tilde{l}_{ij} \tilde{L}^{i\alpha\mu}) = \sqrt{-g} \overset{\text{gauge}}{\tilde{\nabla}}_\mu (\tilde{l}_{ij} \tilde{L}^{i\alpha\mu}),$$

where $\overset{\text{gauge}}{\tilde{\nabla}}_\mu$ means a covariant derivative with respect to the connection $\tilde{\omega}^\alpha_\beta$ on E and ω_E on $Q(E, G)$ at once. The Eqs. (9.4–4a) are equivalent to the second condition of Eq. (5.5).

Let us consider a case with a trivial group $G_0 = \{\varepsilon\}$. This means a complete broken G symmetry group and $M = G/G_0 \simeq G$, $V \simeq E \times G$. Thus we get that the Higgs' field Φ_i^c transforms according to the adjoint representation of the group H and G and it does not obey any constraints. They are satisfied trivially. Thus it transforms according to $\text{Ad}_H \otimes \text{Ad}_G$. If we choose $H = G$ one gets $\text{Ad}_G \otimes \text{Ad}_G$.

The Higgs' potential looks like:

$$V(\Phi) = \frac{\ell_{ab}}{V_G} \int_G \sqrt{|\tilde{\ell}|} d^{n_2} x [2\tilde{\ell}^{[ij]} (C_{cd}^a \Phi_i^c \Phi_j^d - \Phi_k^a f_{ij}^k) \tilde{\ell}^{[mn]} (C_{ef}^b \Phi_m^e \Phi_n^f - \Phi_p^b f_{mn}^p) + \\ - \tilde{\ell}^{im} \tilde{\ell}^{jn} L^a_{ij} (C_{cd}^b \Phi_m^c \Phi_n^d - \Phi_e^b f_{mn}^e)]. \quad (9.21)$$

The kinetic part of the Higgs' field lagrangian is as follows:

$$\mathcal{L}_{\text{kin}}(\overset{\text{gauge}}{\nabla} \Phi) = \frac{1}{V_G} \int_G \sqrt{|\tilde{\ell}|} d^{n_2} x (\ell_{ab} \tilde{\ell}^{bn} L^{a\mu g}_b \overset{\text{gauge}}{\nabla}_\mu \Phi_n^b), \quad (9.22)$$

where

$$\ell_{dc} g_{\mu\beta} g^{\gamma\mu} L^d_{\gamma a} + \ell_{cd} \tilde{\ell}_{am} \tilde{\ell}^{mp} L^d_{\beta p} = 2\ell_{cd} \tilde{\ell}_{am} \tilde{\ell}^{mp} \overset{\text{gauge}}{\nabla}_\mu \Phi_p^d \quad (9.23)$$

and

$$\ell_{dc} \tilde{\ell}_{mb} \tilde{\ell}^{pm} L^d_{pa} + \ell_{cd} \tilde{\ell}_{am} \tilde{\ell}^{mp} L^d_{bp} = 2\ell_{cd} \tilde{\ell}_{am} \tilde{\ell}^{mp} H^d_{bp}. \quad (9.24)$$

For H^d_{bp} one gets:

$$H^d_{bp} = C_{ac}^d \Phi_b^a \Phi_p^c - \Phi_q^d f_{bp}^q. \quad (9.24a)$$

In the above formulae $\tilde{\ell}_{ij}$ is the same as in sec. 6.

The nonsymmetric tensor on G is left-invariant and on H is right-invariant.

The interaction term in the lagrangian is as follows:

$$\mathcal{L}_{\text{int}}(\Phi, \tilde{A}) = h_{ab} \mu_i^a \tilde{H}^i \int_G \frac{\sqrt{|\tilde{\ell}|}}{V_G} d^{n_2} x \tilde{\ell}^{[pq]} \cdot (C_{cd}^b \Phi_p^c \Phi_q^d - \Phi_e^b f_{pq}^e). \quad (9.25)$$

The cosmological term takes the shape:

$$\lambda_c = \frac{e^{(n+2)\Psi} \alpha_S^2}{\ell_{pl}^2} \tilde{R}(\tilde{\Gamma}) + \frac{e^{n\Psi}}{r^2} R_G, \quad (9.26)$$

where R_G is the scalar curvature of the connection defined on the group G . In order to vanish it we take $\xi = \xi_0$ and $\zeta = \zeta_0$ such that $\tilde{R}(\tilde{\Gamma}(\xi_0)) = 0$ and $R(\zeta_0) = 0$.

$$V_G = \int_G \sqrt{|\tilde{\ell}|} d\mu_G(g) = \int_G \sqrt{|\tilde{\ell}|} d^{n_2} x. \quad (9.27)$$

The dimensional reduction procedure can be described in terms of Ω , Q (see 5.56–58).

In this way Ω decomposes into $\widetilde{\Omega}_E$, $\overset{\vee}{\Omega}$, $\widehat{\Omega}$, and $\overline{L}$ into $\widetilde{L}$, $\widehat{L}$, $\overset{\vee}{L}$ (see paragraphs below formula (6.3), below (6.7) and below (9.20)). Similarly we get a decomposition for $\overline{M}$ and $\overline{Q}$ into $\widetilde{M}$, $\widehat{M}$, $\overset{\vee}{M}$ and into $\widetilde{Q}$, $\widehat{Q}$, $\overset{\vee}{Q}$.

According to Eqs. (4.53–55) one gets for $\overline{L}$, $\overline{M}$ and $\overline{Q}$

$$\begin{aligned}\overline{L} &= \widetilde{L} \oplus \widehat{L} \oplus \overset{\vee}{L}, \\ \overline{M} &= \widetilde{M} \oplus \widehat{M} \oplus \overset{\vee}{M}, \\ \overline{Q} &= \widetilde{Q} \oplus \widehat{Q} \oplus \overset{\vee}{Q},\end{aligned}\tag{9.28}$$

similarly as for Ω

$$\Omega = \widetilde{\Omega} \oplus \widehat{\Omega} \oplus \overset{\vee}{\Omega}\tag{9.29}$$

(see Eq. (4.52)), where

$$\Omega_E = d\omega_E + \frac{1}{2}[\omega_E, \omega_E].\tag{9.30}$$

Let us incorporate in our scheme a hierarchy of a symmetry breaking. In order to do this let us consider a case of the manifold

$$M = M_0 \times M_1 \times \dots \times M_{k-1}\tag{9.31}$$

where

$$\begin{aligned}\dim M_i &= \overline{n}_i, \quad i = 0, 1, 2, \dots, k-1 \\ \dim M &= \sum_{i=0}^{k-1} \overline{n}_i,\end{aligned}\tag{9.32}$$

$$M_i = G_{i+1}/G_i.\tag{9.33}$$

Every manifold M_i is a manifold of vacuum states if the symmetry is breaking from G_{i+1} to G_i , $G_k = G$.

Thus

$$G_0 \subset G_1 \subset G_2 \subset \dots \subset G_k = G.\tag{9.34}$$

We will consider the situation when

$$M \simeq G/G_0.\tag{9.35}$$

This is a constraint in the theory. From the chain (9.34) one gets

$$\mathfrak{g}_0 \subset \mathfrak{g}_1 \subset \dots \subset \mathfrak{g}_k = \mathfrak{g}\tag{9.36}$$

and

$$\mathfrak{g}_{i+1} = \mathfrak{g}_i \dot{+} \mathfrak{m}_i, \quad i = 0, 1, \dots, k.\tag{9.37}$$

The relation (9.35) means that there is a diffeomorphism g onto G/G_0 such that

$$g : \prod_{i=0}^{k-1} (G_{i+1}/G_i) \rightarrow G/G_0. \quad (9.38)$$

This diffeomorphism is a deformation of a product (9.31) in G/G_0 . The theory has been constructed for the case considered before with G_0 and G . The multiplet of Higgs' fields Φ breaks the symmetry from G to G_0 (equivalently from G to G'_0 in the false vacuum case). $\mathfrak{g}_i$ mean Lie algebras for groups G_i and $\mathfrak{m}_i$ a complement in a decomposition (9.37). On every manifold M_i we introduce a radius r_i (a “size” of a manifold) in such a way that $r_i > r_{i+1}$. On the manifold G/G_0 we define the radius r as before. The diffeomorphism g induces a contragradient transformation for a Higgs field Φ in such a way that

$$g^*\Phi = (\Phi_0, \Phi_1, \dots, \Phi_{k-1}). \quad (9.39)$$

The fields Φ_i , $i = 0, \dots, k-1$.

In this way we get the following decomposition for a kinetic part of the field Φ and for a potential of this field:

$$\mathcal{L}_{\text{kin}}(\overset{\text{gauge}}{\nabla} \Phi) = \sum_{i=0}^{k-1} \mathcal{L}_{\text{kin}}^i(\overset{\text{gauge}}{\nabla} \Phi_i) \quad (9.40)$$

$$V(\Phi) = \sum_{i=0}^{k-1} V^i(\Phi_i) \quad (9.41)$$

$$\mathcal{L}_{\text{kin}}^i(\overset{\text{gauge}}{\nabla} \Phi_i) = \frac{1}{V_i} \int_{M_i} \sqrt{|\tilde{g}_i|} d^{\tilde{n}_i} x \left(l_{ab} g_i^{\tilde{b}_i \tilde{n}_i} L_{\tilde{b}_i}^{a\mu} \overset{\text{gauge}}{\nabla}_\mu \Phi_{i\tilde{n}_i}^b \right) \quad (9.42)$$

where

$$V_i = \int_{M_i} \sqrt{|\tilde{g}_i|} d^{\tilde{n}_i} x \quad (9.43)$$

$$\tilde{g}_i = \det(g_i \tilde{b}_i \tilde{a}_i). \quad (9.44)$$

$g_i \tilde{b}_i \tilde{a}_i$ is a nonsymmetric tensor on a manifold M_i .

$$\begin{aligned} V^i(\Phi_i) = & \frac{l_{ab}}{V_i} \int_{M_i} \sqrt{|\tilde{g}_i|} d^{\tilde{n}_i} x \left[2g^{[\tilde{m}_i \tilde{n}_i]} \left(C_{cd}^a \Phi_{i\tilde{m}_i}^c \Phi_{i\tilde{n}_i}^d - \mu_{\tilde{a}_i}^a f_{\tilde{m}_i \tilde{n}_i}^{\hat{l}_i} - \Phi_{i\tilde{e}_i}^a f_{\tilde{m}_i \tilde{n}_i}^{\tilde{e}_i} \right) \right. \\ & \times g^{[\tilde{a}_i \tilde{b}_i]} \left(C_{ef}^b \Phi_{i\tilde{a}_i}^e \Phi_{i\tilde{b}_i}^f - \mu_{\tilde{a}_i}^b f_{\tilde{a}_i \tilde{b}_i}^{\hat{j}_i} - \Phi_{i\tilde{a}_i}^b f_{\tilde{a}_i \tilde{b}_i}^{\tilde{d}_i} \right) \\ & \left. - g_i^{\tilde{a}_i \tilde{m}_i} g_i^{\tilde{b}_i \tilde{n}_i} L_{\tilde{a}_i \tilde{b}_i}^a \left(C_{cd}^b \Phi_{i\tilde{m}_i}^c \Phi_{i\tilde{n}_i}^d - \mu_{\tilde{a}_i}^b f_{\tilde{m}_i \tilde{n}_i}^{\hat{l}_i} - \Phi_{i\tilde{e}_i}^b f_{\tilde{m}_i \tilde{n}_i}^{\tilde{e}_i} \right) \right], \end{aligned} \quad (9.45)$$

$f_{\tilde{a}_i \tilde{b}_i}^{\hat{j}_i}$ are structure constants of the Lie algebra $\mathfrak{g}_i$.

The scheme of the symmetry breaking acts as follows from the group G_{i+1} to G_i (G'_i) (if the symmetry has been broken up to G_{i+1}). The potential $V^i(\Phi_i)$ has a minimum (global or local) for Φ_{crt}^k , $k = 0, 1$. The value of the remaining part of the sum (9.41) for fields Φ_j , $j < i$, is small for the scale of energy is much lower ($r_j > r_i$, $j < i$). Thus the minimum of $V^i(\Phi_i)$ is an approximate minimum of the remaining part of the sum (9.41). In this way we have a descending chain of truncations of the Higgs potential. This gives in principle a pattern of a symmetry breaking. However, this is only an approximate symmetry breaking. The real symmetry breaking is from G to G_0 (or to G'_0 in a false vacuum case). The important point here is the diffeomorphism g .

$$g^* \Phi^b = (\Phi_0^b, \Phi_1^b, \dots, \Phi_{k-1}^b) \quad (9.46)$$

$$\Phi_i^b = \Phi_{i\tilde{a}_i}^b \tilde{\theta}^{\tilde{a}_i}, \quad i = 0, \dots, k-1. \quad (9.47)$$

The shape of g is a true indicator of a reality of the symmetry breaking pattern. If

$$g = \text{Id} + \delta g \quad (9.48)$$

where δg is in some sense small and Id is an identity, the sums (9.40–41) are close to the formulae (6.17) and (6.18). The smallness of δg is a criterion of a practical application of the symmetry breaking pattern (9.34). It seems that there are a lot of possibilities for the condition (9.38). Moreover, a smallness of δg plus some natural conditions for groups G_i can narrow looking for grand unified models. Let us notice that the decomposition of M results in decomposition of cosmological terms

$$\tilde{P} = \sum_{i=0}^{k-1} \tilde{P}_i \quad (9.49)$$

where

$$\tilde{P}_i = \frac{1}{r_i^2 V_i} \int_{M_i} \sqrt{|\tilde{g}_i|} \hat{R}_i(\hat{\Gamma}_i) d^{n_i} x \quad (9.50)$$

where $\hat{\Gamma}_i$ is a nonsymmetric connection on M_i compatible with the nonsymmetric tensor $g_{i\tilde{a}_i\tilde{b}_i}$ and $\hat{R}_i(\hat{\Gamma}_i)$ its curvature scalar. The scalar field Ψ is now a function on a product of M_i ,

$$\Psi = \Psi(x, y_0, \dots, y_{k-1}), \quad y_i \in M_i, \quad i = 0, 1, \dots, k-1.$$

The truncation procedure can be proceeded in several ways. Finally let us notice that the energy scale of broken gauge bosons is fixed by a radius r_i at any stage of the symmetry breaking in our scheme. The possible groups in our symmetry breaking pattern can be taken from section 4 (see 4.62–4.66).

Let us consider Eq. (9.39) in more details. One gets

$$A_{i\tilde{a}_i}^{\tilde{a}}(y) \Phi_{\tilde{a}}^b(y) = \Phi_{\tilde{a}_i}^b(y_i), \quad y \in M, \quad y_i \in M_i \quad (9.51)$$

where

$$g^*(y) = \left(A_0 \left| A_1 \right| A_2 \left| \dots \right| A_{k-1} \right), \quad (9.52)$$

$$A_i = (A_{i\tilde{a}_i}^{\tilde{a}})_{\tilde{a}=1,2,\dots,n_1, \tilde{a}_i=1,2,\dots,\tilde{n}_i}, \quad i = 0, 1, 2, \dots, k \quad (9.53)$$

is a matrix of Higgs' fields transformation.

According to our assumptions one gets also:

$$\left(\frac{r_i}{r}\right)^2 g_{i\tilde{a}_i\tilde{b}_i}(y_i) = A_{\tilde{a}_i}^{\tilde{a}}(y) A_{\tilde{b}_i}^{\tilde{b}}(y) g_{\tilde{a}\tilde{b}}(y). \quad (9.54)$$

For g is an invertible map we have $\det g^*(y) \neq 0$.

We have also

$$n_1 = \sum_{i=0}^{k-1} \tilde{n}_i \quad (9.55)$$

and

$$\Phi_a^b(y) = \sum_{i=0}^{k-1} \tilde{A}_{i\tilde{a}}^{\tilde{a}_i}(y) \Phi_{\tilde{a}_i}^{\tilde{b}_i}(y_i) \quad (9.56)$$

or

$$g^{*-1}(y) = \left(\frac{\tilde{A}_0}{\tilde{A}_1} \right) \left(\frac{\tilde{A}_1}{\vdots} \right) \left(\frac{\vdots}{\tilde{A}_{k-1}} \right) \quad (9.57)$$

$$\tilde{A}_i = (\tilde{A}_{i\tilde{a}}^{\tilde{a}_i})_{\tilde{a}_i=1,2,\dots,\tilde{n}_i, \tilde{a}=1,2,\dots,n_1} \quad (9.58)$$

such that

$$g(y_0, \dots, y_{k-1}) = y \quad (9.59)$$

$$(y_0, y_1, \dots, y_{k-1}) = g^{-1}(y) \quad (9.59a)$$

For an inverse tensor $g^{\tilde{a}\tilde{b}}$ one easily gets

$$\left(\frac{r_i^2}{r^2}\right) g^{\tilde{a}\tilde{b}} = \sum_{i=0}^{k-1} \tilde{A}_{i\tilde{a}_i}^{\tilde{a}} g_i^{\tilde{a}_i\tilde{b}_i} A_{i\tilde{b}_i}^{\tilde{b}}. \quad (9.60)$$

We have

$$r^{2n_1} \det(g_{\tilde{a}\tilde{b}}) = \prod_{i=0}^{k-1} r_i^{2\tilde{n}_i} \det(g_{i\tilde{a}_i\tilde{b}_i}). \quad (9.61)$$

In this way we have for the measure

$$d\mu(y) = \prod_{i=0}^{k-1} d\mu_i(y_i) \quad (9.62)$$

where

$$d\mu(y) = \sqrt{\det g} r^{n_1} d^{n_1} y \quad (9.63)$$

$$d\mu_i(y_i) = \sqrt{\det g_i} r_i^{\bar{n}_i} d^{\bar{n}_i} y_i. \quad (9.64)$$

In the case of $\mathcal{L}_{\text{int}}(\Phi, \tilde{A})$ one gets

$$\mathcal{L}_{\text{int}}(\Phi, \tilde{A}) = \sum_{i=0}^{k-1} \mathcal{L}_{\text{int}}(\Phi_i, \tilde{A}) \quad (9.65)$$

where

$$\mathcal{L}_{\text{int}}(\Phi_i, \tilde{A}) = h_{ab} \mu_i^a \tilde{H}^i \underline{g}_i^{[\tilde{a}_i \tilde{b}_i]} \left(C_{cd}^b \Phi_{i\tilde{a}_i}^c \Phi_{i\tilde{b}_i}^d - \mu_i^b f_{\tilde{a}_i \tilde{b}_i}^{\hat{i}} - \Phi_{\tilde{d}_i}^b f_{\tilde{a}_i \tilde{b}_i}^{\tilde{d}_i} \right) \quad (9.66)$$

where

$$\underline{g}_i^{[\tilde{a}_i \tilde{b}_i]} = \frac{1}{V_i} \int_{M_i} \sqrt{|\tilde{g}_i|} d^{\bar{n}_i} x g_i^{[\tilde{a}_i \tilde{b}_i]}, \quad i = 0, 1, 2, \dots, k-1. \quad (9.67)$$

Moreover, to be in line in the full theory we should consider a chain of groups H_i , $i = 0, 1, 2, \dots, k-1$, in such a way that

$$H_0 \subset H_1 \subset H_2 \subset \dots \subset H_{k-1} = H. \quad (9.68)$$

For every group H_i we have the following assumptions

$$G_i \subset H_i \quad (9.69)$$

and G_{i+1} is a centralizer of G_i in H_i . Thus we should have

$$G_i \otimes G_{i+1} \subset H_i, \quad i = 0, 1, 2, \dots, k-1. \quad (9.70)$$

We know from elementary particles physics theory that

$$\begin{aligned} G_0 &= U_{\text{el}} \otimes SU(3)_{\text{color}}, \\ G_1 &= SU(2)_{\text{L}} \otimes U(1)_{\text{Y}} \otimes SU(3)_{\text{color}} \end{aligned}$$

and that G_2 is a group which plays the role of H in the case of a symmetry breaking from $SU(2)_{\text{L}} \otimes U_{\text{Y}}(1)$ to $U_{\text{el}}(1)$. However, in this case because of a factor $U(1)$, $M = S^2$. Thus $M_0 = S^2$ and $G_2 \subset H_0$.

It seems that in a reality we have to do with two more stages of a symmetry breaking. Thus $k = 3$. We have

$$M \simeq S^2 \times M_1 \times M_2 \quad (9.71)$$

$$M = G/(U(1) \otimes SU(3)) \quad (9.72)$$

$$M_1 = G_1/(SU(2) \times U(1) \times SU(3))$$

$$U(1) \otimes SU(3) \subset SU(2) \otimes U(1) \otimes SU(3) \subset G_2 \otimes SU(3) \subset G_3 = G \quad (9.73)$$

and

$$G_1 \subset H_1 \subset H \quad (9.74)$$

$$U(1) \otimes SU(3) \otimes G \subset H \quad (9.74a)$$

$$(U(1) \otimes SU(2) \otimes SU(3)) \otimes G_2 \subset H_1 \quad (9.74b)$$

and

$$G_2 \otimes G \subset H_2 = H \quad (9.74c)$$

$$M_2 = G/G_1. \quad (9.75)$$

We can take for G $SU(5)$, $SO(10)$, $E6$ or $SU(6)$. Thus there are a lot of choices for G_2 , H_1 and H .

We can suppose for a trial that

$$G_2 \otimes SU(3) \subset H_0. \quad (9.75a)$$

We have also some additional constraints

$$\text{rank}(G) \geq 4. \quad (9.75b)$$

Thus

$$\text{rank}(H_0) \geq 4. \quad (9.75c)$$

We can try with $F4 = H_0$.

In the case of H

$$\text{rank}(H) \geq \text{rank}(G) + 3 \geq 7. \quad (9.75d)$$

Thus we can try with $E7$, $E8$

$$\text{rank}(H_1) \geq \text{rank}(G_2) + 4$$

$$\text{rank}(H) \geq \text{rank}(G_2) + \text{rank}(G) \geq \text{rank}(G_2) + 4 \geq \text{rank}(G) + 4 \geq 8. \quad (9.75e)$$

In this way we have

$$\text{rank}(H) \geq 8. \quad (9.75f)$$

Thus we can try with

$$H = E8. \quad (9.75g)$$

But in this case

$$\text{rank}(G_2) = \text{rank}(G) = 4.$$

This seems to be nonrealistic. For instance, if $G = SO(10)$, $E6$,

$$\begin{aligned}\text{rank}(SO(10)) &= 5 \\ \text{rank } E6 &= 6.\end{aligned}$$

In this case we get

$$\begin{aligned}\text{rank}(H) &= 9 \\ \text{rank}(H) &= 10\end{aligned}$$

and H could be $SU(10)$, $SO(18)$, $SO(20)$.

In this approach we try to consider additional dimensions connecting to the manifold M more seriously, i.e. as physical dimensions, additional space-like dimensions. We remind to the reader that gauge-dimensions connecting to the group H have different meaning. They are dimensions connected to local gauge symmetries (or global) and they cannot be directly observed. Simply saying we cannot travel along them. In the case of a manifold M this possibility still exists. However, the manifold M is diffeomorphically equivalent to the product of some manifolds M_i , $i = 0, 1, 2, \dots, k-1$, with some characteristic sizes r_i .

The radii r_i represent energy scales of symmetry breaking. The lowest energy scale is a scale of weak interactions (Weinberg-Glashow-Salam model) $r_0 \simeq 10^{-16}$ cm. In this case this is a radius of a sphere S^2 . The possibility of this “travel” will be considered in section 16. In this case a metric on a manifold M can be dependent on a point $x \in E$ (parametrically).

It is interesting to ask on a stability of a symmetry breaking pattern with respect to quantum fluctuations. This difficult problem strongly depends on the details of the model. Especially on the Higgs sector of the practical model. In order to preserve this stability on every stage of the symmetry breaking we should consider remaining Higgs’ fields (after symmetry breaking) with zero mass. According to S. Weinberg, they can stabilize the symmetry breaking in the range of energy

$$\frac{1}{r_i} \left(\frac{\hbar}{c} \right) < E < \frac{1}{r_{i+1}} \left(\frac{\hbar}{c} \right), \quad i = 0, 1, 2, \dots, k-1, \quad (9.76)$$

i.e. for a symmetry breaking from G_{i+1} to G_i .

In this place we can introduce deconfinement parameters $\Lambda(G_i) = \frac{\alpha_s}{r_i} \left(\frac{\hbar}{c} \right)$ which introduce a scale of energy for a symmetry breaking.

It seems that in order to create a realistic grand unified model based on nonsymmetric Kaluza–Klein (Jordan–Thiry) theory it is necessary to nivel cosmological terms. This could be achieved in some models due to choosing constants ξ and ζ and μ . After this we can control the value of those terms, which are considered as a selfinteraction potential of a scalar field Ψ . The scalar field Ψ can play in this context a role of a quintessence.

Let us notice that using (4.51) and (9.56) one gets

$$\sum_{i=0}^{k-1} \tilde{A}_{i\tilde{b}}^{\tilde{a}_i} \Phi_{\tilde{a}_i}^c f_{i\tilde{a}}^{\tilde{b}} = C_{ab}^c \mu_{\tilde{a}}^a \sum_{i=0}^{k-1} \tilde{A}_{i\tilde{a}}^{\tilde{a}_i} \Phi_{\tilde{a}_i}^b. \quad (9.77)$$

In this way we get constraints for Higgs' fields $\Phi_0, \Phi_1, \dots, \Phi_{k-1}$,

$$\Phi_i = (\Phi_{\tilde{a}_i}^b), \quad i = 0, 1, \dots, k-1.$$

Solving these constraints we obtain some of Higgs' fields as functions of independent components. This could result in some cross terms in the potential (9.41) between Φ 's with different i . For example a term

$$V(\Phi'_i, \Phi'_j),$$

where Φ' means independent fields. This can cause some problems in a stability of symmetry breaking pattern against radiative corrections. This can be easily seen from Eq. (4.51) solved by independent Φ' ,

$$\Phi = B\Phi' \quad (9.78)$$

$$\Phi_b^c = B_{b\tilde{c}}^{\tilde{c}\tilde{b}} \Phi'_{\tilde{b}}^{\tilde{c}} \quad (9.79)$$

where B is a linear operator transforming independent Φ' into Φ .

We can suppose for a trial a condition similar to (4.51) for every $i = 0, \dots, k-1$,

$$\Phi_{\tilde{b}_i}^{c_i} f_{i\tilde{a}_i}^{\tilde{b}_i} = \mu_{\tilde{a}_i}^{a_i} \Phi_{\tilde{a}_i}^{b_i} C_{a_i b_i}^{c_i} \quad (9.80)$$

where $C_{a_i b_i}^{c_i}$ are structure constants for the Lie algebra $\mathfrak{h}_i$ of the group H_i . $f_{i\tilde{a}_i}^{\tilde{b}_i}$ are structure constants of the Lie algebra $\mathfrak{g}_{i+1}$, $\tilde{a}_i$ are indices belonging to Lie algebra $\mathfrak{g}_i$ and $\tilde{b}_i$ to the complement $\mathfrak{m}_i$.

In this way

$$\Phi_{\tilde{b}_i}^{c_i} = \Phi_{\tilde{b}_i}^{c_i} \delta_{c_i}^{c_i}. \quad (9.81)$$

In this case we should have a consistency between (9.80) and (9.77) which impose constraints on C, f, μ and C^i, f^i, μ^i where C^i, f^i, μ^i refer to H_i, G_{i+1} . Solving (9.80) via introducing independent fields Φ'_i one gets

$$\Phi_{i\tilde{b}_i}^{c_i} = B_{i\tilde{c}_i\tilde{b}_i}^{c_i\tilde{b}_i} \Phi'_{i\tilde{b}_i}^{\tilde{c}_i}. \quad (9.82)$$

Combining (9.79), (9.81–82) one gets

$$B_{b\tilde{c}}^{\tilde{c}\tilde{b}} \Phi'_{\tilde{b}}^{\tilde{c}} = \sum_{i=0}^{k-1} \tilde{A}_{i\tilde{b}}^{\tilde{a}_i} \delta_{c_i}^{\tilde{c}_i} B_{i\tilde{c}_i\tilde{b}_i}^{c_i\tilde{b}_i} \Phi'_{i\tilde{b}_i}^{\tilde{c}_i}. \quad (9.83)$$

Eq. (9.83) gives a relation between independent Higgs' fields Φ' and Φ'_i . Simultaneously it is a consistency condition between Eq. (4.51) and Eq. (9.80). However, the condition (9.80) seems to be too strong and probably it is necessary to solve a weaker condition

(9.77) which goes to the mentioned terms $V(\Phi'_i, \Phi'_j)$. The conditions (9.80) plus a consistency (9.83) avoid those terms in the Higgs potential. This problem demands more investigations.

It seems that the condition (9.38) could be too strong. In order to find a more general condition we consider a simple example of (9.34). Let $G_0 = \{e\}$ and $K = 2$. In this case we have

$$\{e\} \subset G_1 \subset G_2 = G \quad (9.84)$$

$$M_0 = G_1, \quad M_1 = G/G_1 \quad (9.85)$$

$$g : G_1 \times G/G_1 \rightarrow G. \quad (9.86)$$

In this way $G_1 \times G/G_1$ is diffeomorphically equivalent to G .

Moreover, we can consider a fibre bundle with base space G/G_1 and a structural group G_1 with a bundle manifold G . This construction is known in the theory of induced group representation (see [51] and section 4). The projection $\varphi : G \rightarrow G/G_1$ is defined by $\varphi(g) = \{gG_1\}$. The natural extension of (9.86) is to consider a fibre bundle $(G, G/G_1, G_1, \varphi)$. In this way we have in a place of (9.86) a local condition

$$g_U : G_1 \times U \xrightarrow{\text{in}} G \quad (9.87)$$

where $U \subset G/G_1$ is an open set. Thus in a place of (9.38) we consider a local diffeomorphism

$$g_U : M_0 \times M_1 \times \dots \times M_{k-1} \xrightarrow{\text{in}} G/G_0. \quad (9.88)$$

where

$$U = U_0 \times U_1 \times \dots \times U_{k-1},$$

$U_i \subset M_i$, $i = 0, 1, 2, \dots, k-1$, are open sets. Moreover we should define projectors φ_i , $i = 0, 1, 2, \dots, k-1$,

$$\varphi_i : G/G_0 \rightarrow G_{i+1}/G_i, \quad (9.89)$$

i.e.

$$\varphi_i(\{gG_0\}) = \{g_{i+1}G_i\}, \quad (9.90)$$

$$g \in G, \quad g_{i+1} \in G_{i+1}, \quad G_0 \subset G_i \subset G_{i+1} \subset G$$

in a unique way. This could give us a fibration of G/G_0 in $\prod_{i=0}^{k-1} (G_{i+1}/G_i)$.

For $g \in G_{i+1}$ we simply define

$$\varphi_i(\{gG_0\}) = \{gG_i\}. \quad (9.91)$$

If $g \in G$, $g \notin G_{i+1}$, we define

$$\varphi_i(\{gG_0\}) = \{G_i\}. \quad (9.92)$$

Thus in general

$$\varphi_i(\{gG_0\}) = \{p(g)G_i\} \quad (9.93)$$

where

$$p(g) = \begin{cases} g, & g \in G_{i+1} \\ e, & g \notin G_{i+1}. \end{cases} \quad (9.94)$$

Thus in a place of (9.38) we have to do with a structure

$$\left\{ G/G_0, \prod_{i=0}^{k-1} (G_{i+1}/G_i), \varphi_0, \varphi_1, \dots, \varphi_{k-1} \right\} \quad (9.95)$$

such that

$$g_U \circ \varphi|_U = \text{id} \quad (9.96)$$

where

$$\varphi|_U = \prod_{i=0}^{k-1} \varphi_i|_{U_i}. \quad (9.97)$$

This generalizes (9.38) to the local conditions (9.88). Now we can repeat all the considerations concerning a decomposition of Higgs' fields using local diffeomorphisms g_U (g_U^*) in the place of g (g^*). Let us also notice that in the chain of groups it would be interesting to consider as G_2

$$G_2 = SU(2)_L \otimes SU(2)_R \otimes SU(4)$$

suggested by Salam and Pati, where $SU(4)$ unifies $SU(3)_{\text{color}} \otimes U(1)_Y$. This will be helpful in our future consideration concerning extension to supersymmetric groups, i.e. $U(2|2)$ which unifies $SU(2)_L \otimes SU(2)_R$ to the super Lie group $U(2|2)$ considered by Mohapatra. Such models on the phenomenological level incorporate fermions with a possible extension to the supersymmetric $SO(10)$ model. They give a natural framework for lepton flavour mixing going to the neutrino oscillations incorporating see-saw mechanism for mass generations of neutrinos. In such approaches the see-saw mechanism is coming from the grand unified models. Our approach after incorporating manifolds with anticommuting parameters, super Lie groups, super Lie algebras and in general supermanifolds (superfibrebundles) can be able to obtain this. However, it is necessary to develop a formalism (in the language of supermanifolds, superfibrebundles, super Lie groups, super Lie algebras) for nonsymmetric connections, nonsymmetric Kaluza–Klein (Jordan–Thiry) theory. In particular we should construct an analogue of Einstein–Kaufmann connection for supermanifold, a nonsymmetric Kaluza–Klein (Jordan–Thiry) theory for superfibrebundle with super Lie group. In this way we should define first of all a nonsymmetric tensor on a super Lie group and afterwards a nonsymmetric metrization of a superfibrebundle.

Let us notice that on every stage of symmetry breaking, i.e. from G_{i+1} to G_i , we have to do with group G'_i (similar to the group G'_0). Thus we can have to do with a true and a false vacuum cases which may complicate a pattern of a symmetry breaking.

10. Properties of the Scalar Field Ψ . Cosmological Drifting of the Mass Scale

In section 9 we get the equation for scalar field Ψ (see (9.6)). It has the following shape:

$$\hat{\square}\Psi + I(\Psi) = 0, \quad (10.1)$$

where $\hat{\square}$ is a wave operator (differential linear operator of the second order) and $I(\Psi)$ describes interaction of Ψ with Higgs' field, Yang–Mills' field and cosmological terms. In the terms of the effective gravitational constant the field Ψ is defined:

$$\Psi = -\frac{1}{(n+2)} \cdot \ln \left(\frac{G_{\text{eff}}}{G_N} \right). \quad (10.2)$$

Simultaneously the field Ψ enters the equation (8.21) and (8.22). It means that there exists an effective scale of masses for broken gauge bosons. It is

$$m_{\text{eff}} = e^{-\Psi} m_{\tilde{A}} \simeq e^{-\Psi} \frac{\alpha_S}{r} \frac{\hbar}{c}, \quad (10.3)$$

The field Ψ enters also the effective cosmological constant for “false vacuum” case. We get:

$$\lambda_{c1} = e^{(n-2)\Psi} \frac{4}{\hbar c r^2} \left(\frac{\ell_{pl}^2}{r^2} \right) V(\Phi_{\text{crt}}^1). \quad (10.4)$$

Using effective scale of masses one gets:

$$\lambda_{c1} = 4e^{(n+2)\Psi} \frac{m_{\text{eff}}^4}{\alpha_S^4} \left(\frac{\hbar^3}{c^5} \right) \ell_{pl}^2 V(\Phi_{\text{crt}}^1). \quad (10.5)$$

Let us suppose that we have a situation corresponding to one of the minima of the potential $V(\Phi)$. It means that:

$$\begin{aligned} \mathcal{L}_{\text{int}}(\tilde{A}, \Phi) &= 0, \\ V(\Phi) &= V(\Phi_{\text{crt}}^k), \\ \mathcal{L}_{\text{kin}}(\overset{\text{gauge}}{\nabla} \Phi) &= N^{\mu\nu} M_{ij}^2(\Phi_{\text{crt}}^k) \tilde{A}_\mu^i \tilde{A}_\nu^j, \\ \mathcal{L}_{\text{YM}} &= 0. \end{aligned} \quad (10.6)$$

Let us suppose that the cosmological terms are negligible. In this case we have for $I(\Psi)$:

$$I(\Psi) = \frac{4e^{-2\Psi}}{r^2} N^{\mu\nu} M_{ij}^2(\Phi_{\text{crt}}^k) \tilde{A}_\mu^i \tilde{A}_\nu^j + \frac{(n-2)}{r^4} e^{(n-2)\Psi} V(\Phi_{\text{crt}}^k). \quad (10.7)$$

In the vacuum state the fields $\tilde{A}_\mu^i$ and Ψ fluctuate only around this state. Thus

$$\begin{aligned} \tilde{A}_\mu^i &= \tilde{A}_{0\mu}^i + \delta\tilde{A}_\mu^i, \\ \Psi &= \Psi_0 + \delta\Psi. \end{aligned} \quad (10.8)$$

It means that:

$$\begin{aligned} I(\Psi) &\simeq \left(\frac{4e^{-2\Psi_0}}{r^2} \eta^{\mu\nu} M_{ij}^2(\Phi_{\text{crt}}^k) \tilde{A}_{0\mu}^i \tilde{A}_{0\nu}^j + \frac{(n-2)}{r^4} e^{(n-2)\Psi_0} V(\Phi_{\text{crt}}^k) \right) + \\ &+ \left(\frac{-8e^{-2\Psi_0}}{r^2} \eta^{\mu\nu} M_{ij}^2(\Phi_{\text{crt}}^k) \tilde{A}_{0\mu}^i \tilde{A}_{0\nu}^j + \frac{(n-2)^2}{r^4} e^{(n-2)\Psi_0} V(\Phi_{\text{crt}}^k) \right) \delta\Psi = \\ &= d_1 + d_2 \delta\Psi, \end{aligned} \quad (10.9)$$

d_1, d_2 are constants and we put $N^{\mu\nu} = \bar{M} \eta^{\mu\nu}$ (in zero order of approximation). And from (10.1) we get:

$$\square(\delta\Psi) + \left(\frac{d_1}{\bar{M}} \right) + \left(\frac{d_2}{\bar{M}} \right) \delta\Psi = 0. \quad (10.10)$$

It means that (10.1) becomes the Klein–Gordon equation for the fluctuation of the scalar field Ψ and $\delta\Psi$ is a massive field. Thus it has Yukawa-type behaviour.

$$\delta\Psi \sim \frac{1}{r} e^{-\alpha r} + \text{const}, \quad \alpha > 0. \quad (10.11)$$

For the field Ψ we get:

$$\Psi \sim \text{const}' + \frac{1}{r} e^{-\alpha r}, \quad \alpha > 0. \quad (10.12)$$

Due to this we preserve the weak equivalence principle for the gravitational field. The scalar forces connected to the field Ψ are of the short range. This is possible if $d_2 > 0$. The first term in d_2 is greater than zero and we have no troubles in the true vacuum case ($V(\Phi_{\text{crt}}^0) = 0$). In the case of the false vacuum the situation may be different ($d_2 \leq 0$). Thus the world built over the false vacuum state can violate the weak equivalence principle. Summing up we get that in the case of true vacuum the field Ψ is massive with short range. It is in accordance with the weak equivalence principle, the universal fall of all bodies in gravitational field. This property is connected, from cosmological point of view of course, to space dependence of Ψ (or G_{eff}). It is interesting to ask what will happen if $G_{\text{eff}} = G_{\text{eff}}(t)$. It means that $\Psi = \Psi(t)$ depends on the time t only. Now it is a cosmological time. In this case $\Psi(t)$ plays the role of the cosmological

factor. Let us estimate the dependence on time for the effective scale of masses, m_{eff} , using some estimations for G_{eff} . We have (see [56], [57]):

$$\eta = \frac{\Delta G_{\text{eff}}}{G_{\text{eff}} \Delta t} = \frac{10^{-11} - 10^{-10}}{\text{yr}}, \quad (10.13)$$

$$\eta = \frac{d}{dt} \ln G_{\text{eff}}(t) = -(n+2) \left(\frac{d\Psi}{dt} \right). \quad (10.14)$$

Thus we get

$$\Delta\Psi = -\frac{\eta}{(n+2)} \Delta t. \quad (10.15)$$

For the effective scale of mass one gets:

$$\omega = \frac{\Delta m_{\text{eff}}}{m_{\text{eff}} \Delta t} = \frac{d}{dt} \ln m_{\text{eff}}(t) = -\frac{d}{dt} \Psi(t) = -\frac{\Delta\Psi}{\Delta t}. \quad (10.16)$$

Thus we get

$$\omega = \frac{\eta}{(n+2)} \quad (10.17)$$

and

$$\omega = \frac{10^{-12} - 10^{-11}}{\text{yr}}, \quad (10.18)$$

if we suppose that $n \geq 24$ ($n = \dim G$ and for $G = \text{SU}(5)$ we can have $n = 24$). Thus we are able to connect changing of the effective gravitational constant with very small drifting of the scale of masses for the broken gauge bosons. This drifting is really very small and it does not contradict experimental data. It is worth to notice that the sign of changing of G_{eff} and m_{eff} is the same. If G_{eff} increases then m_{eff} increases too and vice-versa. This behaviour is in agreement with Mach's principle.

Let us remind to the reader the Dirac large number hypothesis. He gets that

$$\frac{e^2}{G_N m_e m_p} \simeq 10^{40}, \quad (10.19)$$

where m_e , m_p mean electron and proton mass, respectively, and

$$\frac{t}{e^2/m_e c^3} \simeq 10^{40}, \quad (10.20)$$

where t is an age of the Universe and $\frac{e^2}{m_e c^3}$ is a time needed for light to pass a distance of a classical radius of an electron.

If those large numbers are equal forever one gets

$$\frac{e^2}{G_N m_e m_p} \simeq \frac{t}{e^2/m_e c^3} \quad (10.21)$$

and we get an interesting cosmological dependence

$$\frac{1}{t} = \left(\frac{m_p c^3}{e^4} \right) G m_e^2. \quad (10.22)$$

The mass of a proton is established by Λ_{QCD} and we do not expect the cosmological drifting of Λ_{QCD} . Thus $m_p = \text{const.}$ However, in the case of m_e is different

$$m_e \sim m_{\text{EW eff}} \sim \frac{1}{r} e^{-\Psi} \sim m_W e^{-\Psi} \quad (10.23)$$

and

$$G = G_{\text{eff}} = G_N e^{-(n+2)\Psi}. \quad (10.24)$$

Thus we get

$$\frac{1}{t} = C e^{-(n+4)\Psi} \quad (10.25)$$

where C is a constant.

And finally

$$\Psi(t) = \frac{1}{n+4} \ln(Ct), \quad (10.26)$$

which gives an evolution law for a scalar field Ψ . This could be incorporated as an ansatz for some cosmological models, where

$$C \simeq \frac{10^{20}}{t_{\text{pl}}} \quad (10.27)$$

and t_{pl} is a Planck time.

11. Fermion Number, $\mathbb{R}_+$ and $U(1)_F$ Invariance

In N.G.T. and in Einstein Unified Field Theory there exists a gauge invariance for the 2-form of curvature

$$\overline{\Omega}^\alpha{}_\beta(\overline{W}) = d\overline{W}^\alpha{}_\beta + \overline{W}^\alpha{}_\beta \wedge \overline{W}^\gamma{}_\beta \quad (11.1)$$

such that

$$\overline{\Omega}^\alpha{}_\beta(\overline{W}) = \overline{\Omega}^\alpha{}_\beta(\overline{W}'), \quad (11.2)$$

where

$$\overline{W}^\alpha{}_\beta \rightarrow \overline{W}'^\alpha{}_\beta = \overline{W}^\alpha{}_\beta - \frac{2}{3}\delta^\alpha{}_\beta d\phi \quad (11.3)$$

and where ϕ is a function on E . This results as a gauge transformation for the 1-form $\overline{W}$

$$\overline{W} \rightarrow \overline{W}' = \overline{W} + d\phi. \quad (11.4)$$

It is easy to see that the 2-form of curvature for the connection $W^{\tilde{A}}{}_{\tilde{B}}$ is also invariant under (11.4)

$$\Omega^{\tilde{A}}{}_{\tilde{B}}(W) = \Omega^{\tilde{A}}{}_{\tilde{B}}(W'), \quad (11.5)$$

where

$$W'^{\tilde{A}}{}_{\tilde{B}} = W^{\tilde{A}}{}_{\tilde{B}} - \frac{4}{3(m+2)}\delta^{\tilde{A}}{}_{\tilde{B}}d\phi. \quad (11.6)$$

In N.G.T. (see [34], [47], [65], [66]) the vector field $\overline{W}_\mu$ is coupled to the fermion current (fermion number plays in this theory the role of the second gravitational charge). Thus (11.4) results in the conservation law for the fermion current. Fermion current is simultaneously the source of the skew-symmetric part of the metric (see for details [34], [47], [65]). The fermion current for a Dirac particle has the same shape as an electric current

$$S^\mu \sim \overline{\Psi}\gamma^\mu\Psi. \quad (11.7)$$

We have only a different coupling constant between S^μ and $\overline{W}_\mu$.

One can ask what is the origin of the transformation (11.3). It is very well known that the linear connection (coefficients of the connection) $\overline{W}^\alpha{}_\beta$ obeys the following law of transformation: (the linear connection is not a geometrical quantity of σ -type in a Shouten sense as for example tensor, vector, spinor, 2-form of curvature, etc., see Ref. [59] for more details).

$$A^\mu{}_\nu \overline{W}'^\nu{}_\beta = \overline{W}^\mu{}_\nu A^\nu{}_\beta + dA^\mu{}_\beta, \quad (11.8)$$

if we change basic forms $\bar{\theta}^\alpha$ into $\bar{\theta}'^\alpha$ such that

$$\bar{\theta}^\alpha = A^\alpha_\beta \bar{\theta}'^\beta, \quad (11.9)$$

where A^α_β is a matrix valued function on the manifold E (space-time) with values in $\text{GL}(4, \mathbb{R})$, $\det(A^\alpha_\beta) \neq 0$. If we restrict $\text{GL}(4, \mathbb{R})$ to the subgroup $\text{GL}_\downarrow(4, \mathbb{R}) = \{(A^\alpha_\beta), \det(A^\alpha_\beta) > 0\}$ we may consider the following transformation:

$$A^\alpha_\beta = \delta^\alpha_\beta e^{-\frac{2}{3}\phi} \quad (11.10)$$

and we get from (11.8) Eq. (11.3). It is easy to see that (11.10) is a dilatation. Let us write every matrix A from $\text{GL}_\downarrow(4, \mathbb{R})$ in the form:

$$A = e^B. \quad (11.11)$$

This is possible, because $\det A > 0$ and let us decompose B into two parts

$$B^\alpha_\beta = (B^\alpha_\beta - b\delta^\alpha_\beta) + b\delta^\alpha_\beta = B'^\alpha_\beta - b\delta^\alpha_\beta, \quad (11.12)$$

where $b = 1/4B^\alpha_\alpha$ and B'^α_β is a traceless matrix. According to (11.12) one gets

$$A = e^b e^{B'}. \quad (11.13)$$

But

$$\det A = e^{\text{Tr } B} = e^{4b}. \quad (11.14)$$

Comparing (11.13), (11.14) and (11.10) one gets:

$$\phi = -\frac{3}{8} \ln(\det A), \quad (11.15)$$

where $A \in \text{GL}_\downarrow(4, \mathbb{R})$. In this way we have the decomposition

$$\text{GL}_\downarrow(4, \mathbb{R}) = \mathbb{R}_+ \otimes \text{SL}(4, \mathbb{R}), \quad (11.16)$$

where $\mathbb{R}_+ = \{e^\rho, \rho \in \mathbb{R}\}$ and $\text{SL}(4, \mathbb{R}) = \{A \in \text{GL}(4, \mathbb{R}), \det A = 1\}$. The gauge transformation (11.3), (11.4) and (11.6) are the gauge transformations corresponding to $\mathbb{R}_+$ in the decomposition (11.16). Transformation (11.4) induces the following dilatation transformations for spinor fields Ψ and $\bar{\Psi}$ (see Ref. [92])

$$\Psi \rightarrow \Psi' = \Psi e^{-i\beta\phi}, \quad (11.17)$$

$$\bar{\Psi} \rightarrow \bar{\Psi}' = \bar{\Psi} e^{i\beta\phi}, \quad (11.18)$$

where β is a constant connected to the universal coupling constant a from Moffat's theory. In this way we have compactification of the group $\mathbb{R}_+$ from the decomposition (11.16) (see Ref. [92] for details) and spinors transform according to the following group:

$$\mathbf{U}(1)_F \otimes \text{SL}(2, \mathbb{C}). \quad (11.19)$$

In the case of electromagnetic fields we have

$$\mathbf{U}(1)_{el} \otimes \mathrm{SL}(2, \mathbb{C}) \quad (11.20)$$

and the transformation law for Ψ and $\bar{\Psi}$

$$\Psi \rightarrow \Psi' = \Psi e^{-i \frac{e}{\hbar c} \chi}, \quad (11.21)$$

$$\bar{\Psi} \rightarrow \bar{\Psi}' = \bar{\Psi} e^{i \frac{e}{\hbar c} \chi}, \quad (11.22)$$

$$A \rightarrow A' = A + d\chi. \quad (11.23)$$

The difference between (11.19) and (11.20) is significant, because $\mathbf{U}(1)_F$ is a compactification of $\mathbb{R}_+$ and $\mathbf{U}(1)_{el}$ is compact from the very beginning. But there is a more fundamental difference, $\mathbb{R}_+$ (dilatations) belong to space-time symmetries and $\mathbf{U}(1)_{el}$ to the internal symmetry of the electromagnetic field. It is easy to see that we can get (11.19) from the following scheme

$$\mathrm{GL}(4, \mathbb{R}) \rightarrow \mathrm{GL}_{\downarrow}(4, \mathbb{R}) \rightarrow \mathbb{R}_+ \otimes \mathrm{SL}(4, \mathbb{R}) \rightarrow \mathbb{R}_+ \otimes \mathrm{SO}(1, 3) \rightarrow \mathbf{U}(1)_F \otimes \mathrm{SL}(2, \mathbb{C}) \quad (11.24)$$

and $\mathbf{U}(1)_F$ is a compactification of $\mathbb{R}_+$ in the following sense:

$$\mathbf{U}(1)_F = \left\{ e^{-i\psi}, \psi = \frac{3}{8} \ln(\det(A^\mu{}_\nu)), \det(A^\mu{}_\nu) > 0 \right\}. \quad (11.25)$$

Fermion number is independently connected to the internal symmetries of elementary particles. We know that there are no forces with long range connected to fermion number (weak equivalence principle). In N.G.T. there are also no forces with long range connected to this charge. The theory satisfies the weak equivalence principle — universal fall of all bodies. In the Newtonian approximation of N.G.T. there is no Coulomb force connecting to the $\bar{W}_\mu$ potential or to the skew-symmetric part of the metric $g_{[\mu\nu]}$. On the other hand fermion number is conserved in a similar manner to that of electric charge and it is usually connected to the $\mathbf{U}(1)_F$ internal invariance.

In the Grand Unified Theory based on the $\mathrm{SO}(10)$ group (see [93]), fermion number is one of the generators of the Lie algebra of $\mathrm{SO}(10)$, Y_F , and $F = B - L$, where B is the baryon number and L the lepton number. Thus we have the local symmetry $\mathbf{U}(1)_F$. After spontaneous symmetry breaking the intermediate gauge boson corresponding to Y_F in the adjoint representation obtains a rest mass and we have no forces of long range connecting to F .

This symmetry breaking is the spontaneous symmetry breaking and the lagrangian of the theory is invariant under local transformations of $\mathrm{SO}(10)$ (thus under $\mathbf{U}(1)_F$ too). Symmetry is breaking only due to choosing a concrete representative of the vacuum, which is here degenerated.

In the Grand Unified Theory based on $\mathrm{SU}(5)$ we have no $\mathbf{U}(1)_F$ local invariance, we have only global $\mathbf{U}(1)_F$ invariance.

In the $\mathrm{SO}(10)$ G.U.T. we get after symmetry breaking $\mathbf{U}(1)_F$ global invariance and the fermion number $F = B - L$ is conserved. Space-time symmetries and internal

symmetries are usually disconnected. Only some geometrical approaches similar to dimensional reduction or Kaluza–Klein theory or maybe supersymmetry are able to connect these symmetries treating them as “space-time” symmetries on many-dimensional manifolds.

In our theory we have this situation. We have a many-dimensional manifold and after dimensional reduction we get gravity coupled to the Yang–Mills’ field and to the Higgs’ field with spontaneous symmetry breaking and the Higgs’ mechanism. Simultaneously gravitation in our theory is described by N.G.T. Thus we should try to connect $\mathbb{R}_+$ symmetry from N.G.T. with $\mathbf{U}(1)_F$ internal symmetry. There are two possibilities for such relations.

1) Let us suppose that $\mathbf{U}(1)_F$ is a subgroup of the group G

$$\mathbf{U}(1)_F \subset G \quad (11.26)$$

in such a way that $Y_F \in \mathfrak{m}$. This means that after symmetry breaking the intermediate boson corresponding to Y_F (for “true” vacuum case) obtains the rest mass and we will have no long range forces connecting to F . This is similar to the $\mathrm{SO}(10)$ G.U.T.

Now let us suppose that there exists a homomorphism between $\mathrm{GL}_\downarrow(4, \mathbb{R})$ and G

$$\varepsilon : \mathrm{GL}_\downarrow(4, \mathbb{R}) \rightarrow G, \quad (11.27)$$

such that

$$\varepsilon(A) = e^{-(i\frac{3}{8} \ln(\det A) Y_F)}, \quad (11.28)$$

where $A = (A^\mu{}_\nu)$, $\det A > 0$.

(11.28) is really the homomorphism between $\mathrm{GL}_\downarrow(4, \mathbb{R})$ and $\mathbf{U}(1)_F$. It projects $\mathrm{GL}_\downarrow(4, \mathbb{R})$ to a one-parameter subgroup of G . This has the result that dilatations on the space-time from $\mathbb{R}_+$ induce internal rotations of $\mathbf{U}(1)_F$.

2) Let us suppose that the group H has the following structure

$$\mathbf{U}(1)_F \not\subset G, \quad G \otimes G_0 \subset H, \quad \mathbf{U}(1)_F \subset H. \quad (11.29)$$

Let there exists a homomorphism between $\mathrm{GL}_\downarrow(4, \mathbb{R})$ and H

$$\eta : \mathrm{GL}_\downarrow(4, \mathbb{R}) \rightarrow H, \quad (11.30)$$

such that

$$\eta(A) = e^{-(i\frac{3}{8} \ln(\det A) Y_F)}, \quad (11.31)$$

where $A = (A^\mu{}_\nu)$, $\det A > 0$. Now dilatations from the space-time also induce internal $\mathbf{U}(1)_F$ rotations. However the situation is different. Now Y_F does not belong to the Lie algebra of the group G and due to the Wang condition there is no gauge boson connecting to $\mathbf{U}(1)_F$ (to fermion number). There is no gauge field defined on space-time connecting to Y_F . The $\mathbf{U}(1)_F$ symmetry acts only on the Higgs’ multiplet and it is really global. Thus we have a situation similar to $\mathrm{SU}(5)$ G.U.T.

Moreover the correct interpretation of Y_F can be obtained if we consider a fundamental representation of G (if $\mathbf{U}(1)_F \subset G$). In this case we put fermion fields into

the representation and eigenvalues of Y_F tell us what is the meaning of F . In the case of $\mathbf{U}(1)_F \subset H$ and $\mathbf{U}(1)_F \not\subset G$ the interpretation of F is more complex. We should consider a fundamental representation of H and try to span its space by fermion fields looking for eigenvalues of Y_F . However the problem of fermion fields in the Nonsymmetric Jordan–Thiry Theory seems to be interesting and it will be done elsewhere.

We do not identify our group G with $\mathrm{SU}(5)$ or $\mathrm{SO}(10)$ from G.U.T. The relationship between these classical G.U.T.’s and dimensional reduction procedures seems to be more complex (see Ref. [80]). The feature concerning the second possibility of the place of the $\mathbf{U}(1)_F$ group seems to be more general than for $\mathbf{U}(1)_F$ only. The dimensional reduction scheme offers us a treatment of global symmetries in elementary particle physics. Let S be a group such that the group H has a structure:

$$S \subset H, \quad S \not\subset G, \quad G \otimes G_0 \subset H. \quad (11.32)$$

We have here S in a place of $\mathbf{U}(1)_F$ in (11.29). Due to the Wang condition we do not get any gauge field corresponding to the gauge group S . This group is really global. Thus the dimensional reduction scheme allows us to consider both global and local symmetries. Both symmetries can be unified by the one group H . The group S acts, of course, on the Higgs’ field Φ and would be broken spontaneously, without the Higgs’ mechanism. The group S is the symmetry group of the potential V . Thus we should get after symmetry breaking from G to G_0 massless scalars, the so-called pseudo-Goldstone bosons. The number of those zero modes is equal or greater $\dim S$. They can get masses through radiative corrections. The correct interpretation of generators of the Lie algebra of S can be obtained in a similar way as Y_F . In the next section we consider in details the first possibility (1)).

12. Fermion Number as the Second Gravitational Charge

$\overline{W}_\mu$ and $\tilde{A}^F_\mu$ Potentials

Let us consider the principal fibre bundle $\hat{P}$ of frames over the manifold E (space-time) with the structural group

$$\mathrm{GL}_\downarrow(4, \mathbb{R}) = \{(A^\alpha_\beta), \det(A^\alpha_\beta) > 0\}. \quad (12.1)$$

Let $\hat{\pi}$ be a projection $\hat{\pi} : \hat{P} \rightarrow E$. For every $x \in E$ we have that

$$F_x = \hat{\pi}^{-1}(\{x\}) \cong \mathrm{GL}_\downarrow(4, \mathbb{R}), \quad (12.2)$$

i.e., the fibre F_x is equivalent to the structural group $\mathrm{GL}_\downarrow(4, \mathbb{R})$. Let $\mathcal{W}$ be a connection on this bundle and let us decompose the Lie group $\mathrm{GL}_\downarrow(4, \mathbb{R})$ into

$$\mathrm{GL}_\downarrow(4, \mathbb{R}) = \mathbb{R}_+ \otimes \mathrm{SL}(4, \mathbb{R}), \quad (12.3)$$

where

$$\mathrm{SL}(4, \mathbb{R}) = \{A \in \mathrm{GL}(4, \mathbb{R}), \det A = 1\}$$

and

$$\mathbb{R}_+ = \left\{ e^\rho, \rho = \frac{1}{4} \ln(\det A), A \in \mathrm{GL}_\downarrow(4, \mathbb{R}) \right\}.$$

The decomposition (12.3) can be used to construct two subbundles of $\hat{P} : \hat{P}_1$ and $\hat{P}_2$ see Fig. 7. They are defined over E with structural groups $\mathbb{R}_+$ and $\mathrm{SL}(4, \mathbb{R})$. For $\mathrm{SL}(4, \mathbb{R})$ is a closed subgroup of $\mathrm{GL}_\downarrow(4, \mathbb{R})$, the transport of the connection from $\hat{P}$ to $\hat{P}_1$ and $\hat{P}_2$ is possible (see Ref. [59]).

In general the condition for a reduction of the principal bundle $P(M, G)$ to a subbundle $P'(M, G')$ where G' is a Lie subgroup of G is as follows. There is an open covering $\{U_\alpha\}$ of M with a set of transition functions $\{\Psi_{\beta\alpha}\}$ with values in G' . We suppose that this condition is satisfied in our cases (see Ref. [59]).

In this way we have, for the connection $\mathcal{W}$, a decomposition

$$\overline{\mathcal{W}} = \widetilde{\mathcal{W}} \oplus \widehat{\mathcal{W}}, \quad (12.4)$$

where $\widetilde{\mathcal{W}}$ is a 1-form with values in the Lie algebra $\mathrm{SL}(4, \mathbb{R})$ and $\widehat{\mathcal{W}}$ is a 1-form with values in the Lie algebra of $\mathbb{R}_+$. $\widetilde{\mathcal{W}}$ is a connection on $\hat{P}_2$ and $\widehat{\mathcal{W}}$ on $\hat{P}_1$. Thus we have

$$\widehat{\mathcal{W}} = \mathcal{W}Z, \quad (12.5)$$




Fig. 7. Principal fibre bundles $\hat{P}_1$, $\hat{P}_2$, Q_F and $\hat{P}$

where $\mathcal{W}$ is a 1-form with real values and Z is a generator of the Lie algebra $\mathbb{R}_+$. Let us take a local section f of the bundle $\hat{P}$

$$f : E \rightarrow \hat{P}. \quad (12.6)$$

One gets:

$$f^* \overline{\mathcal{W}} = (\overline{W}^\alpha_{\beta\gamma} \overline{\theta}^\gamma) X^\beta_\alpha, \quad (12.7)$$

where X^β_α , are generators of the Lie algebra of the group $\text{GL}_\downarrow(4, \mathbb{R})$. We can define the second connection $\widetilde{\mathcal{W}}$ (coefficients of a connection $\widetilde{\mathcal{W}}^\alpha_\beta$) according to Eq. (12.7) in terms of $\overline{W}^\alpha_\beta$.

Let us consider the general transformation law for the coefficients of the connection $\overline{W}^\alpha_\beta$

$$A^\lambda_\mu \overline{W}'^\mu_\nu = \overline{W}^\lambda_\mu A^\mu_\nu + dA^\lambda_\mu, \quad (12.8)$$

where A^λ_μ is a matrix valued function from $\text{GL}_\downarrow(4, \mathbb{R})$ and a frame transforms as

$$\overline{\theta}^\alpha = A^\alpha_\beta \overline{\theta}'^\beta. \quad (12.9)$$

If we choose

$$A^\alpha_\beta = e^{-2/3\phi} \delta^\alpha_\beta \quad (12.10)$$

we get

$$\overline{W}'^\lambda_\mu = \overline{W}^\lambda_\mu - \frac{2}{3} \delta^\lambda_\mu d\phi \quad (12.11)$$

or

$$\overline{W}'^\lambda{}_\mu = \overline{\omega}^\lambda{}_\mu - \frac{2}{3}\delta^\lambda{}_\mu \overline{W}', \quad (12.12)$$

where

$$\overline{W}' = \overline{W} + d\phi \quad (12.13)$$

and

$$\overline{\omega}^\lambda{}_\mu = f^* \widetilde{\mathcal{W}}^\lambda{}_\mu, \quad \widetilde{\mathcal{W}} = \widetilde{\mathcal{W}}^\lambda{}_\mu X^\mu{}_\lambda.$$

Thus it is possible to consider (12.10) and (12.13) as a gauge transformation for the 1-form $\overline{W}$. This was done in Refs. [34], [26], [27], [47], [65], [66]. It is easy to see that the 2-form of the curvature for the connection $\overline{\mathcal{W}}^\alpha{}_\beta$, $\Omega^\alpha{}_\beta(\overline{\mathcal{W}})$ is invariant under this transformation (see Refs. [26], [27]).

We can call this transformation the Einstein- λ -transformation (see Ref. [3]). This is also true for the curvature of the $\mathcal{W}^{\tilde{A}}{}_{\tilde{B}}$ connection, i.e. $\Omega^{\tilde{A}}{}_{\tilde{B}}$ (see Refs. [26], [27]). Let us consider the decomposition (12.4) for the connection $\overline{\mathcal{W}}$ corresponding to the decomposition (12.3) for the group $\text{GL}_\downarrow(4, \mathbb{R})$. Thus for every local section f of the bundle $\widehat{P}$ we get:

$$f^* \overline{\mathcal{W}} = \overline{\omega}^\lambda{}_\mu \widetilde{X}^\mu{}_\lambda + \overline{W} Z, \quad (12.14)$$

$\widetilde{X}^\mu{}_\lambda$ are generators of the Lie algebra of $\text{SL}(4, \mathbb{R})$. This means that we have chosen the decomposition of the Lie algebra of $\text{GL}_\downarrow(4, \mathbb{R})$ into $\text{SL}(4, \mathbb{R})$ and $\mathbb{R}$ ($\mathbb{R}$ is a real axis). Thus for every section of $\widehat{P}$ or $\widehat{P}_2$ and $\widehat{P}_1$ one gets

$$f^* \widetilde{\mathcal{W}} = \overline{\omega}^\lambda{}_\mu \widetilde{X}^\mu{}_\lambda, \quad (12.15)$$

$$f^* \widehat{\mathcal{W}} = \overline{W}_v \overline{\theta}^v Z. \quad (12.16)$$

Let us consider the decomposition (11.3) and compare it with formula (11.15). We have

$$\phi = -\frac{3}{8} \ln(\det A). \quad (12.17)$$

The group $\mathbb{R}_+$ is called the dilatation group. Thus the gauge transformation (12.13) for the 1-form $\overline{W}$ is connected to the dilatation subgroup of $\text{GL}_\downarrow(4, \mathbb{R})$. Let us consider the principal bundle Q (see Fig. 2A) with the structural group G and with a projection π_E over E (space-time). On this bundle we have defined a 1-form of a connection $\widetilde{\omega}_E$. (see Fig. 7).

Let $\mathbf{U}(1)_F$ be a subgroup of the group G such that

$$\mathbf{U}(1)_F \not\subset G_0 \quad (12.18)$$

and let us call this group the fermion charge group. Eq. (12.18) means that this subgroup is broken spontaneously and the Higgs' mechanism makes massive the gauge boson corresponding to this group. Let $Y_F \in \mathfrak{g}$ be a generator of $\mathbf{U}(1)_F$. For $\mathbf{U}(1)_F$ is

broken spontaneously $Y_F \in \mathfrak{m}$. Let us suppose that there exists the following homomorphism between $\mathrm{GL}_{\downarrow}(4, \mathbb{R})$ and G

$$\sigma : \mathrm{GL}_{\downarrow}(4, \mathbb{R}) \rightarrow G, \quad (12.19)$$

such that

$$\sigma(A) = e^{-(i\frac{3}{8} \ln(\det A) Y_F)}. \quad (12.20)$$

In this way we project the group $\mathrm{GL}_{\downarrow}(4, \mathbb{R})$ on the $\mathbf{U}(1)_F$ subgroup of G . The Eq. (12.20) is really the smooth homomorphism between $\mathbb{R}_+$ and $\mathbf{U}(1)_F$. Simultaneously (12.20) can be made a smooth morphism between typical fibres of the bundles $\hat{P}$ and Q . This suggests the following. Let us consider the morphism of the bundles $\hat{P}$ and Q

$$\Sigma : \hat{P} \rightarrow Q \quad \text{or} \quad (\Sigma_1 : \hat{P}_1 \rightarrow Q_F), \quad (12.21)$$

such that we have (12.20) and on the space-time E (base manifold)

$$\mathrm{id}_E : E \rightarrow E, \quad (12.22)$$

where id_E means the identity on E . The morphism Σ has the following meaning. The fibre bundle $\hat{P}$ is reduced to the principal fibre bundle $\hat{P}_1$ over E with a structural group $\mathbb{R}_+$. Simultaneously the principal fibre bundle Q is reduced to the $\mathbf{U}(1)_F$ fibre bundle over E i.e. Q_F . This is possible because of the property of the $\mathbf{U}(1)_F$ subgroup of G and the definition of $\mathbb{R}_+$ subgroup of $\mathrm{GL}_{\downarrow}(4, \mathbb{R})$. The transport of the connections of both bundles to their subbundles is possible. We can write our construction as a chain of morphisms:

$$\hat{P} \xrightarrow{r_1} \hat{P}_1 \xrightarrow{\Sigma_1} Q_F \xleftarrow{r_2} Q,$$

where r_1 means a reduction of $\hat{P}$ to $\hat{P}_1$, r_2 a reduction of Q to Q_F and Σ_1 a morphism induced by (12.20). Let us consider the contragradient transformation for Σ (a pull-back), Σ^* and apply it for ω_E . One gets:

$$\Sigma^*(\tilde{\omega}_E) = \widehat{\mathcal{W}} \quad (12.23)$$

or

$$\Sigma^*(\tilde{\omega}_E) = \Sigma^*(\tilde{\omega}_E^F Y_F) = \mathcal{W}Z. \quad (12.24)$$

Let us take a local section f of $\hat{P}$. One finds

$$-\frac{3}{2} \tilde{A}^F_{\mu} \bar{\theta}^{\mu} = \overline{W}_{\mu} \bar{\theta}^{\mu}. \quad (12.25)$$

Thus we get a relationship between the gauge field $\tilde{A}^F_{\mu}$, corresponding to the fermion number F , and the field $\overline{W}_{\mu}$ which couples the fermion current in the nonsymmetric theory of gravitation. For the curvature of the connection $\tilde{\omega}_E$, $\tilde{\Omega}_E$ one gets:

$$\Sigma^*(\tilde{\Omega}_E) = \Sigma^*(\tilde{H}^F Y_F) = \Sigma^*(d\tilde{\omega}_E^F Y_F) = \hat{\Omega} = d\widehat{\mathcal{W}} = d\mathcal{W}Z. \quad (12.26)$$

One easily finds that:

$$-\frac{4}{3}\overline{W}_{[\mu,\nu]} = \tilde{F}^F_{\mu\nu} = \partial_\mu \tilde{A}^F_\nu - \partial_\nu \tilde{A}^F_\mu. \quad (12.26a)$$

In this way, we can make the following statements:

- 1) local gauge invariance on the space-time E connected to the $\mathbb{R}_+$ group is the same as the local gauge invariance $\mathbf{U}(1)_F$ for the fermion number.
- 2) the gauge field $\tilde{A}^F_\mu$ and $\overline{W}_\mu$ are the same. We have that

$$\overline{W}^\lambda{}_\mu = \overline{\omega}^\lambda{}_\mu + \delta^\lambda_\mu (\tilde{A}^F_\nu \overline{\theta}^\nu) \quad (12.27)$$

and in the gravitational lagrangian

$$-\frac{4}{3} g^{[\mu\nu]} \overline{W}_{[\mu,\nu]} = g^{[\mu\nu]} \tilde{H}^F_{\mu\nu} \quad (12.28)$$

Due to the spontaneous symmetry breaking the field $\tilde{A}^F_\mu$ is massive and has short distance. Thus the Lorentz force term which appears for $W_{[\mu,\nu]}$ is of short distance

$$\frac{2}{3} f \tilde{g}^{(\mu\nu)} W_{[\mu,\alpha]} u^\alpha = -2 f \tilde{g}^{(\mu\nu)} \tilde{H}^F_{\mu\alpha} u^\alpha, \quad (12.29)$$

(see Ref. [95]). In this way the weak equivalence principle is satisfied. This is really impossible to do in pure N.G.T., where the Lorentz force term (or Coriolis force term) appears as a term of long distance forces and it is necessary to remove it using the so called splitting of the conservation laws (see Refs. [34], [47], [65]). We do not need these additional assumptions, which are not of geometric nature. Moreover, the splitting of the conservation laws is an integral part of Moffat's theory. It seems that it provides a contradiction with the field equations with electromagnetic or scalar sources in N.G.T. (see Ref. [34]). It seems that it is also in contradiction with the Lagrange formulation of hydrodynamics. However the splitting of the conservation laws is necessary in N.G.T. in order to get the equation of motion for the test particle from conservation laws (Bianchi identities) as in G.R.T. In this manner Moffat fits the anomalous perihelion precession of Mercury and Icarus in the presence of a non-zero quadrupole moment of the sun

Otherwise, he gets an equation with a Lorentz force term (Coriolis force term) for $W_{[\mu,\nu]}$.

In some new formulations of N.G.T. (Ref. [95]) J.W. Moffat abandons the splitting of the conservation laws getting equation of motion for a test particle with Lorentz-like (Coriolis force term).

The additional force term in the equation of motion has been connected to the "fifth force" (Refs. [54], [53]). However, it seems that the fifth force term (if this force really exists) should be of exponential decaying, not $\frac{1}{r^5}$ as in Moffat's approach ([95]). In our meaning the "fifth force" is rather scalar than vector-like.

The conditions (12.23), (12.24), (12.25) are really constraints in the theory. Thus we should add to the full lagrangian the term:

$$\frac{8\pi G_N}{c^4} \tilde{L}^\mu \left(\tilde{A}^F_\mu + \frac{2}{3} \overline{W} \right), \quad (12.30)$$

where

$$\tilde{\lambda}^\mu = \sqrt{-g} \lambda^\mu \quad (12.31)$$

is a Lagrange multiplier and because of the gauge invariance ($\mathbb{R}_+$ and $\mathbf{U}(1)_F$) we suppose

$$\partial_\mu \tilde{\lambda}^\mu = 0. \quad (12.32)$$

The Lagrange multiplier becomes a source for $g_{[\mu\nu]}$

$$g^{[\mu\nu]}_{,\nu} = -\kappa \tilde{\lambda}^\mu = 4\pi a^2 \tilde{S}^\mu, \quad \kappa = \frac{8\pi G_N}{c^4}. \quad (12.33)$$

Thus it has a natural physical interpretation as a fermion current. It is coupled to the $\tilde{A}^F_\mu$ and simultaneously to $\overline{W}_\mu$. In this way the fermion charge plays the role of the second gravitational charge. The field $\overline{W}_\mu$ disappears from the theory. In the place of $\overline{W}_\mu$ we have $\tilde{A}^F_\mu$ which is massive and due to this the weak equivalence principle is satisfied. The Lorentz force term (Coriolis force term) for $\tilde{A}^F_\mu$ is of short range. From the Nonsymmetric Gravitational Theory we know that there are not any Lorentz forces and Coulomb potentials connected to $g_{[\mu\nu]}$. Thus the weak equivalence principle is satisfied and we have two gravitational charges: a mass and a fermion charge.

In this way the argument raised by C. Will (see 4th of Ref. [47]) does not concern our case. The arguments is of course valid in the case of N.G.T., because there $\overline{W}$ is not of a short range.

Let us notice that we choose the second possibility of the role of $\mathbf{U}(1)_F$ in our scheme from Ref. [26] and sec. 11. This corresponds to the situation similar for SO(10) G.U.T., i.e. the first possibility from Ref. [26] and sec. 11. It seems that this possibility is more natural. Some possibilities concerning the intermediate stages of the symmetry breaking will be considered elsewhere.

Finally we can say that the $SU(5)$ scheme of G.U.T. seems to be in contradiction with experiment (proton decay). For this kind of scheme we cannot proceed with the construction presented here. $\mathbf{U}(1)_F$ is global and $Y_F \in (\text{Lie algebra of the group } H)$, and $Y_F \notin$.

Let us conclude that in this section we have obtained the physical interpretation of the Einstein- λ -transformation (Ref. [3]). It seems that this transformation is a local $\mathbf{U}(1)_F$ gauge transformation corresponding to the fermion-number generator in G.U.T. In this way the problem seems to be solved. The Einstein Unified Field Theory in a real version was used as a theory of the pure gravitational field. It was necessary to

combine it with the Yang–Mills' and Higgs' field via the Nonsymmetric Jordan–Thiry Theory.

Let us consider the lagrangian of the theory and find his part which contains the gauge field $\tilde{A}^F_\mu = -\frac{2}{3}\overline{W}_\mu$. In order to do this we consider Eq. (6.1) with the $\tilde{H}^F_{\mu\nu}$ contribution. One gets:

$$\tilde{Q}^j_i g_{\delta\beta} g^{\sigma\delta} \tilde{L}^i_{\sigma\alpha} + g_{\alpha\delta} g^{\sigma\delta} \tilde{L}^j_{\beta\sigma} = 2g_{\alpha\delta} g^{\delta\sigma} \tilde{H}^j_{\beta\sigma}, \quad (12.34)$$

where

$$\tilde{Q}^j_i = \tilde{\ell}^{kj} \tilde{\ell}_{ik}.$$

If $j = F$ we have

$$\tilde{Q}^F_i g_{\delta\beta} g^{\sigma\delta} \tilde{L}^i_{\sigma\alpha} + g_{\alpha\delta} g^{\delta\sigma} \tilde{L}^F_{\beta\sigma} = 2g_{\alpha\delta} g^{\delta\sigma} \tilde{H}^F_{\beta\sigma}. \quad (12.35)$$

For

$$\tilde{H}^F_{\mu\nu} = -\frac{4}{3}W_{[\mu,\nu]} \quad (12.36)$$

one gets

$$Q^F_i g_{\delta\beta} g^{\sigma\delta} \tilde{L}^i_{\sigma\alpha} + g_{\alpha\delta} g^{\delta\sigma} \tilde{L}^F_{\beta\sigma} = -\frac{8}{3}g_{\alpha\delta} g^{\delta\sigma} W_{[\beta,\sigma]}. \quad (12.37)$$

This means that the field W_μ has a contribution to the full induction tensor of the theory.

The Yang–Mills' field lagrangian with a direct contribution of W_μ looks as follows:

$$\mathcal{L}_{\text{YM}}(W) = -\frac{1}{6\pi} \tilde{\ell}_{\text{FF}} \left[\frac{8}{3} (g^{[\mu\nu]} \overline{W}_{[\mu,\nu]})^2 + \tilde{L}^{F\mu\nu} \overline{W}_{[\mu,\nu]} \right]. \quad (12.38)$$

There is also a usual part of N.G.T. lagrangian containing $\overline{W}_\mu$ field i.e.

$$\frac{2}{3} g^{[\mu\nu]} \overline{W}_{[\mu,\nu]}. \quad (12.39)$$

We have also a contribution of the field $\overline{W}_\mu$ in the kinetic part of the lagrangian for a Higgs' field if the gauge derivative of the Higgs' field contains $\tilde{A}^F_\mu = -\frac{2}{3}\overline{W}_\mu$ i.e.

$$\begin{aligned} e^* (\overset{\text{gauge}}{\nabla}_\mu \Phi^a_{\tilde{a}}) &= \partial_\mu \Phi^a_{\tilde{a}} + \frac{g}{\hbar c} (C^a_{de} \alpha^e_j \tilde{A}^j_\mu \Phi^a_{\tilde{a}} + f^{\tilde{b}}_{\tilde{a}j} \tilde{A}^j_\mu \Phi^a_{\tilde{b}}) = \\ &= \partial_\mu \Phi^a_{\tilde{a}} - \frac{3g}{2\hbar c} (C^a_{de} \alpha^e_F \overline{W}_\mu \Phi^a_{\tilde{a}} + f^{\tilde{c}}_{\tilde{a}F} \overline{W}_\mu \Phi^a_{\tilde{c}}) + \\ &\quad + \frac{g}{\hbar c} \sum_{j \neq F} (C^a_{de} \alpha^e_j \tilde{A}^j_\mu \Phi^a_{\tilde{a}} + f^{\tilde{c}}_{\tilde{a}j} \tilde{A}^j_\mu \Phi^a_{\tilde{c}}) = \\ &= \overset{\text{gauge}}{\nabla}_\mu \Phi^a_{\tilde{a}} + \frac{g}{\hbar c} \sum_{j \neq F} (C^a_{de} \alpha^e_j \tilde{A}^j_\mu \Phi^a_{\tilde{a}} + f^{\tilde{c}}_{\tilde{a}j} \tilde{A}^j_\mu \Phi^a_{\tilde{c}}). \end{aligned} \quad (12.40)$$

Thus one gets:

$$\mathcal{L}_{\text{kin}}^W(\nabla_\mu^{\text{gauge}} \Phi) = (\nabla_\mu^w \Phi_a^{\text{gauge}} L^{\mu\tilde{a}}{}_a)_{av}. \quad (12.41)$$

Taking everything together we have:

$$\begin{aligned} \mathcal{L}(\overline{W}) &= \frac{2}{3} g^{[\mu\nu]} \overline{W}_{[\mu,\nu]} - e^{-(n+2)\Psi} \frac{\lambda^2}{3} \tilde{\ell}_{\text{FF}} \left[\frac{8}{3} (g^{[\mu\nu]} \overline{W}_{[\mu,\nu]})^2 + \tilde{L}^{\text{F}\mu\nu} \overline{W}_{[\mu,\nu]} \right] + \\ &\quad - \frac{\lambda^2}{2r^2} e^{-2\Psi} (\nabla_\mu^w \Phi_a^{\text{gauge}} L^{\mu\tilde{a}}{}_a)_{av}. \end{aligned} \quad (12.42)$$

However because of the Eq. (6.1) and Eq. (6.4) we have the $\overline{W}_\mu$ field contribution via an induction tensor $\tilde{L}^i{}_{\mu\nu}$ and $L^{\mu\tilde{a}}{}_a$ in the remaining parts of $\mathcal{L}_{\text{YM}}$ and $\mathcal{L}^{\text{kin}}(\nabla_\mu^{\text{gauge}} \Phi)$. Finally let us rewrite the field equations *modulo* constraint $\tilde{A}^{\text{F}}{}_\mu = -\frac{2}{3} \overline{W}_\mu$.

$$\overline{R}_{(\alpha\beta)}(\overline{T}) = \frac{8\pi K}{c^4} \left(T_{(\alpha\beta)}^{\text{tot}} - \frac{1}{2} g_{(\alpha\beta)} T \right), \quad (12.43)$$

$$\overline{R}_{[[\alpha\beta],\lambda]}(\overline{T}) = \frac{8\pi}{c^4} \left(K T_{[\alpha\beta]}^{\text{tot}} - \frac{1}{2} g_{[[\alpha\beta]} T \right)_{,\lambda}, \quad (12.44)$$

where

$$T_{\alpha\beta}^{\text{tot}} = T_{\alpha\beta}^{\text{gauge}} + T_{\alpha\beta}^{\text{scal}} + T_{\alpha\beta}(\Phi) + T_{\alpha\beta}^{\text{int}} + \Lambda g_{\alpha\beta} \quad (12.45)$$

and

$$\begin{aligned} T &= g^{\alpha\beta} T_{\alpha\beta}^{\text{tot}}, \\ \tilde{g}^{[\mu\nu]}{}_{,\nu} &= 4\pi \tilde{S}^\mu, \end{aligned} \quad (12.46)$$

$$g_{\mu\nu,\sigma} - g_{\rho\nu} \tilde{\Lambda}^\rho{}_{\mu\sigma} - g_{\mu\rho} \tilde{\Lambda}^\rho{}_{\sigma\nu} = 0, \quad (12.47)$$

where

$$\begin{aligned} \tilde{S}^\mu &= \frac{3\lambda^2 \alpha_S^2}{c\hbar} \sqrt{-g} \frac{\delta}{\delta \overline{W}_\mu} \left(e^{-(n+2)\Psi} \mathcal{L}_{\text{YM}} + \frac{e^{-2\Psi}}{4\pi r^2} \mathcal{L}^{\text{kin}}(\nabla_\mu^{\text{gauge}} \Phi) + \right. \\ &\quad \left. - \frac{e^{(n-2)\Psi}}{2\pi r^4} \mathcal{L}_{\text{int}}(\Phi, \tilde{A}) \right), \end{aligned} \quad (12.48)$$

$$\tilde{\Lambda}_{\mu\sigma} = \tilde{\Gamma}^\rho{}_{\mu\sigma} + D^\rho{}_{\mu\sigma}(S) \quad (12.49)$$

and

$$g_{\rho\nu} D^\rho{}_{\mu\sigma} + g_{\mu\rho} D^\rho{}_{\sigma\nu} = \frac{4\pi}{3} S^\rho (g_{\mu\sigma} g_{\rho\nu} - g_{\mu\rho} g_{\sigma\nu} + g_{\mu\nu} g_{[\sigma\rho]}), \quad (12.50)$$

$\tilde{\tilde{R}}_{\alpha\beta}(\tilde{\tilde{T}})$ is a Moffat–Ricci tensor for a connection $\tilde{\tilde{\omega}}^\alpha{}_\beta = \tilde{\tilde{\Gamma}}^\alpha{}_{\beta\gamma} \theta^\gamma$. It is easy to see that we have sources for equation (9.4) and (9.5). This is similar to the situation with the second pair of Maxwell equations in the Nonsymmetric Kaluza–Klein Theory in comparison to the classical Kaluza–Klein Theory.

The above equations have been obtained as results of variations with respect to $g_{\mu\nu}$ and

$$\overline{W}^\lambda{}_\mu = \tilde{\omega}^\lambda{}_\mu - \frac{2}{3}\delta^\lambda{}_\mu \overline{W} = \tilde{\omega}^\lambda{}_\mu + \delta^\lambda{}_\mu \tilde{A}^F.$$

Moreover if we vary with respect to

$$\tilde{\omega}_E = \tilde{\omega}_E^F X_F + \sum_{i \neq F} \tilde{\omega}_E^i X_i$$

the variation with respect to the first part of ω is included in the variation of $\overline{W}^\lambda{}_\mu$ because of

$$e^* \tilde{\omega}_E^F = \tilde{A}^F = -\frac{2}{3} \overline{W}, \quad (12.51)$$

Thus we get Eq. (9.7) for $j \neq F$ only.

We can easily calculate the form of S^μ . One gets

$$\begin{aligned} \tilde{S}^\alpha = & \frac{-8G_N \hbar}{c^4} \left\{ \nabla_\mu^{\text{gauge}} (\tilde{\ell}_{iF} \tilde{L}^{i\alpha\mu}) + 2 \tilde{g}^{[\alpha\beta]} \nabla_\beta^{\text{gauge}} (\tilde{h}_{iF} g^{[\mu\nu]} \tilde{H}^i{}_{\mu\nu}) + \right. \\ & + \frac{2\sqrt{-g}}{r^2} \alpha_S \sqrt{\hbar c} e^{n\Psi} \left[\ell_{ab} g^{\tilde{b}\tilde{n}} g^{\mu\alpha} L^a{}_{\mu\tilde{b}} (\Phi_{\tilde{n}}^d C^b{}_{dc} \alpha_F^c + f_{n\tilde{F}}^{\tilde{e}} \Phi_{\tilde{e}}^b + \right. \\ & + \left. \left(\frac{\delta L_{\beta\tilde{b}}^a}{\delta \nabla_\alpha^{\text{gauge}} \Phi_{\tilde{v}}^w} \right) \ell_{ab} g^{\tilde{b}\tilde{n}} g^{\beta\mu} (\nabla_\mu^{\text{gauge}} \Phi_{\tilde{n}}^b) (\Phi_{\tilde{w}}^d C^w{}_{dc} \alpha_F^c + f_{\tilde{w}F}^{\tilde{e}} \Phi_{\tilde{e}}^w) \right]_{av} + \\ & + 4\sqrt{-g} \frac{e^{2n\Psi}}{r^2} h_{ab} \mu_k^a \tilde{\ell}_{iF} \tilde{\ell}^{ki} \tilde{g}^{[a\tilde{b}]} \nabla_\mu^{\text{gauge}} \left\{ g^{[\mu\alpha]} \times \right. \\ & \times \left[\frac{\alpha_s}{\sqrt{\hbar c}} C_{cd}^b \Phi_{\tilde{a}}^c \Phi_{\tilde{b}}^d - \frac{\hbar c}{\alpha_s} \mu_{\tilde{a}\tilde{b}}^b f_{\tilde{a}\tilde{b}}^{\tilde{i}} - \phi_{\tilde{a}}^b f_{\tilde{a}\tilde{b}}^{\tilde{d}} \right] \Big\} + \\ & + \left. (n+2) \partial_\beta \Psi [\tilde{\ell}_{iF} \tilde{L}^{i\beta\alpha} - 2 \tilde{g}^{[\beta\alpha]} (\tilde{h}_{iF} g^{[\mu\nu]} \tilde{H}^i{}_{\mu\nu})] \right\}. \end{aligned} \quad (12.52)$$

The remaining equations i.e. Eqs. (9.6), (9.8) are the same substituting for $\tilde{H}^F{}_{\mu\nu}$, $-\frac{4}{3} \overline{W}_{[\mu,\nu]}$ and for $\tilde{A}^F{}_\mu$, $-\frac{2}{3} \overline{W}_\mu$.

13. Equation of Motion for a Test Particle in the Nonsymmetric Jordan–Thiry Theory

Let us derive the equation of motion for a test particle in our theory from the Bianchi identities in the hydrodynamic limit.

In N.G.T. one derives the following Bianchi identity (see Refs. [5], [97], [98])

$$\frac{1}{2}(G_{\rho\nu} \, \underset{\sim}{g}^{\alpha\nu})_{,\alpha} + \frac{1}{2}(G_{\nu\rho} \, \underset{\sim}{g}^{\nu\alpha})_{,\alpha} + \frac{1}{2} \underset{\sim}{G}_{\varepsilon\nu} \, g^{\varepsilon\nu}_{,\rho} = 0, \quad (13.1)$$

where

$$G_{\rho\nu} = \overline{R}_{\rho\nu}(\overline{\Gamma}) - \frac{1}{2}g_{\rho\nu}\overline{R}(\overline{\Gamma}), \quad (13.2)$$

$$\underset{\sim}{G}_{\varepsilon\nu} = \sqrt{-g}G_{\varepsilon\nu}, \quad (13.3)$$

$\overline{R}_{\rho\nu}(\overline{\Gamma})$ is the Moffat–Ricci tensor for the connection $\overline{\omega}^\alpha_\beta$ and $R(\overline{\Gamma})$ is the scalar of curvature.

From Eq. (9.3) one gets:

$$R_{\alpha\rho}(\overline{W}) - \frac{1}{2}g_{\alpha\rho}R(\overline{W}) = \frac{8\pi G}{c^4}e^{-(n+2)\Psi} \overset{\text{tot}}{T}_{\alpha\beta}, \quad (13.4)$$

where

$$\overset{\text{tot}}{T}_{\alpha\rho} = \overset{\text{gauge}}{T}_{\alpha\rho} + T_{\alpha\rho}(\Phi) + \overset{\text{scal}}{T}_{\alpha\rho} + g_{\alpha\rho}\Lambda + \overset{\text{int}}{T}_{\alpha\rho}. \quad (13.5)$$

If we suppose the constraints (12.30) we have to use an effective energy-momentum tensor

$$\overset{\text{eff}}{T}_{\alpha\rho} = \overset{\text{tot}}{T}_{\alpha\rho} + \overset{\text{constraints}}{T}_{\alpha\rho}, \quad (13.6)$$

where

$$\overset{\text{constraints}}{T}_{\alpha\rho} = e^{(n+2)\Psi} \left[\lambda_\alpha \left(\overline{W}_\rho + \frac{2}{3}\tilde{A}^{\text{F}}_\rho \right) - \frac{1}{2}g_{\alpha\rho}\lambda^\mu \left(\overline{W}_\mu + \frac{2}{3}\tilde{A}^{\text{F}}_\mu \right) \right]. \quad (13.7)$$

Now Eq. (13.4) takes the form:

$$R_{\alpha\rho}(\overline{W}) - \frac{1}{2}g_{\alpha\rho}R(\overline{W}) = \frac{8\pi G}{c^4}e^{-(n+2)\Psi} \overset{\text{eff}}{T}_{\alpha\rho}. \quad (13.8)$$

Simultaneously for Eq. (9.4) we have

$$\underset{\sim}{g}^{[\mu\nu]}_{,\nu} = \frac{3}{2} \underset{\sim}{\lambda}^\mu = 4\pi a^2 \underset{\sim}{S}^\mu. \quad (13.9)$$

From Eq. (13.1) one derives:

$$[e^{-(n+2)\Psi}(g^{\alpha\nu}\overset{\text{eff}}{\mathcal{T}}_{\nu\rho} + g^{\nu\alpha}\overset{\text{eff}}{\mathcal{T}}_{\nu\rho}),_{\alpha} + g^{\mu\nu},_{\rho}e^{-(n+2)\Psi}\overset{\text{eff}}{T}_{\mu\nu} + \frac{2}{3}W_{[\rho,\nu]}\overset{\text{eff}}{\mathcal{S}}^{\nu} = 0. \quad (13.10)$$

Let us suppose that:

1) The field configuration described by Eqs. (9.3)–(9.8) and constraints (12.30) correspond to a situation close to the “true” vacuum case.

2) The hydrodynamic limit for this field configuration.

From 1) we have (see Eq. (10.12)):

$$e^{-(n+2)\Psi} \simeq \text{const}, \quad (13.11)$$

$$\Lambda \simeq 0, \quad (13.12)$$

$$-\frac{2}{3}\overline{W}_{[\mu,\nu]} = \tilde{A}^{\text{F}}_{[\mu,\nu]} \simeq \frac{1}{r^2}e^{-\beta r} \simeq 0, \quad (13.13)$$

where $\frac{1}{\beta}$ is the range of the gauge field $\tilde{A}^{\text{F}}_{\mu}$.

Thus from Eq. (13.10) one gets:

$$(g^{\alpha\nu}\overset{\text{eff}}{\mathcal{T}}_{\rho\nu} + g^{\nu\alpha}\overset{\text{eff}}{\mathcal{T}}_{\nu\rho}) + g^{\mu\nu},_{\rho}\overset{\text{eff}}{\mathcal{T}}_{\mu\nu} = 0. \quad (13.14)$$

2) means that our field configuration is described by the macroscopical (averaged) hydrodynamic quantities: u^{α} (four-velocity of a fluid), p (pressure), ρ (density), T (temperature) and $F_{\mu\nu}$ (macroscopic electromagnetic field) which is the only gauge field with long range.

$$\overset{\text{eff}}{T}_{\rho\nu} = \overset{\text{eff}}{T}_{\rho\nu}(u^{\alpha}, p, \rho, T, F_{\mu\beta}). \quad (13.15)$$

Thus Eq. (13.14) would be treated as an energy-momentum conservation law in N.G.T. for phenomenological sources.

Let us derive the equation for a test particle from Eq. (13.14). In order to do this let us go to the limit of one uncharged particle. This means:

$$p = 0, \quad F_{\mu\nu} = 0, \quad (13.16a)$$

$$T_{\rho\nu} = m_0 g_{\rho\alpha} g_{\beta\nu} u^{\alpha} u^{\beta}, \quad (13.16b)$$

$$du = d(\sqrt{-g}u_{\nu}\bar{\theta}^{\nu}) = 0. \quad (13.16c)$$

The last equation gives:

$$\overset{\text{eff}}{u}^{\alpha},_{\alpha} = 0, \quad (13.17)$$

where u^{α} is a four-velocity of a test particle and m_0 its mass. Putting Eqs. (13.16b) and (13.17) into (13.14) one gets after some algebra:

$$\frac{d^2 x^{\gamma}}{dt^2} + \left\{ \begin{matrix} \gamma \\ \alpha\beta \end{matrix} \right\} \left(\frac{dx^{\alpha}}{dt} \right) \left(\frac{dx^{\beta}}{dt} \right) = 0, \quad (13.18)$$

where $u^\alpha = \frac{dx^\alpha}{dt}$ and $\{\gamma_{\alpha\beta}\}$ are Christoffel symbols formed for the metric $g_{(\alpha\beta)}$.

Thus we get that a test particle (without spin and electric charge) moves along a geodetic line for the metric $g_{(\alpha\beta)}$. The Eq. (13.6) fits the anomalous perihelion shift of Mercury and Icarus in the presence of a nonzero mass quadrupole moment for the sun if we use the symmetric part of the spherically symmetric, static solution in N.G.T. (see Ref. [45]). In this limit we suppose that Mercury and Icarus can be treated as test particles in the gravitational field of the sun.

It works also very well in the case of the anomalous periastron movement of the closed binary system DI Hercules (DI Her i.e. HD175227), which contradicts G.R.T. predictions up to 21σ (standard deviations) (see Refs. [67], [99]). The equation fits the observational data. In the case of different closed binary systems (AG Per, AS Cam, α Vir, V541 Cyg, V1143 Cyg, V889 Aql) with strange movement of the periastron the equation can also fit the observational data. Using the equation under above conditions we can also fit the data for X -ray bursters (Ref. [49]).

Let us notice that the Eq. (13.15) could be treated as a fundamental equation in any post-Newtonian approximation for N.G.T. using the hydrodynamic formalism as in Ref. [100].

Finally let us consider the Newtonian limit for Ψ . Let us expand $e^{-(n+2)\Psi}$ into a power series:

$$e^{-(n+2)\Psi} \cong 1 - (n+2)\Psi + \dots \quad (13.19)$$

The field Ψ has a Yukawa-type behaviour i.e.

$$\Psi \cong C' + \frac{C}{r}e^{-\gamma r}, \quad (13.20)$$

where $\gamma, C', C = \text{const}$ and $\gamma, C > 0$. One gets:

$$e^{-(n+2)\Psi} \simeq (1 + C') - \frac{(n+2)C}{r}e^{-\gamma r}. \quad (13.21)$$

We can consider a correction to the Newtonian potential coming from the scalar field Ψ . One has:

$$V_{\text{eff}}(r) = -G_{\text{eff}} \frac{M}{r} = -(\tilde{G}e^{-(n+2)\Psi}) \frac{M}{r} = -G_N \frac{M}{r} + \frac{\alpha}{r^2}e^{-\gamma r}, \quad (13.22)$$

where $G_N = \tilde{G}(1 + C')$ plays the role of the Newton constant and

$$\alpha = \tilde{G}C(n+2) > 0. \quad (13.23)$$

Thus the correction to the Newtonian potential has a repulsive character. Putting Eq. (13.18) into linearized equation of motion for a test particle we can get a composition-dependent repulsive correction with a short range. This could be considered as something similar to the fifth force (Refs. [53], [54]). We can also try to get an equation

of motion from the conservation laws under conditions (13.17) and (13.19), i.e.

$$(g^{\alpha\nu} \overset{\text{eff}}{\tilde{T}}_{\rho\nu} + g^{\nu\alpha} \overset{\text{eff}}{\tilde{T}}_{\nu\rho}) + g^{\mu\nu}{}_{,\rho} - (n+2)\Psi g^{\mu\nu}{}_{,\rho} \overset{\text{eff}}{\tilde{T}}_{\mu\nu} = 0. \quad (13.24)$$

Recently J. W. Moffat and his coworkers have renamed ([10]) N.G.T. into N.G.E.T. (Nonsymmetric Gravitational and Electromagnetic Theory), taking the Lagrangian for the electromagnetism from Nonsymmetric Kaluza–Klein Theory ([18]). They found nonsingular solutions in the case for pure gravitational field with interesting physical interpretations ([102], [103]).

14. On a Cosmological Origin of the Mass of the Scalar Field Ψ (or ρ). Cosmological Models with Quintessence and Inflation

Let us consider the lagrangian in our theory supposing that Yang–Mills' and Higgs' fields are zero. One gets in terms of the field Ψ (see Eq. (7.5)):

$$\mathcal{L} = \left[\bar{R}(\bar{W}) + \frac{\lambda^2}{4} \left(-e^{(n-2)\Psi} \frac{V(0)}{r^4} - \mathcal{L}_{\text{scal}}(\Psi) + \frac{8}{\lambda^4} e^{(n+2)\Psi} \tilde{R}(\tilde{\Gamma}) + \frac{4e^{n\Psi}}{\lambda^2 r^2} \tilde{P} \right) \right]. \quad (14.1)$$

If we suppose that we have to do with a cosmological background i.e. $\bar{R}(\bar{W}) = R_0 = \text{const}$ and Ψ is constant we get in terms of $\rho = e^{-\Psi}$ ($\rho > 0$)

$$\mathcal{L} = \left[R_0 + \frac{\lambda^2}{4\rho^{n+2}} \left(-\rho^4 \frac{V(0)}{r^4} + \frac{8}{\lambda^4} \tilde{R}(\tilde{\Gamma}) + 4\rho^2 \frac{\tilde{P}}{\lambda^2 r^2} \right) \right] = W(\rho). \quad (14.2)$$

Eq. (14.2) is a self-interacting potential for a scalar field ρ . The potential has a critical point if

$$\frac{dW}{d\rho} = 0, \quad (14.3)$$

i.e.

$$-(n-1) \left(\frac{\lambda^2 V(0)}{4r^4} \right) \rho^4 + n \left(\frac{\tilde{P}}{4r^2} \right) \rho^2 + 4(n+2) \frac{\tilde{R}(\tilde{\Gamma})}{\lambda^2} = 0. \quad (14.4)$$

Eq. (14.4) has real positive solutions if

$$\tilde{P}^2 \geq - \left(\frac{16(n-1)(n+2)}{n^2} \right) \left(\frac{\tilde{R}(\tilde{\Gamma}) V(0)}{\lambda^2} \right) \quad (14.5a)$$

and

$$\frac{n\lambda^2 \tilde{P}}{V(0)} \leq \frac{\sqrt{n^2 \lambda^2 \tilde{P}^2 + 16(n-1)(n+2)V(0)\tilde{R}(\tilde{\Gamma})}}{V(0)}. \quad (14.5b)$$

In this case we have:

$$\rho_{0,1,2}^{\pm} = \pm \frac{r}{\lambda} \sqrt{\frac{\sqrt{n^2 \lambda^2 \tilde{P}^2 + 16(n-1)(n+2)V(0)\tilde{R}(\tilde{\Gamma})} \pm n\lambda \tilde{P}}{V(0)}} \quad (14.6)$$

i.e. in general two positive real roots. The potential W has a minimum at $\rho = \rho_0$ if

$$\left. \frac{d^2 W}{d\rho^2} \right|_{\rho=\rho_0} > 0 \quad (14.7)$$

i.e.

$$\left(\frac{-(n-1)(n-2)\frac{\lambda^2 V(0)}{4r^4}\rho^4 + \frac{n(n+1)\underline{P}}{4r^2}\rho^2 + (n+1)(n+2)\frac{4\tilde{R}(\tilde{\Gamma})}{\lambda^2}}{\rho^{n+2}} \right) \bigg|_{\rho=\rho_0} > 0. \quad (14.8)$$

All of these conditions could be easily satisfied if $\tilde{R}(\tilde{\Gamma}) = 0$ and $\frac{\tilde{P}}{V(0)} > 0$. The condition (14.8) can be satisfied for only 2 roots of Eq. (14.4). For $\tilde{R}(\tilde{\Gamma}) = 0$ Eq. (14.6) has the shape:

$$\rho_0^\pm = \pm \frac{r}{\lambda} \sqrt{\frac{n\tilde{P}}{(n-1)V(0)}}. \quad (14.6')$$

In terms of a scalar field Ψ one gets:

$$W(\rho(\Psi)) = W(\rho_0) + \frac{dW}{d\Psi} \bigg|_{\Psi=\Psi_0} \Psi' + \frac{1}{2} \frac{d^2 W}{d\Psi^2} \bigg|_{\Psi=\Psi_0} (\Psi')^2 + \dots, \quad (14.9)$$

where

$$\Psi = -\ln \rho_0 + \Psi' = \Psi_0 + \Psi'.$$

One finds

$$W(\Psi) = W(\rho_0) + \frac{\rho_0^2}{2} \frac{d^2 W}{d\rho^2} \bigg|_{\rho=\rho_0} (\Psi')^2 + \text{higher order terms in } \Psi'. \quad (14.10)$$

For (14.7) is satisfied we can write:

$$\rho_0^2 \frac{d^2 W}{d\rho^2} \bigg|_{\rho=\rho_0} = \frac{\lambda^2}{4} m_0^2 \overline{M} \quad (14.11)$$

and rewrite Eq. (14.2) in terms of Ψ' . One easily gets:

$$\mathcal{L} = R_0 + \frac{\lambda^2}{4} \left(\Lambda_0 + \frac{m_0^2}{2} \overline{M} (\Psi')^2 + \text{higher order terms} \right), \quad (14.12)$$

where

$$\frac{\lambda^2}{4} \Lambda_0 = W(\rho_0) - R_0,$$

is a new cosmological constant.

Thus one can write Eq. (14.1) in terms of Ψ' using Eq. (14.12).

$$\mathcal{L} = \bar{R}(\bar{W}) + \frac{\lambda^2}{4} \left(-\mathcal{L}_{\text{scal}}(\Psi') + \bar{M} \frac{m_0^2}{2} (\Psi')^2 + \text{higher order terms in } \Psi' + \Lambda \right). \quad (14.13)$$

Thus the cosmological background can induce a mass for the scalar field $\Psi(\rho)$ via usual mechanism known in solid state physics if (14.5a), (14.5b), (14.7) are satisfied. In this way a mass m_0 for Ψ' has a cosmological origin and the field Ψ or ρ has a short range (Yukawa-type) behaviour. Moreover we should redefine a field Ψ' .

The scalar field $\Psi(\rho)$ is connected to the effective gravitational constant and the mechanism presented in this section gives us a cosmological reason for a slow change of the gravitational constant. This mechanism gives also some explanation for the origin of the possible “fifth force” similar in spirit to the Fujii theory (see Ref. [96]). It is easy to see that if we choose $\rho_0 > 0$ the reflection symmetry of $W(\rho)$ is spontaneously broken.

Let us notice that the value R_0 does not enter to the formulae for ρ_0 and m_0 . Thus we can abandon in principle a constancy of R_0 and consider a slow changing in time of $\bar{R}$ i.e.

$$\bar{R}(t) = R_0 - 24\pi G_N d_0 H_0 (t - t_0), \quad (14.14)$$

where R_0 is the scalar curvature at the present epoch time t_0 , d_0 is a present rest mass galaxies density and H_0 is the Hubble constant.

Finally let us calculate the mass of the scalar field Ψ' and the cosmological constant Λ_0 .

$$m_0 = \frac{4}{\lambda} \sqrt{\frac{-(n-2)(n-1) \frac{\lambda^2 V(0)}{4r^4} \rho_0^4 + \frac{n(n+1) \tilde{P}}{4r^4} \rho_0^2 + (n+2)(n+3) \frac{4\tilde{R}(\tilde{\Gamma})}{\lambda^2}}{\bar{M} \rho_0^n}} \quad (14.15)$$

and

$$\Lambda_0 = \frac{\rho_0^4 \frac{V(0)}{r^4} + \rho_0^2 \frac{\tilde{P}}{4r^2 \lambda^2} + \frac{4\tilde{R}(\tilde{\Gamma})}{\lambda^2}}{\rho_0^{n+2}}, \quad (14.16)$$

where ρ_0 is given by Eq. (14.6). In the case $\tilde{R}(\tilde{\Gamma}) = 0$ one easily gets

$$m_0 = \left(\frac{\lambda^{\frac{n}{2}}}{r} \right) \frac{\sqrt{\tilde{P}(n-2n-2)}}{4\sqrt{|\bar{M}|}} \left(\frac{n\tilde{P}}{(1-n)V(0)} \right)^{-\frac{n}{4}} \quad (14.17)$$

and a range

$$r_0 = \frac{\hbar}{m_0 c}, \quad (14.17a)$$

$$A_0 = (\pm 1)^n \frac{\lambda^{n+2} \tilde{P}}{(1-n)r^{n-2}} \left(\frac{n \tilde{P}}{(n-1)V(0)} \right)^{-\frac{n}{2}}. \quad (14.18)$$

We can calculate $V(0)$ using Eq. (6.18). One gets:

$$V(0) = \frac{\ell_{ab}}{V_2} \int_M \sqrt{|\tilde{g}|} d^{n_1} x (2g^{[\tilde{m}\tilde{n}]} g^{[\tilde{b}\tilde{n}]} \mu^a_{\hat{i}} \mu^b_{\hat{j}} f^{\hat{i}}_{\tilde{m}\tilde{n}} f^{\hat{j}}_{\tilde{a}\tilde{b}} - g^{\tilde{a}\tilde{m}} g^{\tilde{b}\tilde{n}} \mu^b_{\hat{i}} f^{\hat{i}}_{\tilde{m}\tilde{n}} L^a_{\tilde{a}\tilde{b}}(0)), \quad (14.19)$$

where

$$\ell_{dc} g_{\tilde{m}\tilde{b}} g^{\tilde{c}\tilde{m}} L^d_{\tilde{c}\tilde{a}}(0) + \ell_{cd} g_{\tilde{a}\tilde{m}} g^{\tilde{m}\tilde{c}} L^d_{\tilde{b}\tilde{c}}(0) = -2\ell_{cd} g_{\tilde{a}\tilde{m}} g^{\tilde{m}\tilde{c}} \mu^d_{\hat{i}} f^{\hat{i}}_{\tilde{b}\tilde{c}}. \quad (14.20)$$

In the symmetric case one easily finds:

$$V(0) = -\frac{1}{V_2} \int_M \sqrt{|\tilde{g}|} d^{n_1} x g^{\tilde{a}\tilde{m}} g^{\tilde{b}\tilde{n}} f^{\hat{i}}_{\tilde{m}\tilde{n}} f^{\hat{j}}_{\tilde{a}\tilde{b}} h_{ab} \mu^a_{\hat{j}} \mu^b_{\hat{j}}. \quad (14.21)$$

Let us consider the effective gravitational constant G_{eff} close to the minimum of $W(\rho)$. One gets:

$$G_{\text{eff}} = G_0 e^{-(n+2)\Psi} = G_N e^{-(n+2)/\sqrt{|M|}\Psi'} \cong G_N \left(1 - \frac{\alpha}{r} e^{-(r/r_0)} \right), \quad (14.22)$$

where

$$G_N = G_0 \rho_0^{n+2} \quad (14.23)$$

and

$$0 < \alpha = \frac{n+2}{\sqrt{|M|}} \gamma, \quad (14.24)$$

(γ is a positive constant).

According to new observational data [104] concerning distances of type Ia supernovae it seems that we need some kind of “dark energy” which drives the evolution of the Universe. This “dark energy” can be considered as a cosmological constant or more general as cosmological terms in field equations (in the lagrangian). In the case of cosmological model this type of “dark energy”—“vacuum energy” is a cause to accelerate the evolution of the Universe (a scale factor $R(t)$) (see Ref. [105]). The cosmological constant is negligible on the level of the Solar System and on the level of the Galaxy. Moreover it can be important if we consider even nonrelativistic movement of galaxies in a cluster of galaxies (see Ref. [106]). In some papers considered cosmological terms result in changing with time of a cosmological constant (see Ref. [107]). Some of them introduce additional scalar field (or fields) in order to give a field-theoretical description of such an evolution of a cosmological “constant”. Those scalar fields are independent in general of the additional scalar fields in inflationary models.

Thus the inflation field (or fields in multicomponent inflation, which can be the same as some of Higgs' fields from G.U.T.-models) can be different from those fields. In particular considering scalar-tensor theories of gravitation results in so called quintessence models (see Ref. [108]). Moreover in such theories there is a natural field-theoretical background for an inconstant "gravitational constant". In such a way this quintessence field can be used in twofold ways. First as a source of change in space and time of a gravitational constant. Secondly as a source of cosmological terms leading to the model of quintessence and a change in time of a cosmological "constant". In our theory we have a natural occurrence of these phenomena due to the scalar field Ψ (or ρ). Let us consider the lagrangian of our theory paying a special attention to the part involving the scalar field Ψ , i.e.:

$$\begin{aligned}
L = & \overline{R}(\overline{W}) + 8\pi G_N \left(e^{-(n+2)\Psi} \mathcal{L}_{\text{YM}}(\tilde{A}) + \frac{e^{-2\Psi}}{4\pi r^2} \mathcal{L}_{\text{kin}}(\overset{\text{gauge}}{\nabla} \Psi) \right. \\
& \left. - \frac{e^{(n-2)\Psi}}{8\pi r^2} V(\Phi) - \frac{e^{(n-2)\Psi}}{2\pi r^4} \mathcal{L}_{\text{int}}(\Phi, \tilde{A}) \right) \\
& - 8\pi G_N \mathcal{L}_{\text{scal}}(\Psi) + e^{n\Psi} \left(\frac{e^{2\Psi} \alpha_s^2}{l_{\text{pl}}^2} \tilde{R}(\tilde{\Gamma}) + \frac{\tilde{P}}{r^2} \right),
\end{aligned} \tag{14.25}$$

where all the terms are defined by Eqs (6.16–22).

$$\mathcal{L}_{\text{scal}}(\Psi) = \left(\overline{M} \tilde{g}^{(\gamma\nu)} + n^2 g^{[\mu\nu]} g_{\delta\mu} \tilde{g}^{(\delta\gamma)} \right) \Psi_{,\nu} \Psi_{,\gamma}$$

and $\overline{M}$ is defined by Eq. (6.27a). We put $c = \hbar = 1$.

Now we rewrite the lagrangian (14.25) in the following form.

$$\begin{aligned}
L = & \overline{R}(\overline{W}) - 8\pi G_N e^{-(n+2)\Psi} L_{\text{matter}} \\
& - 8\pi G_N \mathcal{L}_{\text{scal}}(\Psi) + e^{n\Psi} \left(\frac{e^{2\Psi}}{l_{\text{pl}}^2} \alpha_s^2 \tilde{R}(\tilde{\Gamma}) + \frac{\tilde{P}}{r^2} \right),
\end{aligned} \tag{14.25a}$$

where in L_{matter} we include all the terms from Eq. (14.25) with Yang-Mills' fields, Higgs' fields, their interactions and coupling to the scalar field Ψ .

The effective gravitational constant is defined by

$$G_{\text{eff}} = G_N e^{-(n+2)\Psi}$$

in such a way that the lagrangian of the Yang-Mills field in L_{matter} is without any factor involving scalar field Ψ . If the scalar field Ψ is constant (e.g. $\Psi = 0$) we can redefine all the fields in such a way that we get ordinary (standard) lagrangians for these fields.

Let us consider the situation after a spontaneous symmetry breaking and simplify to the case of $g_{[\mu\nu]} = 0$. We get

$$L = \tilde{\tilde{R}} - 8\pi G_{\text{eff}} L_{\text{matter}} - 8\pi G_N L_{\text{scal}}(\Psi) + 8\pi G_N U(\Psi) \tag{14.26}$$

where

$$L_{\text{scal}} = \overline{M} g^{\gamma\nu} \Psi_{,\nu} \cdot \Psi_{,\gamma}, \quad \overline{M} > 0 \quad (14.27)$$

$$8\pi G_N U(\Psi) = -\frac{V(\Phi_{\text{crt}}^K)}{r^4} l_{\text{pl}}^2 e^{(n-2)\Psi} + e^{(n+2)\Psi} \left(\frac{\alpha_s^2 \tilde{R}(\tilde{\Gamma})}{l_{\text{pl}}^2} \right) + \frac{\tilde{P}}{r^2} e^{n\Psi}. \quad (14.28)$$

We get the following equations

$$\tilde{\tilde{R}}_{\mu\nu} - \frac{1}{2} \tilde{\tilde{R}} g_{\mu\nu} = 8\pi G_{\text{eff}}^{\text{matter}} T_{\mu\nu}^{\text{matter}} + 8\pi G_N^{\text{scal}} T_{\mu\nu}^{\text{scal}}, \quad (14.29)$$

where $\tilde{\tilde{R}}_{\mu\nu}$ and $\tilde{\tilde{R}}$ are a Ricci tensor and a scalar curvature for a Riemannian geometry generated by $g_{\mu\nu} = g_{(\mu\nu)}$,

$$T^{\text{matter}}{}^{\mu\nu} = (p + \rho) u^\mu u^\nu - p g^{\mu\nu} \quad (14.30)$$

is an energy-momentum tensor for a matter considered as a radiation plus a dust.

$$8\pi G_N^{\text{scal}} T_{\mu\nu}^{\text{scal}} = 8\pi G_N \left(\frac{\overline{M}}{2} g_{\mu\nu} \cdot (g^{\alpha\beta} \Psi_{,\alpha} \cdot \Psi_{,\beta}) - \overline{M} \Psi_{,\mu} \cdot \Psi_{,\nu} \right) + g_{\mu\nu} \bar{\lambda}_{cK} \quad (14.31)$$

where

$$\bar{\lambda}_{cK} = 2e^{(n-2)\Psi} \frac{m_{\tilde{A}}^4}{\alpha_s^4} l_{\text{pl}}^2 V(\Phi_{\text{crt}}^K) - e^{(n+2)\Psi} \frac{\alpha_s^2}{2l_{\text{pl}}^2} \tilde{R}(\tilde{\Gamma}) - e^{n\Psi} \frac{m_{\tilde{A}}^2}{2\alpha_s^2} \tilde{P}, \quad (14.32)$$

where $m_{\tilde{A}}$ is a scale of a mass for massive Yang-Mills' fields (after a spontaneous symmetry breaking for a true vacuum case). It is convenient to write

$$\bar{\lambda}_{cK} = \frac{e^{(n-2)\Psi}}{2} \alpha_K - \frac{e^{n\Psi}}{2} \gamma - \frac{e^{(n+2)\Psi}}{2} \beta \quad (14.33)$$

$$\begin{aligned} \alpha_K &= \alpha_K(\xi, \zeta, m_{\tilde{A}}, \alpha_s) \\ \beta &= \beta(\xi, \alpha_s, m_{\tilde{A}}) \\ \gamma &= \gamma(\zeta, m_{\tilde{A}}, \alpha_s) \end{aligned} \quad (14.33a)$$

$K = 0, 1$ corresponds to true and false vacuum case, i.e.

$$V(\Phi_{\text{crt}}^0) = 0, \quad V(\Phi_{\text{crt}}^1) \neq 0, \quad (14.34)$$

$$\alpha_0 = 0, \quad \alpha_1 \neq 0. \quad (14.34a)$$

For a scalar field Ψ we have the following equation:

$$\begin{aligned} 16\pi G_N \overline{M} g^{\alpha\beta} \left(\tilde{\tilde{\nabla}}_\alpha (\partial_\beta \Psi) \right) + (n-2) e^{(n-2)\Psi} \alpha_K - (n+2) e^{(n+2)\Psi} \beta \\ - n e^{n\Psi} \gamma - (n+2) G_{\text{eff}} T = 0 \end{aligned} \quad (14.35)$$

where $T = \rho - 3p$ is a trace of an energy-momentum tensor for a matter field. For we are interested in cosmological models we take for a metric tensor a Robertson-Walker metric:

$$ds^2 = dt^2 - R^2(t) \left[\frac{dr^2}{1 - kr^2} + r^2 d\theta^2 + r^2 \sin^2 \theta d\varphi^2 \right], \quad k = -1, 0, 1 \quad (14.36)$$

and we suppose that Ψ, ρ, p are functions of t only.

One gets

$$\begin{aligned} \frac{1}{M_{\text{pl}}^2} \ddot{\Psi} &= \frac{3}{M_{\text{pl}}^2} H \dot{\Psi} - \left(\frac{n-2}{\overline{M}} \right) \alpha_K e^{(n-2)\Psi} + \frac{(n+2)}{\overline{M}} \beta e^{(n+2)\Psi} \\ &+ \frac{n}{\overline{M}} e^{n\Psi} \gamma + \frac{(n+2)}{M_{\text{pl}}^2} e^{-(n+2)\Psi} \cdot (\rho - 3p) \end{aligned} \quad (14.37)$$

$$M_{\text{pl}} = \left(\sqrt{8\pi G_N} \right)^{-1} = \frac{m_{\text{pl}}}{\sqrt{8\pi}}.$$

In this case we get standard equations for a cosmological model adapted to our theory

$$\frac{3\ddot{R}}{R} = -8\pi G_{\text{eff}} \left(\frac{1}{2}(\rho + 3p) \right) - 8\pi G_N \left(\frac{1}{2}(\rho_\Psi + 3p_\Psi) \right), \quad (14.38)$$

$$R\ddot{R} + 2\dot{R} + 2k = 8\pi G_{\text{eff}} \left(\frac{1}{2}(\rho - p) \right) R^2 + 8\pi G_N \left(\frac{1}{2}(\rho_\Psi - p_\Psi) \right) R^2. \quad (14.39)$$

Using Eqs (14.38–39) one easily gets

$$H^2 + k = \frac{8\pi G_{\text{eff}}}{3} \rho + \frac{8\pi G_N}{3} \rho_\Psi \quad (14.40)$$

where $H = \frac{\dot{R}}{R}$ is a Hubble constant

$$\begin{aligned} 8\pi G_N \rho_\Psi &= 8\pi G_N \frac{\overline{M}}{2} \dot{\Psi}^2 + \frac{1}{2} \overline{\lambda}_{cK} \\ 8\pi G_N p_\Psi &= 8\pi G_N \frac{\overline{M}}{2} \dot{\Psi}^2 - \frac{1}{2} \overline{\lambda}_{cK}. \end{aligned} \quad (14.41)$$

Let us consider a cosmological model for a “false vacuum” case, i.e. without matter and only with a scalar Ψ and a vacuum energy $V(\Phi_{\text{crt}}^1) \neq 0$. In this case $\rho = p = 0$ and we get

$$2\overline{M}\ddot{\Psi} - \frac{6\overline{M}\dot{R}}{R}\dot{\Psi} - M_{\text{pl}}^2 \frac{d\lambda_{c1}}{d\Psi} = 0 \quad (14.42)$$

$$\dot{R}^2 + k = \frac{1}{3} \left(\frac{1}{2} \frac{\overline{M}}{M_{\text{pl}}^2} \dot{\Psi}^2 + \frac{1}{2} \lambda_{c1} \right) R^2 \quad (14.43)$$

$$\frac{3\ddot{R}}{R} = -\frac{\overline{M}}{M_{\text{pl}}^2}\dot{\Psi}^2 + \frac{1}{2}\lambda_{c1}. \quad (14.44)$$

Let us take

$$\Psi = \Psi_1 = \text{const.} \quad (14.45)$$

Thus we get

$$\frac{d\lambda_{c1}}{d\Psi}(\Psi_1) = 0 \quad (14.46)$$

and

$$(n+2)\beta x_1^4 + n\gamma x_1^2 - (n-2)\alpha_1 = 0 \quad (14.47)$$

$$\dot{R}^2 + k = \frac{1}{6}\lambda_{c1}(x_1)R^2 \quad (14.48)$$

$$\frac{3\ddot{R}}{R} = \frac{1}{2}\lambda_{c1}(x_1) \quad (14.49)$$

$$x_1 = e^{\Psi_1}. \quad (*)$$

One gets from Eq. (14.49)

$$\overline{R}(t) = R_0 e^{H_0 t} \quad (14.50)$$

where

$$H_0 = \sqrt{\frac{\lambda_{c1}(x_1)}{6}} \quad (14.51)$$

is Hubble constant (really constant). From Eq. (14.47) we obtain

$$x_1 = \sqrt{\frac{-n\gamma + \sqrt{n^2\gamma^2 + 4(n^2 - 4)\alpha_1\beta}}{2(n+2)\beta}}. \quad (14.52)$$

For $\alpha_1 > 0$, $\beta > 0$ we get

$$\sqrt{n^2\gamma^2 + 4(n^2 - 4)\alpha_1\beta} > n|\gamma| \quad (14.53)$$

$$\lambda_{c1}(x_1) = \frac{x_1^{n-2}}{2(n+2)^2\beta} \left[n\gamma^2 - \gamma\sqrt{n^2\gamma^2 + 4(n^2 - 4)\alpha_1\beta} + 4\alpha_1\beta(n+2) \right] \quad (14.54)$$

$$H_0 = \frac{x_1^{(n-2)/2}}{2(n+2)\sqrt{3}\beta} \left[n\gamma^2 - \gamma\sqrt{n^2\gamma^2 + 4(n^2 - 4)\alpha_1\beta} + 4\alpha_1\beta(n+2) \right]^{1/2}. \quad (14.55)$$

Thus we get an exponential expansion of the Universe. Using Eq. (14.48) we get also $k = 0$ (i.e. a flatness of a space).

In this way we get de Sitter model of the Universe. This is of course a very special solution to the Eqs (14.42–44) with very special initial conditions

$$\left. \begin{aligned} \overline{R}(0) &= R_0 \\ \frac{d\overline{R}}{dt}(0) &= H_0 R_0 \\ \Psi(0) &= \Psi_1 \\ \frac{d\Psi}{dt}(0) &= 0 \end{aligned} \right\} \quad (14.56)$$

Let us disturb the solution by a small perturbation and examine its stability. Let

$$\begin{aligned} \Psi &= \Psi_1 + \varphi, \quad |\varphi|, |\dot{\varphi}| \ll \Psi_1 \\ R &= \overline{R} + \delta R, \quad |\delta R|, |\delta \dot{R}| \ll \overline{R}. \end{aligned} \quad (14.57)$$

One gets in a linear approximation

$$2\overline{M}\ddot{\varphi} + 6\overline{M}H_0\dot{\varphi} + M_{\text{pl}}^2 \frac{d^2\lambda_{c1}}{d\Psi^2}(\Psi_1)\varphi = 0 \quad (14.58)$$

$$\frac{3\delta\ddot{R}}{\overline{R}} = -\overline{M}\dot{\varphi}^2 \quad (14.59)$$

$$2\overline{R}\delta\dot{R} + k = \frac{1}{6}\overline{M}\dot{\varphi}^2(\overline{R}^2 + \overline{R}\delta R). \quad (14.60)$$

One gets

$$\varphi = \varphi_0 e^{-\frac{3}{2}H_0 t} \sin\left(\frac{\sqrt{p}}{2}t + \delta\right) \quad (14.61)$$

where (we take for simplicity $M = \frac{\overline{M}}{M_{\text{pl}}^2}$)

$$p = -\Delta = \frac{1}{2M} \frac{d^2\lambda_{c1}}{d\Psi^2}(\Psi_1) - 9H_0^2 > 0 \quad (14.62)$$

if

$$\begin{aligned} M < M_0 &= \frac{4}{3} \\ &\times \frac{n(n+2)\gamma\sqrt{n^2\gamma^2 + 4(n^2-4)\alpha_1\beta} - 4(n+2)^2(n-1)\alpha_1\beta - n^2(n+2)\gamma^2}{n\gamma^2 + 4(n+2)\alpha_1\beta - \gamma\sqrt{n^2\gamma^2 + 4(n^2-4)\alpha_1\beta}}. \end{aligned} \quad (14.63)$$

Thus we get an exponential decay of the solution φ , i.e. a damped oscillation around $\Psi = \Psi_1$. It means the solution $\Psi = \Psi_1$ is stable against small perturbations of initial conditions for Ψ , i.e.

$$\Psi(0) = \Psi_1 + \varphi_0 \sin \delta \quad (14.64)$$

$$\frac{d\Psi}{dt}(0) = \varphi_0 \left(-\frac{3}{2}H_0 \sin \delta + \cos \delta\right). \quad (14.65)$$

Making φ_0 and δ sufficiently small we can achieve smallness of perturbations of initial conditions. The exponential decay of the solution to Eq. (14.58) can be satisfied also for some different conditions than Eq. (14.63) if we consider aperiodic case. However, we do not discuss it here.

Thus we get

$$0 \leq \frac{1}{2} \overline{M} \dot{\varphi}^2 \leq a^2 e^{-3H_0 t} \quad (14.66)$$

where a^2 is a constant.

From Eq. (14.59) one gets

$$0 \leq \delta \ddot{R} \leq a^2 e^{-4H_0 t}, \quad (14.67)$$

which means that $\delta \ddot{R} \sim 0$ and $\delta \dot{R} = O(\delta R(0))$. It means $\delta R = \delta R(0) + \delta \dot{R}(0)t$. Moreover from Eq. (14.60) we get that $k = 0$ due to the fact that δR is a linear function of t . In this way we get

$$R(t) \cong \overline{R}(t) + \delta R(0) + \delta \dot{R}(0)t = \left(R_0 + \frac{\delta R(0) + \delta \dot{R}(0)t}{e^{H_0 t}} \right) e^{H_0 t}. \quad (14.68)$$

This means that the small perturbations for initial conditions for R result in a perturbation of R_0 in such a way that R_0 is perturbed by a quickly decaying function. Thus our de Sitter solution is stable under small perturbations and is an attractor for any small perturbed initial data. One can think that this evolution will continue forever. However, we should remember that we are in a “false” vacuum regime and this configuration of Higgs’ fields is unstable. There is a stable configuration—a “true” vacuum case for which $\alpha_K = 0$ ($K = 0$). For $\alpha = 0$ the considered de Sitter evolution cannot be continued. Thus we should take under consideration a second order phase transition in the configuration of Higgs’ fields from metastable state to stable state, from “false” vacuum to “true” vacuum, from Φ_{crt}^1 to Φ_{crt}^0 . In this case the Higgs fields play a rôle of an order parameter. Let us suppose that the scale time of this phase transition is small in comparison to $\frac{1}{H_0}$ and let it take place locally. We suppose locally a conservation of a density of an energy. Thus

$$T_{44}^{\text{scal}} = T_{44}^{\text{scal}} + T_{44}^{\text{matter}}. \quad (14.69)$$

We suppose also that the scalar field Ψ will be close to the new equilibrium (new minimum for cosmological terms) and that the matter will consist of a radiation only:

$$T_{44}^{\text{scal}} = \frac{1}{2} \lambda_{c1}(x_1) \quad (14.70a)$$

$$T_{44}^{\text{scal}} + T_{44}^{\text{matter}} = \frac{1}{2} \lambda_{c0}(x_0) + 8\pi G_N \frac{\rho_r}{x_0^{n+2}}. \quad (14.70b)$$

For $x_0 = e^{\Psi_0}$ is a new equilibrium point we have:

$$\frac{d\lambda_{c0}}{d\Psi}(\Psi_0) = 0 \quad (14.71)$$

$$\lambda_{c0} = -e^{(n+2)\Psi} \beta - e^{n\Psi} \gamma \quad (14.72)$$

$$\frac{d\lambda_0}{d\Psi} = -e^{n\Psi} ((n+2)\beta e^{2\Psi} + n\gamma). \quad (14.73)$$

One gets

$$e^{\Psi_0} = x_0 = \sqrt{\frac{n|\gamma|}{(n+2)\beta}} \quad (14.74)$$

and supposing

$$\gamma < 0 \quad (14.75)$$

thus

$$\lambda_{c0}(\Psi_0) = \frac{2|\gamma|^{\frac{n}{2}+1} n^{\frac{n}{2}}}{\beta^{\frac{n}{2}} (n+2)^{\frac{n}{2}+1}}. \quad (14.76)$$

From Eqs (14.70ab) one gets

$$\begin{aligned} \rho_r = \frac{1}{16\pi G_N} \left\{ \frac{x_1^{n-2}}{(n+2)^2 \beta} \left[n\gamma^2 - \gamma \sqrt{n^2 \gamma^2 + 4(n^2 - 4)\alpha_1 \beta} + 2\alpha_1 \beta (n+4) \right] \right. \\ \left. - \frac{2|\gamma|^{\frac{n}{2}+1}}{\beta^{\frac{n}{2}} (n+2)^{\frac{n}{2}+1}} \right\} \frac{n^{\frac{n+2}{2}} \gamma^{\frac{n+2}{2}}}{(n+2)^{\frac{n+2}{2}} \beta^{\frac{n+2}{2}}}. \end{aligned} \quad (14.77)$$

This gives us matching condition for a second order phase transition and simultaneously it is an initial condition for a new epoch of an evolution of the Universe plus a condition $\dot{\Psi}(t_r) = 0$, $R(t_r) = \overline{R}(t_r)$, where t_r is a time for a phase transition to occur. Notice we have simply $\Psi_0 \neq \Psi_1$. Thus we have to do with a discontinuity for a field Ψ . Let us consider Hubble's constants for both phases of the Universe

$$\begin{aligned} H_0^2 &= \frac{\lambda_{c1}(x_1)}{6} \\ H_1^2 &= \frac{\lambda_{c0}(x_0)}{6} = \frac{|\gamma|^{\frac{n}{2}+1} n^{\frac{n}{2}}}{3\beta^{\frac{n}{2}} (n+2)^{\frac{n}{2}+1}} \\ H_0^2 &\neq H_1^2 \end{aligned} \quad (14.79)$$

Summing up we get

$$\begin{aligned} R(t_r) &= \overline{R}(t_r) = R_0 e^{+H_0 t_r} \\ \dot{\Psi}_1(t_r) &= \dot{\Psi}_0(t_r) = 0 \\ \Psi_1 &\neq \Psi_0 \\ H_0^2 &\neq H_1^2 \end{aligned} \quad (14.80)$$

Thus we see that the second order phase transition in the configuration of Higgs' fields results in the first order phase for an evolution of the Universe. We get discontinuity for Hubble constants and values of scalar field before and after phase transition.

Let us calculate deceleration parameters before and after phase transition

$$q = -\frac{R\ddot{R}}{\dot{R}^2} \quad (14.81)$$

$$q = -1 \quad (14.82)$$

before phase transition and

$$q = -1 \quad (14.83)$$

after phase transition.

Let us come back to the Eqs (14.38), (14.40), (14.37). Let us rewrite Eq. (14.37) in the following form

$$\overline{M}\ddot{\Psi} + \frac{6\overline{M}\dot{R}}{R}\dot{\Psi} + M_{\text{pl}}^2 \frac{d\lambda_{c0}}{d\Psi} = 0. \quad (14.84)$$

(Remember we now have to do with a radiation for which the trace of an energy-momentum tensor is zero.)

Let us suppose that $\dot{\Psi} \approx 0$. Thus Eq. (14.84) simplifies

$$\frac{1}{M_{\text{pl}}^2} M\ddot{\Psi} + \frac{d\lambda_{c0}}{d\Psi} = 0 \quad (14.85)$$

and we get the first integral of motion

$$\frac{\overline{M}}{2M_{\text{pl}}^2} \dot{\Psi}^2 + \lambda_{c0} = \frac{1}{2} \bar{\delta} = \text{const.} \quad (14.86)$$

$$\overline{M}\dot{\Psi}^2 = M_{\text{pl}}^2 (\bar{\delta} - 2\lambda_{c0}). \quad (14.86a)$$

One gets from Eqs (14.38) and (14.40)

$$\frac{3\ddot{R}}{R} = -\frac{1}{M_{\text{pl}}^2} \cdot \frac{\rho_r}{e^{(n+2)\Psi}} - \frac{1}{2} \left(\frac{2\overline{M}}{M_{\text{pl}}^2} \dot{\Psi}^2 - \lambda_{c0} \right) \quad (14.87)$$

$$\dot{R}^2 + k = \frac{1}{3M_{\text{pl}}^2} \cdot \frac{\rho_r}{e^{(n+2)\Psi}} R^2 + \frac{1}{3} \left(\frac{\overline{M}}{2M_{\text{pl}}^2} \dot{\Psi}^2 + \frac{1}{2} \lambda_{c0} \right) R^2. \quad (14.88)$$

Using Eq (14.86a) we get:

$$\frac{3\ddot{R}}{R} = -\frac{1}{M_{\text{pl}}^2} \cdot \frac{\rho_r}{e^{(n+2)\Psi}} - \bar{\delta} + \frac{5}{2} \lambda_{c0} \quad (14.89)$$

$$\dot{R}^2 + k = \frac{1}{3M_{\text{pl}}^2} \cdot \frac{\rho_r}{e^{(n+2)\Psi}} R^2 + \frac{1}{6} (\bar{\delta} - \lambda_{c0}) R^2. \quad (14.90)$$

One can derive from Eqs (14.89–90)

$$\frac{d}{dt}(R\dot{R}) = -\frac{1}{6}\bar{\delta}R^2 + \frac{2}{3}\lambda_{c0}R^2 - k \quad (14.91)$$

$$\lambda_{c0} = -e^{n\Psi}(\beta e^{2\Psi} + \gamma). \quad (14.92)$$

Let us take $k = 0$ and $\delta = 0$ and let us change independent and dependent variables in (14.91) using (14.92) and (14.86a). One gets

$$2y^2(y^2 - 1)\frac{d^2f}{dy^2} + y((n+4)y^2 - (n+1)y - 2)\frac{df}{dy} = \frac{4\bar{M}}{3}(y^2 - 1)f(y) \quad (14.93)$$

where

$$y = \sqrt{\frac{\beta}{|\gamma|}}e^\Psi \quad (14.94)$$

and

$$f = R^2. \quad (14.95)$$

It is easy to see that $y^2 > 1$ for

$$\frac{d\Psi}{dt} = \pm \frac{M_{\text{pl}}}{\sqrt{\bar{M}}} \cdot \frac{|\gamma|^{\frac{n+2}{4}}}{\beta^{\frac{n}{4}}} \sqrt{y^n(y^2 - 1)}. \quad (14.96)$$

Moreover according to our assumptions $\dot{\Psi} \approx 0$ and this can be achieved only if $0 < y - 1 < \varepsilon$, where ε is sufficiently small.

Thus for further investigations we take $z = y - 1$, $y = z + 1$. One gets

$$2z(z+2)(z+1)^2\frac{d^2f}{dz^2} + (z+1)((n+4)z^2 + (n+7)z - (n+3))\frac{df}{dz} - \frac{4\bar{M}}{3}z(z+2)f(z) = 0. \quad (14.97)$$

Let us consider Eqs (14.89–90) supposing $k = 0$ and $\bar{\delta} = 0$.

After eliminating $\ddot{R}$ (via differentiation of Eq. (14.90) with respect to time t) and changing independent and dependent variables one gets:

$$\frac{d}{dy} \left[\left(8\pi G_N \frac{\tilde{\rho}_r}{y^{n+2}} + \frac{1}{2}y^n(y^2 - 1) \right) f^2 \right] = -\frac{d}{dy}(f^2)y^n(y^2 - 1) \quad (14.98)$$

where

$$\bar{\rho}_r = \rho_r \frac{|\gamma|^n}{\beta^{n+1}}. \quad (14.99)$$

It is convenient for further investigations to consider a parameter

$$\bar{r} = \frac{4\bar{M}}{3}. \quad (14.100)$$

In such a way we get

$$2y^2(y^2 - 1)\frac{d^2 f}{dy^2} + y((n+4)y^2 - (n+1)y - 2)\frac{df}{dy} - \bar{r}(y^2 - 1)f(y) \quad (14.101)$$

and

$$2z(z+2)(z+1)^2\frac{d^2 f}{dz^2} + (z+1)((n+4)z^2 + (n+7)z - (n+3))\frac{df}{dz} - \bar{r}z(z+2)f(z) = 0. \quad (14.102)$$

One can transform Eq. (14.98) into

$$\frac{d}{dy} \left[\left(8\pi G_N \frac{\tilde{\rho}_r}{y^{n+2}} + \frac{3}{2} y^n (y^2 - 1) \right) f^2 \right] = y^{n-1} [(n+2)y^2 - n] f^2. \quad (14.103)$$

Changing the independent variable from y to $z = y - 1$ one gets:

$$\begin{aligned} \frac{d}{dz} \left[\left(8\pi G_N \frac{\tilde{\rho}_r}{(z+1)^{n+2}} + \frac{3}{2} (z+1)^n z(z+2) \right) f^2 \right] \\ = (z+1)^{n-1} [(n+2)(z+1)^2 - n] f^2. \end{aligned} \quad (14.104)$$

Let us come back to Eq. (14.96) in order to find time-dependence of Ψ . One gets

$$\int \frac{d\Psi}{\sqrt{\beta e^{(n+2)\Psi} - |\gamma| e^{n\Psi}}} = \pm \frac{t - t_1}{\sqrt{M}} M_{\text{pl}} \quad (14.105)$$

or

$$\int \frac{dx}{x \sqrt{\beta x^{n+2} - |\gamma| x^n}} = \pm \frac{t - t_1}{\sqrt{M}} M_{\text{pl}}. \quad (14.105a)$$

If $n = 2l$, where l is a natural number, one gets for $\beta > 0$

$$\begin{aligned} \left(\frac{|\gamma|}{\beta} \right)^{-\frac{1}{2}} \left\{ \sum_{k=1}^l \sum_{p=0}^k \frac{\binom{k}{p} (-2)^{k-p}}{2p} \left(\sqrt{\frac{\sqrt{\beta}x - \sqrt{|\gamma|}}{\sqrt{\beta}x + \sqrt{|\gamma|}}} \right)^{2p+1} \right. \\ \left. + \frac{1}{2\sqrt{3}} \ln \left(\frac{\sqrt{\sqrt{\beta}x - |\gamma|} - \sqrt{3(\sqrt{\beta}x + |\gamma|)}}{\sqrt{\sqrt{\beta}x - |\gamma|} + \sqrt{3(\sqrt{\beta}x + |\gamma|)}} \right) \right\} = \pm \frac{(t - t_1)}{\sqrt{M}} M_{\text{pl}}. \end{aligned} \quad (14.106)$$

In this case it can be expressed in terms of elementary functions. For a general case of n , β and γ one gets

$$\frac{2x^{-\frac{n}{2}} {}_1F_2\left(\frac{1}{2}, -\frac{n}{4}, \left(1 - \frac{n}{4}\right); \frac{x^2\beta}{\gamma}\right)}{\sqrt{\gamma}n} = \pm \frac{(t - t_1)}{\sqrt{M}} M_{\text{pl}} \quad (14.107)$$

where ${}_1F_2(a, b, c; z)$ is a hypergeometric function. For a small z (around zero) one finds the following solution to Eq. (14.102)

$$f(z) = C_1 z^{\frac{(n+7)}{4}} \left(1 - \frac{(n+7)(5n+23)}{8(n+1)} z \right) + C_2 \left(1 + \frac{\bar{r}}{(n-1)} z^2 \right), \quad (14.108)$$

$C_1, C_2 = \text{const.}$ Thus

$$R(y) = \left[C_1 (y-1)^{\frac{(n+7)}{4}} \left(1 - \frac{(n+7)(5n+23)}{8(n+1)} (y-1) \right) + C_2 \left(1 + \frac{\bar{r}}{(n-1)} (y-1)^2 \right) \right]^{\frac{1}{2}} \quad (14.109)$$

where $y > 1$ (but only a little).

Let us make some simplifications of the formulae taking under consideration that $y - 1$ is very small.

One gets

$$f(y) = C \left(1 + \frac{\bar{r}}{(n-1)} (y-1)^2 \right), \quad C > 0 \quad (14.110)$$

$$R(y) = R_1 \left(1 + \frac{\bar{r}}{(n-1)} (y-1)^2 \right)^{\frac{1}{2}}. \quad (14.111)$$

After some calculations one gets

$$8\pi G_N \tilde{\rho}_r = y^{2(n+1)} \left(-\frac{2(n+5)}{(n+4)(n-1)} y^4 + \frac{10\bar{r}(n+2)}{(n-1)(n+4)} y^3 + \frac{(8\bar{r} - (n-1)(n+4))}{2(n-1)(n+4)} y^2 - \frac{2\bar{r}(n+3)}{(n^2-1)} y + \frac{(\bar{r} - n + 1)}{(n-1)} \right) \quad (14.112)$$

(taking into account that $(y-1)$ is very small). Let us make some simplifications in the formula (14.96) for y close to 1. We find:

$$\int \frac{dy}{\sqrt{y-1}} = \pm \frac{|\gamma|^{\frac{n+4}{4}} \sqrt{2}}{\beta^{\frac{n}{4}} \sqrt{M}} (t - t_1) M_{\text{pl}} \quad (14.113)$$

and finally

$$y = \frac{|\gamma|^{\frac{n+4}{2}}}{2\bar{M}\beta^{\frac{n}{2}}} (t - t_1)^2 + 1 \quad (14.114)$$

such that $y = 1$ for $t = t_1$.

Let us pass to the minimum value for $\tilde{\rho}_r$ with respect to y close to 1. Thus we are looking for a minimum of a function

$$\tilde{\rho} = \frac{1}{8\pi G_N} y^{2(n+1)} \left[-\frac{2(n+5)}{(n+4)(n-1)} y^4 + \frac{10\bar{r}(n+2)}{(n-1)(n+4)} y^3 + \frac{(8\bar{r} - (n-1)(n+4))}{2(n-1)(n+4)} y^2 - \frac{2\bar{r}(n+3)}{(n^2-1)} y + \frac{(\bar{r} - n + 1)}{(n-1)} \right] = \frac{1}{8\pi G_N} y^{2(n+1)} W_4(y) \quad (14.115)$$

for such a y that is close to 1.

Let us write $y = 1 + \eta$, where η is very small and develop $W_4(1 + \eta)$ up to the second order in η . One finds

$$W_4(1 + \eta) \simeq V_2(\eta) \quad (14.116)$$

and

$$V_2(\eta) = a\eta^2 + b\eta + c \quad (14.117)$$

where

$$a = \frac{(-n^2 + 3n(20\bar{r} - 9) + 4(32\bar{r} - 31))}{2(n + 4)(n - 1)} \quad (14.118)$$

$$b = \frac{(-n^3 - 4n^2(7\bar{r} - 3) + 5n(18\bar{r} - 11) + 22(3\bar{r} - 2))}{(n + 4)(n^2 - 1)} \quad (14.119)$$

$$c = \frac{(-3n^3 + 2n^2(9\bar{r} - 8) + n(50\bar{r} - 29) + 4(7\bar{r} - 4))}{2(n + 4)(n^2 - 1)} \quad (14.120)$$

Let us find minimum of $V_2(\eta)$ for $\eta > 0$ such that $V_2(\eta) \geq 0$. For $a < 0$, $b < 0$, $c > 0$ we get

$$V_2(\eta_{\min}) = 0 \quad (14.121)$$

and

$$\eta_{\min} = \frac{b + \sqrt{b^2 + 4|a|c}}{2|a|}. \quad (14.122)$$

In this case

$$(1 + \eta_{\min})^{2(n+1)} W_4(1 + \eta_{\min}) > 0 \quad (14.123)$$

and will be close to the real minimum of the function (14.115). Simultaneously

$$\eta_{\min} < 1.$$

For

$$\tilde{\rho}_r = \frac{|\gamma|^n}{\beta^{n+1}} \sigma T^4 \quad (14.124)$$

(where σ is a Stefan-Boltzmann constant) we get $T_{\min}$ and we call it a T_d (a decoupling temperature-decoupling of matter and radiation). Thus $y_d = 1 + \eta_{\min}$. y_d is reached at a time t_d (a decoupling time)

$$t_d = t_1 + \frac{\eta_{\min} \overline{M} \beta^{\frac{n}{2}}}{2|\gamma|^{\frac{n+4}{2}}} M_{\text{pl}}. \quad (14.125)$$

From that time (t_d) the evolution of the Universe will be driven by a matter (with good approximation a dust matter) and the scalar field Ψ . The radius of the Universe

at $t = t_d$ is equal to

$$R(t_d) = R_1 \left(1 + \frac{\bar{r}}{(n-1)} \eta_{\min}^2 \right)^{\frac{1}{2}}. \quad (14.126)$$

The full field equations (generalized Friedmann equations) are as follows:

$$\left(\frac{\dot{R}}{R} \right)^2 = \frac{8\pi G_{\text{eff}}}{3} \rho_m + \frac{1}{3} \left(\frac{1}{2} \frac{\bar{M}}{M_{\text{pl}}^2} \dot{\Psi}^2 + \frac{1}{2} \lambda_{c0} \right) \quad (14.127)$$

$$\frac{3\ddot{R}}{R} = \frac{8\pi G_{\text{eff}}}{3} \rho_m - \frac{1}{2} \left(2 \frac{\bar{M}}{M_{\text{pl}}^2} \dot{\Psi}^2 - \lambda_{c0} \right) \quad (14.128)$$

$$-\frac{2\bar{M}}{M_{\text{pl}}^2} \ddot{\Psi} - \frac{6\bar{M}}{M_{\text{pl}}^2} \cdot \frac{\dot{R}}{R} \dot{\Psi} - \frac{d\lambda_{c0}}{d\Psi} + (n+2)8\pi\rho_m G_{\text{eff}} = 0. \quad (14.129)$$

Before we pass to those equations we answer the question what is a mass of the scalar field Ψ during the de Sitter phases and the radiation era. One gets:

$$-m_1^2 = \frac{M_{\text{pl}}^2}{2\bar{M}} \frac{d^2 \lambda_{c1}}{d\Psi^2}(\Psi_1) \quad (14.130)$$

$$-m_0^2 = \frac{M_{\text{pl}}^2}{2\bar{M}} \frac{d^2 \lambda_{c0}}{d\Psi^2}(\Psi_0). \quad (14.131)$$

The second question we answer is an evolution of the Universe during a period from t_r to t_1 , i.e. when y is changing from $y = \sqrt{\frac{n}{n+2}}$ to $y = 1$.

This will be simply the de Sitter evolution

$$R(t) = R e^{H_1 t} \quad (14.132)$$

where

$$H_1 = \sqrt{\frac{\lambda_{c0}(x_0)}{6}}. \quad (14.133)$$

This will be unstable evolution and will end at $t = t_1$, i.e. for $\dot{\Psi} \simeq 0$ and value of the field Ψ will be changed to the value corresponding to $y = 1$, i.e.

$$\Psi(t_1) = \frac{1}{2} \ln \frac{\beta}{|\gamma|}. \quad (14.134)$$

Thus we have to do with two de Sitter phases of the evolution with two different Hubble constants H_1 and H_0 . In the third phase (a radiation era) the radius of the Universe is given by the formula

$$R(t) = R_0 e^{H_0 t_r + H_1 (t_1 - t_r)} \cdot \left(1 + \frac{|\gamma|^{n+4}}{3\bar{M}(n-1)\beta^n} (t - t_1)^4 \right)^{\frac{1}{2}} \quad (14.135)$$

and such an evolution ends for $t = t_d$,

$$R(t_d) = R_0 \exp(H_0 t_r + H_1(t_1 - t_r)) \cdot \left(1 + \frac{\bar{r}}{(n-1)} \eta_{\min}^2\right)^{\frac{1}{2}}. \quad (14.136)$$

In both de Sitter phases the equation of state for the matter described by the scalar field Ψ is the same:

$$\rho_\Psi = -p_\Psi. \quad (14.137)$$

In the radiation era we get

$$p_\Psi = \frac{4}{3} \rho_\Psi. \quad (14.138)$$

Now we go to the matter dominated Universe and we change the notation introducing a scalar field Q such that

$$Q = \sqrt{M} \frac{\Psi}{M_{\text{pl}}}. \quad (14.139)$$

In this case we have

$$\rho_Q = \frac{1}{2} \dot{Q}^2 + U(Q) \quad (14.140a)$$

$$p_Q = \frac{1}{2} \dot{Q}^2 - U(Q) \quad (14.140b)$$

where

$$U(Q) = \frac{1}{2} \lambda_{c0} \left(\frac{Q M_{\text{pl}}}{\sqrt{M}} \right) = -\frac{\beta}{2} e^{aQ} + \frac{1}{2} |\gamma| e^{bQ} \quad (14.141)$$

$$a = \frac{(n+2)}{\sqrt{M}} M_{\text{pl}}, \quad b = \frac{n}{\sqrt{M}} M_{\text{pl}}. \quad (14.142)$$

Let us write Eqs (14.127–129) in terms of Q :

$$\ddot{Q} + 3H\dot{Q} + U'(Q) - 8\pi(n+2)G_{\text{eff}}\rho_m = 0 \quad (14.143)$$

where

$$G_{\text{eff}} = G_N e^{-\frac{(n+2)}{\sqrt{M}} M_{\text{pl}} Q} \quad (14.144)$$

$$H^2 = \frac{1}{3} \left(8\pi G_{\text{eff}} \rho_m + \frac{1}{2} \dot{Q}^2 + U \right) \quad (14.145)$$

$$\frac{3\ddot{R}}{R} = \frac{1}{2} \left(8\pi G_{\text{eff}} \rho_m + 2(\dot{Q}^2 - U) \right). \quad (14.146)$$

Let us define an equation of state

$$W_Q = \frac{p_Q}{\rho_Q} = \frac{\frac{1}{2} \dot{Q}^2 - U}{\frac{1}{2} \dot{Q}^2 + U} \quad (14.147)$$

and a variable

$$x_Q = \frac{1 + W_Q}{1 - W_Q} = \frac{\frac{1}{2}\dot{Q}^2}{U}, \quad (14.148)$$

which is a ratio of a kinetic energy to a potential energy density for Q . It is interesting to combine (14.143) and (14.145) using (14.147) and (14.148).

Let us define

$$\Omega_Q = \frac{\frac{1}{2}\dot{Q}^2 + U}{3H^2} \quad (14.149)$$

$$\Omega_m = \frac{8\pi G_{\text{eff}}\rho_m}{3H^2}. \quad (14.150)$$

One gets

$$\Omega_m + \Omega_Q = 1 \quad (14.151)$$

or

$$8\pi G_{\text{eff}}\rho_m = 3(1 - \Omega_Q)H^2. \quad (14.152)$$

We call ρ_m an energy density of a background and for such a matter the equation of state is

$$W_m = \frac{p_m}{\rho_m} = 0. \quad (14.153)$$

It is natural to define

$$\tilde{\rho}_m = 8\pi G_{\text{eff}}\rho_m \quad (14.154)$$

as an effective energy density of a background with the same equation of a state as (14.153).

Under an assumption of slow roll for Q , i.e.

$$\frac{1}{2}\dot{Q}^2 \ll U(Q) \quad (14.155)$$

one gets

$$W_Q \simeq -1, \quad (14.156)$$

i.e. an equation of state of a cosmological constant evolving in time. In this case $x_Q \simeq 0$. This can give in principle an account for an acceleration of an evolution of the Universe if we suppose those conditions for our contemporary epoch.

Let us come back to the inflationary era in our model. For our Higgs' field is multicomponent, i.e. it is a multiplet of Higgs' fields, we have to do with so called multicomponent inflation (see Ref. [109]). In this case we can define slow-roll parameter for our model in equations for Higgs' fields.

One gets from Eq. (9.8)

$$\frac{d}{dt} (L_b^{d0})_{av} - 3H (L_b^{d0})_{av} = -\frac{e^{n\Psi_1}}{2r^2} l^{db} \left\{ \frac{\delta V'}{\delta \Phi_{\tilde{n}}^b} g_{b\tilde{n}} \right\}_{av} \quad (14.157)$$

where $L_{0\tilde{a}}^d = L_{t\tilde{a}}^d$ (a time component of $L_{\mu\tilde{a}}^d$) is defined by

$$l_{dc}L_{0\tilde{a}}^d + l_{cd}g_{\tilde{a}\tilde{m}}g^{\tilde{m}\tilde{c}}L_{0\tilde{c}}^d = 2l_{cd}g_{\tilde{a}\tilde{m}}g^{\tilde{m}\tilde{c}}\frac{d}{dt}(\Phi_{\tilde{c}}^d), \quad (14.158)$$

V' , $\frac{\delta V'}{\delta\Phi_{\tilde{n}}^b}$ and $\left\{\frac{\delta V'}{\delta\Phi_{\tilde{n}}^b}\right\}_{av}$ are defined by Eqs (8.6–12) and Eq. (6.24).

$$3H^2 = 8\pi G_N \left(\frac{e^{(n-2)\Psi_1}}{r^4} V'(\Phi) - \frac{2e^{-2\Psi_1}}{r^2} \mathcal{L}_{\text{kin}}(\dot{\Phi}) + \frac{1}{2} \lambda_{c1}(\Psi_1) \right) \quad (14.159)$$

$$\dot{H} = 8\pi G_N \left(-\frac{2e^{-2\Psi_1}}{r^2} \mathcal{L}_{\text{kin}}(\dot{\Phi}) \right) \quad (14.160)$$

where H is a Hubble constant and

$$\mathcal{L}_{\text{kin}}(\dot{\Phi}) = l_{ab} \left\{ g^{\tilde{b}\tilde{n}} L_{0\tilde{b}}^a \frac{d}{dt} \Phi_{\tilde{n}}^b \right\}_{av}. \quad (14.161)$$

Ψ_1 is a constant value for a field Ψ . The slow-roll parameters are defined by

$$\varepsilon = \frac{l_{ab} \left\{ g^{\tilde{b}\tilde{n}} L_{0\tilde{b}}^a \frac{d}{dt} \Phi_{\tilde{n}}^b \right\}_{av}}{H^2} \quad (14.162)$$

$$\delta = \frac{l_{ab} \left\{ g^{\tilde{b}\tilde{n}} \frac{d}{dt} \left(L_{0\tilde{b}}^a \right) \frac{d}{dt} (\Phi_{\tilde{n}}^b) \right\}_{av}}{H^2 \left(l_{ab} \left\{ g^{\tilde{b}\tilde{n}} L_{0\tilde{b}}^a \frac{d}{dt} (\Phi_{\tilde{n}}^b) \right\}_{av} \right)} \quad (14.163)$$

with the assumptions of the slow roll

$$\varepsilon = O(\eta), \quad \delta = O(\eta) \quad (14.164)$$

for some small parameter η .

Under slow-roll conditions we have

$$3H (L_{0\tilde{b}}^d)_{av} + \frac{e^{n\Psi_1}}{2r^2} l^{db} \left(\frac{\delta V'}{\delta\Phi_{\tilde{n}}^b} g_{\tilde{b}\tilde{n}} \right)_{av} \cong 0 \quad (14.165)$$

$$H^2 \cong \frac{8\pi}{3M_{\text{pl}}^2} \left\{ \frac{e^{(n-2)\Psi_1}}{r^4} V'(\Phi) \right\} + \frac{1}{2} \lambda_{c1}(\Psi_1) \quad (14.166)$$

and with a standard extra assumption

$$\frac{\dot{\delta}}{H} = O(\eta^2). \quad (14.167)$$

We remind that the scalar field Ψ is the more important agent to drive the inflation. However, the full amount of time the inflation takes place depends on an evolution of Higgs' fields (under slow-roll approximation). The evolution must be slow and starting for $\Phi = \Phi_{\text{crt}}^1$. It ends at $\Phi = \Phi_{\text{crt}}^0$.

Thus we have according to Eq. (14.50)

$$\overline{N} = \ln \frac{R(t_{\text{end}})}{R(t_{\text{initial}})} = H_0(t_{\text{end}} - t_{\text{initial}})$$

where

$$\Phi(t_{\text{end}}) = \Phi_{\text{crt}}^0 \quad (14.169)$$

$$\Phi(t_{\text{initial}}) = \Phi_{\text{crt}}^1 \quad (14.170)$$

$$t_{\text{end}} = t_r$$

(we omit indices for Φ).

Thus we should solve Eq. (14.165) for initial condition (14.169) and with an assumption $H = H_0$. We have of course a short period of an inflation with $H = H_1$ from $t_{\text{end}} = t_r$ to $t = t_1$, i.e.

$$\overline{N}_{\text{tot}} = \overline{N} + \overline{N}_1 = H_0(t_r - t_{\text{initial}}) + H_1(t_1 - t_r). \quad (14.171)$$

Let us remind to the reader that $\overline{N}$ is called an amount of inflation. Let us consider inflationary fluctuations of Higgs' fields in our theory. We are ignoring gravitational backreaction for the cosmological evolution is driven by the scalar field Ψ . We write the Higgs field $\Phi_{\tilde{m}}^a$ as a sum

$$\Phi_{\tilde{m}}^a(\vec{r}, t) = \Phi_{\tilde{m}}^a(t) + \delta\Phi_{\tilde{m}}^a(\vec{r}, t) \quad (14.172)$$

where $\Phi_{\tilde{m}}^a(t)$ is a solution of Eq. (14.165) with boundary condition (14.169–170) and $\delta\Phi_{\tilde{m}}^a(\vec{r}, t)$ is a small fluctuation which will be written in terms of Fourier modes

$$\delta\Phi_{\tilde{m}}^a(\vec{r}, t) = \sum_{\vec{K}} \delta\Phi_{\vec{K}\tilde{m}}^a(t) e^{i\vec{K}\vec{r}}. \quad (14.173)$$

The full field equation for Higgs' field in the de Sitter background linearized for $\delta\Phi_{\vec{K}\tilde{m}}^a(t)$ can be written

$$\begin{aligned} \frac{d^2}{dt^2} \left(M_{\vec{K}\tilde{b}}^d \right)_{av} + 3H_0 \frac{d}{dt} \left(M_{\vec{K}\tilde{b}}^d \right)_{av} + \frac{1}{R^2} \vec{K}^2 \left(M_{\vec{K}}^d \right)_{av} \\ = - \frac{e^{n\Psi_1}}{2r^2} l^{db} \left\{ \frac{\delta^2 V' \left(\Phi_{\tilde{b}}^a(t) \right)}{\delta\Phi_{\tilde{n}}^b \delta\Phi_{\tilde{m}}^c} g_{\tilde{b}\tilde{n}} \right\} \delta\Phi_{\vec{K}\tilde{m}}^c(t) \end{aligned} \quad (14.174)$$

where

$$l_{dc} M_{\vec{K}\tilde{b}}^d + l_{cd} g_{\tilde{a}\tilde{m}} g^{\tilde{m}\tilde{c}} M_{\vec{K}\tilde{c}}^d = 2l_{cd} g_{\tilde{a}\tilde{m}} g^{\tilde{m}\tilde{c}} \delta\Phi_{\vec{K}\tilde{c}}^d. \quad (14.175)$$

Let us change dependent and independent variables.

$$d\tau = \frac{dt}{\bar{R}(t)}, \quad \bar{R}(t) = R_0 e^{H_0 t} \quad (14.176)$$

$$R(\tau) = -\frac{1}{H_0 \tau} \quad (14.177)$$

$$-\infty < \tau < 0 \quad (\text{a conformal time}) \quad (14.178)$$

and

$$\delta\Phi_{\vec{K}\vec{m}}^a = \frac{\chi_{\vec{K}\vec{m}}^a}{R}. \quad (14.179)$$

Simultaneously we neglect the term with the second derivative of the potential. Eventually we get

$$\frac{d^2}{d\tau^2} \left(\widetilde{M}_{\vec{K}\vec{b}}^d \right)_{av} - \frac{2}{\tau^2} \frac{d}{d\tau} \left(\widetilde{M}_{\vec{K}\vec{b}}^d \right)_{av} + \vec{K}^2 \left(\widetilde{M}_{\vec{K}\vec{b}}^d \right)_{av} = 0 \quad (14.180)$$

where

$$l_{dc} \widetilde{M}_{\vec{K}\vec{b}}^d + l_{cd} g_{\vec{a}\vec{m}} g^{\vec{m}\vec{c}} \widetilde{M}_{\vec{K}\vec{c}}^d = 2l_{cd} g_{\vec{a}\vec{m}} g^{\vec{m}\vec{c}} \chi_{\vec{K}\vec{c}}^d. \quad (14.181)$$

Let us notice that $|\vec{K}\tau|$ is the ratio of the proper wavenumber $|K|R^{-1}$ to the Hubble radius $\frac{1}{H_0}$. At early times $|K\tau| \gg 1$, the wavelength is small in comparison to Hubble radius and the mode oscillates as in Minkowski space. However, if τ goes to zero, $|K\tau|$ goes to zero too. The wavelength of the mode is stretched far beyond the Hubble radius. It means the mode freezes. Thus the analyzes of the modes are similar to those in classical symmetric inflationary perturbation theory neglecting the fact that we have to do with multicomponent inflation and that the structure of representation space of Higgs' fields should be taken into account.

However, the form of Eq. (14.180) is exactly the same as in one component symmetric theory if we take $\widetilde{M}_{\vec{K}\vec{a}}^d$ a field to be considered. The relation (14.181) between $\widetilde{M}_{\vec{K}\vec{a}}^d$ and $\chi_{\vec{K}\vec{a}}^d$ is linear and under some simple conditions unambiguous.

It is possible to find an exact solution to Eq. (14.180) and Eq. (14.181). One gets

$$\begin{aligned} \chi_{\vec{K}\vec{m}}^a &= C_{1\vec{K}\vec{m}}^a \left(\frac{\tau|\vec{K}| \cos(\tau|\vec{K}|) - \sin(\tau|\vec{K}|)}{\tau} \right) \\ &+ C_{2\vec{K}\vec{m}}^a \left(\frac{\tau|\vec{K}| \sin(\tau|\vec{K}|) + \cos(\tau|\vec{K}|)}{\tau} \right) \end{aligned} \quad (*)$$

and

$$M_{\vec{K}\tilde{b}}^d = N_{1\vec{K}\tilde{b}}^d \left(\frac{\tau|\vec{K}|\cos(\tau|\vec{K}|) - \sin(\tau|\vec{K}|)}{\tau} \right) + N_{2\vec{K}\tilde{b}}^d \left(\frac{\tau|\vec{K}|\sin(\tau|\vec{K}|) + \cos(\tau|\vec{K}|)}{\tau} \right) \quad (**)$$

where

$$l_{dc}N_{i\vec{K}\tilde{b}}^d + l_{cd}g_{\tilde{a}\tilde{m}}g^{\tilde{m}\tilde{c}}N_{i\vec{K}\tilde{c}}^d = 2l_{cd}g_{\tilde{a}\tilde{m}}g^{\tilde{m}\tilde{c}}C_{i\vec{K}\tilde{c}}^d, \quad i = 1, 2. \quad (***)$$

In this way one obtains

$$\delta\Phi_{\vec{K}\tilde{m}}^a = \frac{1}{R_0}e^{-H_0t}\chi_{\vec{K}\tilde{m}}^a \left(-\frac{e^{-H_0t}}{R_0H_0} \right) \quad (****)$$

and using (*) one easily gets

$$\begin{aligned} \delta\Phi_{\vec{K}\tilde{m}}^a &= \tilde{C}_{1\vec{K}\tilde{m}}^a \left(-\frac{|\vec{K}|}{R_0H_0}e^{-H_0t}\cos\left(\frac{|\vec{K}|}{R_0H_0}e^{-H_0t}\right) + \sin\left(\frac{|\vec{K}|}{R_0H_0}e^{-H_0t}\right) \right) \\ &+ \tilde{C}_{2\vec{K}\tilde{m}}^a \left(\frac{|\vec{K}|}{R_0H_0}\sin\left(\frac{|\vec{K}|}{R_0H_0}e^{-H_0t}\right) + \cos\left(\frac{|\vec{K}|}{R_0H_0}e^{-H_0t}\right) \right) \end{aligned} \quad (V*)$$

where

$$\tilde{C}_{i\vec{K}\tilde{m}}^a = -H_0C_{i\vec{K}\tilde{m}}^a. \quad (VI*)$$

However, it is not necessary to use a full exact solution (V*) to proceed an analysis which we gave before.

Thus the primordial value of $R_{\vec{K}}(t)$ is equal to

$$R_{\vec{K}} = - \left[\frac{H_0}{\frac{d}{dt}(\Phi_m^d)} \delta\Phi_{\vec{K}\tilde{m}}^d \right]_{|t=t^*} \quad (14.182)$$

where $R_{\vec{K}}(t)$ is defined in cosmological perturbation theory as

$$R^{(3)}(t) = 4\frac{\vec{K}^2}{R^2}R_{\vec{K}}(t) \quad (14.183)$$

(see Ref. [7]) and t^* is such that $|\vec{K}| = R(t^*)H_0$, i.e.

$$t^* = \frac{1}{H_0} \ln \left[\frac{|\vec{K}|}{R_0H_0} \right], \quad |\vec{K}| = K. \quad (14.184)$$

The primordial value is of course time-independent. We suppose that fluctuations of Higgs' fields (the initial values of them) are independent and Gaussian.

Thus we can repeat some classical results concerning a power spectrum of primordial perturbations. One gets

$$\left[|\delta\Phi_{\vec{K}\tilde{m}}^d|^2\right] = \left(\frac{H_0}{2\pi}\right)^2. \quad (14.185)$$

Thus

$$P_R(K) \cong \left(\frac{H_0}{2\pi}\right)^2 \sum_{\tilde{m},d} \left(\frac{H_0}{\frac{d}{dt}(\Phi_{\tilde{m}}^d)}\right)^2 \Big|_{t=t^*} \quad (14.186)$$

for t^* given by Eq. (14.184).

Taking a specific situation with concrete groups H, G, G_0 and constants ξ, ζ, r we can in principle calculate exactly $P_R(K)$ and a power index n_s . Thus using some specific software packages (i.e. CMBFAST code) we can obtain theoretical curves of CMB (Cosmic Microwave Background) anisotropy including polarization effects.

Let us give the following remark. The condition for slow-roll evolution for Φ can be expressed in terms of the potential V' . If those are impossible to be satisfied for some models (depending on G, G_0, H, G'_0 and parameters ξ, ζ), we can employ a scenario with a tunnel effect from “false” to “true” vacuum and bubbles coalescence. In the last case the time of an inflation (in the first de Sitter phase) depends on the characteristic time of the coalescence.

Let us consider a more general model of the Universe filled with ordinary (dust) matter, a radiation and a quintessence.

From Bianchi identity we have:

$$\frac{d}{dt}\rho_{\text{tot}} + (\rho_{\text{tot}} + p_{\text{tot}})\frac{3\dot{R}}{R} = 0 \quad (14.187)$$

where

$$\rho_{\text{tot}} = \rho_Q + \tilde{\rho}_m + \tilde{\rho}_r \quad (14.188)$$

$$p_{\text{tot}} = p_Q + \tilde{p}_r. \quad (14.189)$$

Let us suppose that the evolution of radiation, ordinary matter (barionic + cold dark matter) and a quintessence are independent. In this way we get independent conservation laws

$$\frac{d}{dt}(R^3\tilde{\rho}_m) = 0 \quad (14.190)$$

$$\frac{d}{dt}(R^4\tilde{\rho}_r) = 0 \quad (14.191)$$

and

$$\dot{\rho}_Q + (\rho_Q + p_Q)\frac{3\dot{R}}{R} = 0. \quad (14.192)$$

One gets

$$\tilde{\rho}_m = \frac{A}{R^3}, \quad A = \text{const.} \quad (14.193)$$

$$\tilde{\rho}_r = \frac{B}{R^4}, \quad B = \text{const.} \quad (14.194)$$

From Eq. (14.192) we get

$$\dot{\rho}_Q + (1 + W_Q)\rho_Q \cdot \frac{3\dot{R}}{R} = 0. \quad (14.195)$$

Substituting Eqs (14.193–194) into Eqs (14.38) and (14.40), remembering Eqs (14.139) and (14.141–142) one gets

$$\left(\frac{\dot{R}}{R}\right)^2 + k = \frac{1}{3M_{\text{pl}}^2} \left(\frac{A}{R} + \frac{B}{R^2}\right) + \frac{1}{3M_{\text{pl}}^2} \rho_Q R^2 \quad (14.196)$$

$$\frac{3\ddot{R}}{R} = -\frac{1}{M_{\text{pl}}^2} \left(\frac{A}{2R^3} + \frac{2B}{3R^4}\right) - \frac{1}{2M_{\text{pl}}^2} (1 + 3W_Q)\rho_Q \quad (14.197)$$

and from Eq. (14.143)

$$\ddot{Q} + 3H\dot{Q} - \frac{(n+2)A}{M_{\text{pl}}^2 R^3} = 0. \quad (14.198)$$

However, now Q and $U(Q)$ are redefined in such a way that $\frac{1}{M_{\text{pl}}^2}$ factor appears before ρ_Q . This is a simple rescaling. We have of course

$$\tilde{\rho}_m = e^{-(n+2)\Psi} \rho_m = e^{-aQ} \rho_m \quad (14.199)$$

$$\tilde{\rho}_r = e^{-(n+2)\Psi} \rho_r = e^{-aQ} \rho_r. \quad (14.200)$$

Thus we get

$$\rho_m = \frac{A}{R^3} e^{aQ} \quad (14.201)$$

$$\rho_r = \frac{B}{R^4} e^{aQ}. \quad (14.202)$$

Let us consider Eqs (14.196–198) and Eq. (14.192). Combining these equations we get the following formula

$$\frac{(n+2)A}{M_{\text{pl}}^2 R^3} \sqrt{(1 + W_Q)\rho_Q} = 0. \quad (14.203)$$

The one way to satisfy this equation is to put $W_Q = -1$. It is easy to see that if $A = 0$ then Eq. (14.203) is satisfied trivially. The condition $A = 0$ does not imply $B = 0$ and we can still have $B \neq 0$. For such a solution $Q = Q_0 = \text{const.}$ and $Q_0 = \frac{\sqrt{M}}{M_{\text{pl}}} \Psi_0$ found

earlier

$$e^{\Psi_0} = x_0 = \sqrt{\frac{n|\gamma|}{(n+2)\beta}} = \frac{1}{\alpha_s} \left(\frac{m_{\tilde{A}}}{m_{\text{pl}}} \right) \sqrt{\frac{n|\underline{\tilde{P}}|}{(n+2)\tilde{R}(\tilde{\Gamma})}}. \quad (14.204)$$

Thus

$$e^{-aQ_0} = e^{-(n+2)\Psi_0} = \alpha_s^{n+2} \left(\frac{m_{\text{pl}}}{m_{\tilde{A}}} \right)^{n+2} \left(\frac{n+2}{n} \right)^{\frac{n+2}{2}} \left(\frac{\tilde{R}(\tilde{\Gamma})}{|\underline{\tilde{P}}|} \right)^{\frac{n+2}{2}} \quad (14.205)$$

or

$$e^{-aQ_0} = \alpha_s^{n+2} \left(\frac{r}{l_{\text{pl}}} \right) \left(\frac{n+2}{n} \right)^{\frac{n+2}{2}} \left(\frac{\tilde{R}(\tilde{\Gamma})}{|\underline{\tilde{P}}|} \right)^{\frac{n+2}{2}}. \quad (14.206)$$

Let us put $W_Q = -1$ into Eqs (14.196–197). One gets

$$\dot{R}^2 + k = \frac{1}{3M_{\text{pl}}^2} \left(\frac{A}{R} + \frac{B}{R^2} + \rho_Q R^2 \right) \quad (14.207)$$

$$\frac{3\ddot{R}}{R} = -\frac{1}{M_{\text{pl}}^2} \left(\frac{A}{2R^3} + \frac{2B}{3R^4} + \rho_Q \right). \quad (14.208)$$

From Eq. (14.207) one gets

$$\frac{\pm R dR}{\sqrt{\frac{\rho_Q}{3M_{\text{pl}}^2} R^4 - kR^2 + \frac{A}{3M_{\text{pl}}^2} R + \frac{B}{3M_{\text{pl}}^2}}} = dt \quad (14.209)$$

if $A = 0$,

$$\frac{\pm R dR}{\sqrt{\frac{\rho_Q}{3M_{\text{pl}}^2} R^4 - kR^2 + \frac{B}{3M_{\text{pl}}^2}}} = dt. \quad (14.210)$$

Taking $x = R^2$ one gets

$$\int \frac{dx}{\sqrt{\frac{\rho_Q}{3M_{\text{pl}}^2} x^2 - kx + \frac{B}{3M_{\text{pl}}^2}}} = \pm 2(t - t_0) \quad (14.211)$$

and finally

$$\pm \int \frac{dy}{\sqrt{y^2 - \frac{3kM_{\text{pl}}^2}{\sqrt{B\rho_Q}} y + 1}} = \frac{2\sqrt{3}\sqrt{\rho_Q}M_{\text{pl}}}{B}(t - t_0) \quad (14.212)$$

where

$$x = \sqrt{\frac{B}{\rho_Q}} y. \quad (14.213)$$

For a flat case $k = 0$ we find:

$$\pm \int \frac{dy}{\sqrt{y^2 + 1}} = \frac{2\sqrt{3}\sqrt{\rho_Q}M_{\text{pl}}}{B}(t - t_0). \quad (14.214)$$

The last formula can be easily integrated and one gets

$$R = \sqrt[4]{\frac{B}{\rho_Q}} \sqrt{\sinh\left(\frac{2\sqrt{3}\sqrt{\rho_Q}M_{\text{pl}}}{B}(t - t_0)\right)} \quad (14.215)$$

where we take a sign $+$ in the integral on the left hand side of Eq. (14.214).

Let us calculate a Hubble parameter (constant) for our model. One gets

$$H = \frac{\dot{R}}{R} = \sqrt{3}M_{\text{pl}} \left(\frac{\sqrt{\rho_Q}}{B}\right) \text{ctgh}\left(\frac{2\sqrt{3}\sqrt{\rho_Q}M_{\text{pl}}}{B}(t - t_0)\right). \quad (14.216)$$

For a deceleration parameter one obtains

$$q = -\frac{\ddot{R}R}{R^2} = -\frac{(2\sinh^2(u) - \cosh(u))}{\cosh(u)\sinh^{\frac{1}{2}}(u)} \quad (14.217)$$

where

$$u = \frac{2\sqrt{3}\sqrt{\rho_Q}M_{\text{pl}}}{B}(t - t_0). \quad (14.218)$$

Thus we get a spatially flat model of a Universe dominated by a radiation which is expanding and an expansion is accelerating.

Let us take $k = 0$ and $B = 0$ in Eq. (14.209). One gets

$$\frac{\pm R dR}{\sqrt{\frac{\rho_Q}{3M_{\text{pl}}^2}R^4 + \frac{A}{3M_{\text{pl}}^2}R}} = dt \quad (14.219)$$

$$R = \sqrt[3]{\frac{A}{\rho_Q}}x, \quad x = R\sqrt[3]{\frac{\rho_Q}{A}}. \quad (14.220)$$

One finally gets

$$\int \frac{x dx}{\sqrt{x(x^3 + 1)}} = \pm \frac{1}{3M_{\text{pl}}} \sqrt{\rho_Q}(t - t_0). \quad (14.221)$$

Let us notice that

$$H^2 = \left(\frac{\dot{R}}{R}\right)^2 = \frac{1}{3M_{\text{pl}}^2} \left(\frac{A}{R^3} + \rho_Q\right) > \frac{\rho_Q}{M_{\text{pl}}^2} \quad (14.222)$$

$$\frac{\ddot{R}R}{\dot{R}^2} = \frac{(2\rho_Q R^4 - A)R}{2(A + \rho_Q R^3)}. \quad (14.223)$$

Thus the model of the Universe is expanding and accelerating.

Let us consider the integral on the left hand side of Eq. (14.221). One gets after some tedious algebra:

$$\begin{aligned} \int \frac{x dx}{\sqrt{x(x^3+1)}} &= \frac{\sqrt{\sqrt{3}+2}}{12} (9-2\sqrt{3}) \ln W + \frac{9\sqrt{2}}{52} \sqrt{4+\sqrt{3}} (6\sqrt{3}+11) \Pi \\ &- \frac{\sqrt{6}}{598} \left(92\sqrt{4\sqrt{3}+3} (141\sqrt{3}+272) + (9\sqrt{3}-15) \sqrt{598} \sqrt{16+9\sqrt{3}} \right) F \end{aligned} \quad (14.224)$$

where

$$\begin{aligned} W &= \left(-20 - \frac{9\sqrt{3}}{2} + \frac{(16-7\sqrt{3})}{2} \left(\frac{2x+\sqrt{3}-1}{2x-\sqrt{3}-1} \right)^2 \right. \\ &+ 2\sqrt{6-\sqrt{3}} \sqrt{\left[\left(\frac{(2x+\sqrt{3}-1)}{(2x-\sqrt{3}-1)} \right)^2 + 7 - 4\sqrt{3} \right] \left[\left(\frac{(2x+\sqrt{3}-1)}{(2x-\sqrt{3}-1)} \right)^2 - 1 + \frac{\sqrt{3}}{2} \right]} \\ &\times \left. \left(\left(\frac{(2x+\sqrt{3}-1)}{(2x-\sqrt{3}-1)} \right)^2 - 1 \right)^{-1} \right), \end{aligned} \quad (14.225)$$

$$\Pi = \Pi \left(\arccos \left(\left(\frac{(2x+(\sqrt{3}-1))}{(2x-(\sqrt{3}+1))} \right) \sqrt{2(2+\sqrt{3})} \right), (\sqrt{3}-1), \sqrt{\frac{(5-2\sqrt{3})}{13}} \right) \quad (14.226)$$

$$F = F \left(\arccos \left(\left(\frac{(2x+(\sqrt{3}-1))}{(2x-(\sqrt{3}+1))} \right) \sqrt{2(2+\sqrt{3})} \right), \sqrt{\frac{(5-2\sqrt{3})}{13}} \right) \quad (14.227)$$

where Π is an elliptic integral of the third kind and F is an elliptic integral of the first kind

$$F(u, k) = \int_0^u \frac{d\varphi}{\sqrt{1-k^2 \sin^2 \varphi}} \quad (14.228)$$

$$k = \sqrt{\frac{(5-2\sqrt{3})}{13}} \quad (14.229)$$

$$\Pi(u, n, k) = \int_0^u \frac{d\varphi}{(1-n \sin^2 \varphi) \sqrt{1-k^2 \sin^2 \varphi}} \quad (14.230)$$

$$n = \sqrt{3} - 1 \quad (14.231)$$

and with the same modulus as $F(u, k)$

$$k^2 = \frac{5 - 2\sqrt{3}}{13}.$$

However, the most important condition for an existence of a physical solution is coming from the first part of a sum in the right hand side of Eq. (14.224).

We should have

$$W > 0. \quad (14.232)$$

In order to analyze condition (14.232) let us write W in the following way

$$W = \frac{-20 - \frac{9\sqrt{3}}{2} + \frac{(16-7\sqrt{3})}{2}y + 2\sqrt{6-\sqrt{3}}\sqrt{(y+7-4\sqrt{3})\left(y-1+\frac{\sqrt{3}}{2}\right)}}{y-1} \quad (14.233)$$

where

$$y = \left(\frac{2x + \sqrt{3} - 1}{2x - \sqrt{3} - 1} \right)^2. \quad (14.234)$$

Obviously $y \geq 0$.

We have two possibilities:

$$\text{I} \quad y > 1 \quad (14.235)$$

and

$$-20 - \frac{9\sqrt{3}}{2} + \frac{(16-7\sqrt{3})}{2}y + 2\sqrt{6-\sqrt{3}}\sqrt{(y+7-4\sqrt{3})\left(y-1+\frac{\sqrt{3}}{2}\right)} > 0 \quad (14.236)$$

$$\text{II} \quad y < 1 \quad (14.237)$$

and

$$-20 - \frac{9\sqrt{3}}{2} + \frac{(16-7\sqrt{3})}{2}y + 2\sqrt{6-\sqrt{3}}\sqrt{(y+7-4\sqrt{3})\left(y-1+\frac{\sqrt{3}}{2}\right)} < 0. \quad (14.238)$$

The first possibility can easily be solved:

$$\text{I} \quad y > 4.0612 \quad (14.239)$$

or for the second possibility

$$\text{II} \quad \frac{2 - \sqrt{3}}{2} \leq y < 1. \quad (14.240)$$

We have finally

$$0.7916 < x < \frac{\sqrt{3}+1}{2}, \quad (14.241)$$

$$\frac{3\sqrt{3}-5}{2} < x < \frac{1}{2}. \quad (14.242)$$

For

$$x > \frac{\sqrt{3}+1}{2} \quad (14.243)$$

we get

$$x < 3.072. \quad (14.244)$$

Let us notice the following: the function W has a finite limit at $x = \frac{\sqrt{3}+1}{2}$. Thus we can remove a singularity at $x = \frac{\sqrt{3}+1}{2}$. In this way we get

$$0.7916 < x < 3.072 \quad (14.245)$$

or

$$\frac{3\sqrt{3}-5}{2} < x < \frac{1}{2}. \quad (14.246)$$

This simply means that the solution exists only for

$$0.7916 \sqrt[3]{\frac{A}{\rho_Q}} < R < 3.072 \sqrt[3]{\frac{A}{\rho_Q}} \quad (14.247)$$

or

$$\frac{3\sqrt{3}-5}{2} \sqrt[3]{\frac{A}{\rho_Q}} < R < \frac{1}{2} \sqrt[3]{\frac{A}{\rho_Q}}. \quad (14.248)$$

Let us calculate a density of a quintessence energy:

$$\begin{aligned} \rho_Q &= M_{\text{pl}}^2 U(Q_0) = -\frac{1}{2} e^{n\psi_0} (\beta e^{2\psi_0} - |\gamma|) \\ &= \left(\frac{n}{n+2} \right)^{\frac{n}{2}} \left(\frac{m_{\tilde{A}}}{\alpha_s^{2(n+1)}} \right) \left(\frac{m_{\tilde{A}}}{m_{\text{pl}}} \right)^n \left(\frac{|\underline{\tilde{P}}|}{\tilde{R}(\tilde{\Gamma})} \right)^{\frac{n}{2}} |\underline{\tilde{P}}|. \end{aligned} \quad (14.249)$$

An energy density of a matter is equal to

$$\rho_m = \frac{A}{R^3} e^{aQ_0} \simeq \rho_Q e^{aQ_0} \quad (14.250)$$

and

$$\frac{\rho_m}{\rho_Q} \simeq e^{aQ_0} = \left(\frac{1}{\alpha_s} \right)^{n+2} \left(\frac{m_{\tilde{A}}}{m_{\text{pl}}} \right)^{n+2} \left(\frac{n}{n+2} \right)^{\frac{n+2}{2}} \left(\frac{\tilde{R}(\tilde{\Gamma})}{|\underline{\tilde{P}}|} \right)^{\frac{n+2}{2}}. \quad (14.251)$$

Let us consider the behaviour of the “effective” gravitational constant in our model

$$\begin{aligned}
G_{\text{eff}} &= G_N e^{-(n+2)\Psi_0} = G_N e^{-aQ_0} = G_N \left(\frac{1}{x_0} \right)^{n+2} \\
&= G_N \alpha_s^{2(n+2)} \left(\frac{m_{\text{pl}}}{m_{\tilde{A}}} \right)^{\frac{n+2}{2}} \left(\frac{\tilde{R}(\tilde{\Gamma})}{|\tilde{P}|} \right)^{\frac{n+2}{2}} \left(\frac{n+2}{n} \right)^{\frac{n+2}{2}}
\end{aligned} \tag{14.252}$$

Thus G_{eff} is a constant. For we are living in that model we should rescale the constant and consider $G_{\text{eff}}(Q_0)$ (Eq. (14.252)) a Newton constant. Thus

$$G_N = G_0 \alpha_s^{2(n+2)} \left(\frac{m_{\text{pl}}}{m_{\tilde{A}}} \right)^{\frac{n+2}{2}} \left(\frac{n+2}{n} \right)^{\frac{n+2}{2}} \left(\frac{\tilde{R}(\tilde{\Gamma})}{|\tilde{P}|} \right)^{\frac{n+2}{2}} \tag{14.253}$$

where G_0 is a different constant responsible for a strength of gravitational interactions in earlier epochs of an evolution of the Universe. It means we should write

$$G_{\text{eff}} = \left(G_N \alpha_s^{-2(n+2)} \left(\frac{m_{\tilde{A}}}{m_{\text{pl}}} \right)^{\frac{n+2}{2}} \left(\frac{n}{n+2} \right)^{\frac{n+2}{2}} \left(\frac{|\tilde{P}|}{\tilde{R}(\tilde{\Gamma})} \right)^{\frac{n+2}{2}} \right) \cdot e^{-(n+2)\Psi}. \tag{14.254}$$

The obtained solution (i.e. (14.224)) is for $W_Q = -1$. However, this is not an attractor of the dynamical equations. This is similar to the tracker solution of Steinhardt (see Ref. [111]). Moreover we cannot apply his method because our equation for a scalar field Ψ (or Q) is different. It contains an additional term with a trace of the energy-momentum tensor (matter + radiation). Only with a radiation filled Universe our equation is the same (if we identify $\tilde{\rho}_r$ with Steinhardt density of a matter). In this case we could apply Steinhardt tracker solution method and apply his criterion for a potential $U(Q)$. In our case $U(Q)$ is given by Eq. (14.141)

$$U'(Q) = -\frac{M_{\text{pl}}}{2\sqrt{M}} ((n+2)\beta e^{aQ} - n|\gamma|e^{bQ}) \tag{14.255}$$

$$U''(Q) = -\frac{M_{\text{pl}}^2}{4\sqrt{M}} ((n+2)^2\beta e^{aQ} - n^2|\gamma|e^{bQ}). \tag{14.256}$$

The Steinhardt criterion consists in finding

$$\Gamma = \frac{U''U}{(U')^2} = \frac{((n+2)^2\beta e^{aQ} - n^2|\gamma|e^{bQ})(\beta e^{aQ} - |\gamma|e^{bQ})}{((n+2)\beta e^{aQ} - n|\gamma|e^{bQ})^2}. \tag{14.257}$$

To get a tracker solution we need

$$\Gamma \geq 1. \tag{14.258}$$

In our case

$$\Gamma = 1 - \frac{4\beta|\gamma|e^{(a+b)Q}}{((n+2)\beta e^{aQ} - n|\gamma|e^{bQ})^2} \quad (14.259)$$

$$\Gamma \simeq 1 - \frac{4|\gamma|}{(n+2)\beta} e^{(b-a)Q} = 1 - \frac{4|\gamma|}{\beta(n+2)} e^{-\frac{2M_{\text{pl}}}{\sqrt{M}}Q} = 1 - \frac{4|\gamma|}{\beta(n+2)} e^{-2\Psi}. \quad (14.260)$$

Thus asymptotically we are close to $\Gamma = 1$. Moreover the Steinhardt criterion is only a sufficient condition. Our radiation filled Universe seems to be unstable due to appearing of a matter (a dust matter). Moreover a quintessential period of a Universe model has a severe restrictions of a value of a radius. Thus the solution with $W_Q = -1$ and a matter independently evolving cannot evolve forever.

In order to answer a question what is a further evolution of the Universe we come back to Einstein equations with a scalar field Ψ and matter sources. Supposing as usual a Robertson-Walker metric and a spatial flatness of the metric we write once again a Bianchi identity

$$\left(\frac{T^{\text{scal}}{}^{\mu\nu}}{M_{\text{pl}}^2} + \frac{1}{M_{\text{pl}}^2} e^{-(n+2)\Psi} T^{\text{matter}}{}^{\mu\nu} \right)_{;\nu} = 0 \quad (14.261)$$

where

$$T^{\text{matter}}{}^{\mu\nu} = \rho_m u^\mu u^\nu \quad (14.262)$$

is an energy-momentum tensor for a dust matter. One gets

$$\begin{aligned} \frac{1}{M_{\text{pl}}^2} e^{-(n+2)\Psi} \left(\rho_m + \dot{\rho}_m + \rho_m \frac{3\dot{R}}{R} - (n+2)\rho_m \dot{\Psi} \right) \\ + \frac{d\lambda_{c0}}{d\Psi} \dot{\Psi} - \frac{\overline{M}}{M_{\text{pl}}^2} \dot{\Psi} \ddot{\Psi} - \frac{\overline{M}}{M_{\text{pl}}^2} \dot{\Psi}^2 \frac{3\dot{R}}{R} = 0. \end{aligned} \quad (14.263)$$

We make the following ansatz concerning an evolution of a matter density

$$\rho_m = \rho_0 e^{(n+2)\Psi} \quad (14.264)$$

$$\rho_0 = \text{const.} \quad (14.265)$$

In that moment the scalar field Ψ and the matter ρ_m interact nontrivially.

Using Eqs (14.264–265), (14.263) and an equation for a scalar field Ψ one obtains

$$\frac{\rho_0}{M_{\text{pl}}^2} - \frac{(n+2)}{M_{\text{pl}}^2} \rho_0 + \frac{\rho_0}{M_{\text{pl}}^2} \frac{3\dot{R}}{R} = 0. \quad (14.266)$$

Using Eq. (14.266) and Eq. (14.127) one gets

$$\frac{d\Psi}{dt} = \pm \frac{\sqrt{2}M_{\text{pl}}}{\sqrt{3\overline{M}}} \sqrt{\left(\frac{n+1}{M_{\text{pl}}}\right)^2 - \frac{3\rho_0}{M_{\text{pl}}^2} - \frac{3}{2}\lambda_{c0}} \quad (14.267)$$

$$\int \frac{dx}{x\sqrt{\delta + 3\beta x^{n+2} + 3\gamma x^n}} = \pm \frac{M_{\text{pl}}}{\sqrt{\overline{M}}}(t - t_0) \quad (14.268)$$

where

$$\delta = 2 \left(\frac{n+1}{M_{\text{pl}}}\right)^2 - \frac{3\rho_0}{M_{\text{pl}}^2} \quad (14.269)$$

$$x = e^\Psi. \quad (14.270)$$

Using Eq. (14.268) and Eq. (14.127) one gets

$$\frac{d}{dt} \ln R = \frac{(n+1)}{M_{\text{pl}}} \quad (14.271)$$

or

$$R = R_0 e^{\left(\frac{n+1}{M_{\text{pl}}}\right)(t-t_0)}. \quad (14.271a)$$

Let us consider Eq. (14.268) and let us suppose that $x < 1$ (and practically small). In this case we can simplify

$$\int \frac{dx}{x\sqrt{\delta + 3\beta x^{n+2} + 3\gamma x^n}} \simeq \int \frac{dx}{x\sqrt{\delta + 3\gamma x^n}} \quad (14.272)$$

and finally we get

$$\Psi = \frac{1}{2n} \ln \left(\frac{\delta}{3|\gamma|} \right) - \frac{1}{n} \sqrt{\frac{2\delta}{3\overline{M}}} M_{\text{pl}}(t - t_0). \quad (14.273)$$

Thus our prediction that $x < 1$ and is small has been justified. Using Eq. (14.273) one easily gets

$$p_\Psi = \frac{\delta M_{\text{pl}}^2}{3n^2} + \frac{\sqrt{|\gamma|\delta}}{2\sqrt{3}} \exp \left(-\sqrt{\frac{2\delta}{3\overline{M}}} M_{\text{pl}}(t - t_0) \right) \quad (14.274)$$

$$\rho_\Psi = \frac{\delta M_{\text{pl}}^2}{3n^2} - \frac{\sqrt{|\gamma|\delta}}{2\sqrt{3}} \exp \left(-\sqrt{\frac{2\delta}{3\overline{M}}} M_{\text{pl}}(t - t_0) \right). \quad (14.275)$$

It is easy to see that

$$\frac{p_\Psi}{\rho_\Psi} \xrightarrow{t \rightarrow \infty} 1. \quad (14.276)$$

Thus in this model we have to do asymptotically (practically very quickly) with a stiff matter equation of state:

$$p = \rho. \quad (14.277)$$

Simultaneously an energy density for a scalar field Ψ is dominated by a kinetic energy which is practically constant and

$$\frac{1}{2}\dot{Q}^2 \cong \frac{\delta M_{\text{pl}}^2}{3n^2}. \quad (14.278)$$

In this sense a dark matter generated by Ψ in the model is in some sense K -essence (not a quintessence).

Let us calculate an energy density for a matter

$$\rho_m = \rho_0 \left(\frac{\delta}{3|\gamma|} \right)^{\frac{(n+2)}{2n}} \exp \left(-\frac{(n+2)}{n} \sqrt{\frac{2\delta}{3M}} M_{\text{pl}}(t - t_0) \right). \quad (14.279)$$

The effective gravitational constant reads

$$G_{\text{eff}} = G_N \left(\frac{3|\gamma|}{\delta} \right)^{\frac{(n+2)}{n}} \exp \left(\left(\frac{n+2}{n} \right) \sqrt{\frac{2\delta}{3M}} M_{\text{pl}}(t - t_0) \right). \quad (14.280)$$

It is easy to write R as a function of Ψ :

$$R = R_0 \exp \left(\frac{n+1}{M_{\text{pl}}^2} \sqrt{M} \int_{x_0}^{e^\Psi} \frac{dx}{x \sqrt{\delta + 3\beta x^{n+2} + 3\gamma x^n}} \right) \quad (14.281)$$

where

$$x_0 = \left(\frac{3|\gamma|}{\delta} \right)^{\frac{2n}{n+2}}. \quad (14.282)$$

Eq. (14.281) can be easily reduced to the Eq. (14.271) using Eq. (14.268).

Thus an energy density of matter goes to zero in an exponential way. The effective gravitational constant is growing exponentially. What does it mean for the future of the Universe? First of all the Universe will be very diluted after some time of such an evolution. Secondly, a relative strength of gravitational interactions will be stronger. This means that if we take substrat particles as cluster of galaxies then in a quite short time any cluster will be a lonely island in the Universe without any communication with other clusters. All the clusters will be beyond a horizon. On the level of a single cluster the strength of gravitational interactions will be stronger and eventually they collapse to form a black hole. In an intermediate time a gravitational dynamics on the level of galaxies' clusters will be governed by a Newtonian dynamics (nonrelativistic) with a gravitational potential corrected by a cosmological constant. The cosmological constant induces a positive pressure (a stiff matter) and will be important up to a moment of sufficiently large gravitational constant. After a sufficiently long time only a classical Newtonian gravity will drive a dynamics of galaxies' clusters. Let us roughly

estimate this effect. In order to do this let us suppose that

$$\rho_\Psi = p_\Psi = \frac{\delta M_{\text{pl}}^2}{3n^2} = \text{const.} \quad (14.283)$$

In an energy momentum we get

$$T^{\mu\nu}_{\text{scal}} = 2 \cdot \frac{\delta M_{\text{pl}}^2}{3n^2} u^\mu u^\nu - \frac{\delta M_{\text{pl}}^2}{3n^2} g^{\mu\nu}. \quad (14.284)$$

However, we have in the place of Newtonian constant G_N a G_{eff} which is a function of cosmological time and in the equation for gravitational field for $g_{\mu\nu}$ we should put

$$\frac{1}{(M_{\text{pl}}^{\text{eff}})^2} T^{\mu\nu}_{\text{scal}} = \frac{\delta}{3n^2} \left(\frac{M_{\text{pl}}}{M_{\text{pl}}^{\text{eff}}} \right)^2 (2u^\mu u^\nu - g^{\mu\nu}) \quad (14.285)$$

where

$$\begin{aligned} M_{\text{pl}}^{\text{eff}} &= \frac{1}{\sqrt{8\pi G_{\text{eff}}}} = M_{\text{pl}} \left(\frac{\alpha_s}{m_{\text{pl}} m_{\tilde{A}}} \right)^{n+2} \left(\frac{2(n+1)^2 - 3\rho_0}{3|\tilde{P}|} \right)^{\frac{n+2}{2}} \\ &\times \exp \left[\left(\frac{1}{2} \left(\frac{n+2}{2} \right) \sqrt{\frac{2\delta}{3M}} \right) M_{\text{pl}}(t - t_0) \right]. \end{aligned} \quad (14.286)$$

Thus

$$\frac{1}{(M_{\text{pl}}^{\text{eff}})^2} T^{\mu\nu}_{\text{scal}} = \tilde{A} e^{-\kappa(t-t_0)} (2u^\mu u^\nu - g^{\mu\nu}) \quad (14.287)$$

where

$$\kappa = \frac{1}{2} \left(\left(\frac{n+2}{2} \right) \sqrt{\frac{2\delta}{3M}} \right) \quad (14.288)$$

$$\tilde{A} = \frac{3n^2}{\delta} \left(\frac{\alpha_s}{m_{\text{pl}} m_{\tilde{A}}} \right)^{-(n+2)} \left(\frac{2(n+1)^2 - 3\rho_0}{3|\tilde{P}|} \right)^{-\left(\frac{n+2}{2}\right)}. \quad (14.289)$$

Consider the following case (the justification will be given below):

$$\frac{1}{(M_{\text{pl}}^{\text{eff}})^2} T^{\mu\nu}_{\text{scal}} = -\tilde{A} e^{-\kappa(t-t_0)} g^{\mu\nu}. \quad (14.290)$$

Now let us solve the following problem. Let $\bar{A}$ be a value $\tilde{A} e^{-\kappa(\bar{t}-t_0)}$ in an established cosmological time $t = \bar{t}$ and the same for $\bar{G} = G_{\text{eff}}(\bar{t})$. Let us consider a gravitational field of a point mass M_0 static and spherically symmetric. Consider a metric

$$ds^2 = B(r) dt^2 - \frac{dr^2}{B(r)} - r^2 (d\theta^2 + \sin^2 \theta d\phi^2) \quad (14.291)$$

in spherical coordinates and Einstein equations

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = -\bar{\Lambda}g_{\mu\nu} \quad (14.292)$$

for the metric (14.291) where

$$\bar{\Lambda} = \tilde{A}e^{-\kappa(\bar{t}-t_0)}. \quad (14.293)$$

Via a standard procedure one obtains

$$B(r) = 1 - \frac{2\bar{G}M_0}{r} - \frac{\bar{\Lambda}r^2}{3}. \quad (14.294)$$

Thus an effective nonrelativistic Newton-like gravitational potential has a shape

$$V(r) = \frac{\bar{G}M_0}{r} + \frac{\bar{\Lambda}r^2}{6}. \quad (14.295)$$

Remembering that $\bar{G}$ is growing exponentially in time and $\bar{\Lambda}$ is decaying in time exponentially we see that the second term is important only for a short time. For a contemporary time

$$\bar{\Lambda} \simeq 10^{-52}\text{m}^{-2} \quad (14.296)$$

according to the Perlmutter data and it could be important on the level of a size of galaxies' cluster. However, in a short time it will be negligible even on that level. Let us calculate a Schwarzschild radius for a mass M_0 with an effective gravitational constant G_{eff} . One gets

$$\begin{aligned} r_g^{\text{eff}} &= \sqrt{2M_0\bar{G}} = \sqrt{2M_0G_{\text{eff}}(\bar{t})} = r_g \left(\frac{m_{\text{pl}}m_{\tilde{A}}}{\alpha_s} \right)^{n+2} \\ &\times \left(\frac{3|\tilde{P}|}{2(n+1)^2 - 3\rho_0} \right)^{\frac{n+2}{2}} \exp \left[\left(\frac{1}{2} \left(\frac{n+2}{2} \right) \sqrt{\frac{2\delta}{3\bar{M}}} \right) M_{\text{pl}}(t - t_0) \right] \end{aligned} \quad (14.297)$$

Thus r_g^{eff} (calculated for an effective gravitational constant $G_{\text{eff}}(\bar{t})$) is growing exponentially. It means that after some time it will be of an order of size of galaxies' cluster. Then galaxies' clusters collapse to form black holes. The time of a collapse is easy to calculate.

$$r_g^{\text{eff}} = r_{\text{cluster}} \quad (14.298)$$

$$\begin{aligned} t_{\text{collapse}} &= t_0 + (n+2)\sqrt{6\bar{M}} \left[\ln \left(\frac{r_{\text{cluster}}}{r_g} \right) + (n+2) \ln \left(\frac{m_{\text{pl}}m_{\tilde{A}}}{\alpha_s} \right) \right. \\ &\quad \left. + \left(\frac{n+2}{2} \right) \ln \left(\frac{3|\tilde{P}|}{2(n+1)^2 - 3\rho_0} \right) \right] \cdot \left[\sqrt{\delta}(n+2)M_{\text{pl}} \right]^{-1} \end{aligned} \quad (14.299)$$

where r_g is a gravitational radius of a mass of a galaxies' cluster with $G = G_N$.

Let us calculate a Hubble constant and a deceleration parameter for our model. One gets

$$H = \frac{\dot{R}}{R} = (n+1)M_{\text{pl}} \quad (14.300)$$

$$q = -\frac{\ddot{R}R}{\dot{R}^2} = -1. \quad (14.301)$$

Summing up we get the following scenario of an evolution. The Universe starts in a “false” vacuum state and its stable evolution is driven by a “cosmological constant” formed from $\tilde{R}(\tilde{\Gamma})$, $\overline{P}$, $V(\Phi_{\text{crt}}^1)$ and Ψ_1 being a minimal solution to an effective self-interacting potential for Ψ . This evolution is stable against small perturbation for Ψ and R . The evolution is exponential and a Hubble constant is calculable in terms of our theory (the nonsymmetric Jordan–Thiry theory). This evolution ends at a moment the “false” vacuum state changes into a “true” vacuum state. In that moment an energy of a vacuum is released into radiation (and a matter). The Universe is reheating and a big-bang scenario begins at a very hot stage. Moreover the evolution is still governed by a scalar field Ψ which attains a new minimum of an effective potential. The effective potential is different and a new value of Ψ , a Ψ_0 , is different too. The evolution is exponential and a new Hubble constant H_1 is also calculable in terms of the underlying theory. In this background we have an evolution of a multiplet of scalar fields Φ_m^a from Φ_{crt}^1 to Φ_{crt}^0 . This goes to a spectrum of fluctuations calculable in the theory. After that time the field Ψ is slowly changing $\dot{\Psi} \approx 0$ and the temperature of the Universe is going down. The evolution of a radiation is in the second inflationary stage adiabatic and afterwards during next phase ($\dot{\Psi} \simeq 0$) nonadiabatic. The last means there is an energy exchange between radiation and a scalar field Ψ . The total amount of an inflation $\overline{N}_{\text{tot}}$ is calculable in the theory. The evolution of a factor $R(t)$ is governed by a simple elementary function of a small rise. After this the radiation condenses into a matter (a dust with zero pressure) and a matter (a dust), a radiation and a scalar field evolve adiabatically without interaction among them. The evolution of a scalar field is governed by a quintessence, i.e. $p_Q = W_Q \rho_Q$, where $W_Q \cong -1$.

The quintessence field Q is a normalized scalar field Ψ . During a radiation era ($\dot{\Psi} \simeq 0$) an equation of state for a scalar field Ψ is different: $p_\Psi = \frac{4}{3}\rho_\Psi$ (similar to stiff matter equation of state). The scale factor $R(t)$ evolves according to some elementary function built from hyperbolic function in radiation + quintessence scenario and according to complicated function formed from the first and the second elliptic integrals (after reversing of those functions). It expands and accelerates. The solution with radiation and quintessence has a restricted behaviour (Eq. (14.247–248)). The energy density of radiation is going to zero. Only a quintessence energy density is constant. The energy of a quintessence is a potential energy dominated. However, the solution cannot be continued up to the end for an interaction between a matter and a scalar field starts to be important. The effective gravitational constant remains constant during the period. Let us remind to the reader that during inflationary epochs

the effective gravitational constant was constant. Only during a radiation era it was slightly changing. In order to keep the Universe to evolve forever it is necessary to find a solution with an interaction between a matter and a scalar field Ψ . Using an appropriate ansatz for an evolution of an energy density of a matter we get such a solution. The scale factor $R(t)$ is exponentially expanding.

An energy density of a matter is going to zero and an energy density of a scalar field is approaching (exponentially) a constant. However, now a scalar field Ψ forms a different form of a matter—a stiff matter with an equation of a state $\rho_\Psi \simeq p_\Psi$ ($W_\Psi \simeq 1$). The effective gravitational constant is growing exponentially in time. In contradistinction with the previous period of an evolution the scalar field Ψ forms a matter with a kinetic energy dominance (i.e. K -essence). In the previous period we have to do with a potential energy dominance (i.e. quintessence).

Let us give some remarks on a spatial curvature of a model of the Universe. In the first period (inflationary scenario) the spatial curvature has to be zero and any perturbations of initial conditions cannot change it. In next periods it is natural to suppose that it is not changed. Thus our model of a Universe is spatially flat. In the moment of change of a phase of an evolution we should apply a matching conditions for an energy density and for a scale factor (a radius of a model of a Universe). However, an evolution in such moments undergoes phase transitions for a law of an evolution of a scalar field and a scale factor, which can be considered as a second order or even a first order phase transition. It is worth to notice that in formulae concerning Hubble constants, deceleration parameters, gravitational constants (effective) meet some parameters from unification theory of fundamental interactions and from cosmology. It seems that we are on a right track to give an account for large number hypothesis by P. Dirac. This demands of course some investigations especially in developing nonsymmetric Kaluza–Klein (Jordan–Thiry) theory. It is also interesting to mention that the scalar field Ψ plays many rôles in our theory. It works as an inflation field during an inflation epoch. (However Higgs' fields Φ_b^a play an important rôle in creating fluctuations spectrum.) It plays also a rôle of a quintessence and a K -essence, except its influence on an effective gravitational constant and on an effective scale of masses in unification of fundamental interactions. This field is massive getting masses from several mechanisms. During inflation phases it acquires mass due to spontaneous symmetry breaking, different in both epochs. It gets a mass from a cosmological background mentioned earlier in this section. It can get a mass from interactions with Higgs' fields (see sec. 10). Thus except its cosmological importance the fluctuations of a scalar field around its cosmological value could be detected (in principle) as massive scalar particles of a large mass. In this way an effective gravitational potential in nonrelativistic limit takes a form

$$V(r) = \frac{G_N M_0}{r} \left(1 - \frac{\alpha}{r} e^{-\left(\frac{r}{r_0}\right)}\right) e^{\kappa(t-t_0)} + \frac{\tilde{A} r^2}{6} e^{-\kappa(t-t_0)} \quad (14.302)$$

where $\kappa, \tilde{A}$ are given by Eqs (14.288–289), r_0 is a range of massive scalar $\delta\Psi$ (a fluctuation of a scalar field Ψ in a cosmological background). This could be detected as a tiny change of a Newton constant in gravitational law (e.g. the fifth force) on short

distances. Let us notice that on short distances and on short time scales an effective gravitational potential reads:

$$V(r) = \frac{G_N M_0}{r} \left(1 - \frac{\alpha}{r} e^{-\left(\frac{r}{r_0}\right)} \right). \quad (14.303)$$

On large distances and short time scales

$$V(r) = \frac{G_N M_0}{r} + \frac{\bar{\Lambda} r^2}{6}. \quad (14.304)$$

On short distances and large time scales

$$V(r) = \frac{G_N M_0}{r} \left(1 - \frac{\alpha}{r} e^{-\left(\frac{r}{r_0}\right)} \right) e^{\kappa(t-t_0)}. \quad (14.305)$$

On intermediate distances and large time scales

$$V(r) = \frac{G_N M_0}{r} e^{\kappa(t-t_0)}. \quad (14.306)$$

The latest being a cause to collapse a cluster of galaxies into a black hole. It seems also possible to consider an effective coupling constant α_s^{eff} for α_s enters $m_{\tilde{A}}$ and $m_{\tilde{A}}^{\text{eff}}$. However

$$m_{\tilde{A}}^{\text{eff}} = e^{-\Psi} \left(\frac{\alpha_s}{r} \right), \quad (14.307)$$

r is a radius of a manifold G/G_0 (e.g. a radius of a sphere S^2). If we rewrite Eq. (14.307) as

$$\frac{m_{\tilde{A}}^{\text{eff}}}{\alpha_s^{\text{eff}}} = \frac{1}{r} e^{-\Psi}, \quad (14.308)$$

it would be quite difficult to distinguish a drift of $m_{\tilde{A}}^{\text{eff}}$ from α_s^{eff} drift. Only a quotient has a definite dependence on the scalar field Ψ . Especially it is interesting to consider a drift of α_{em} (a fine structure constant) reported from several sources. However, without extending our theory to include fermion (spinor) fields this is impossible. Thus it is necessary to construct a nonsymmetric-geometric version with supersymmetry and supergravity including noncommutative (anticommutative) coordinates to settle the problem. The scalar field Ψ can form an infinite tower of massive scalar fields which could have important astrophysical and high-energy consequences (see sec. 16).

Let us consider Eqs (4.51) and (4.56) and changing a base in the Lie algebra $\mathfrak{h}$ we get

$$\mu_i^a = \delta_i^a \quad (14.309)$$

and

$$\alpha_i^c = \delta_i^c. \quad (14.310)$$

Simultaneously for $G \subset H$ ($\mathfrak{g} \subset \mathfrak{h}$) we get

$$f_{\tilde{a}\tilde{b}}^{\hat{i}} = C_{\tilde{a}\tilde{b}}^{\hat{i}} \quad (14.311a)$$

$$f_{\tilde{a}\tilde{b}}^{\tilde{d}} = C_{\tilde{a}\tilde{b}}^{\tilde{d}} \quad (14.311b)$$

where $\tilde{a}, \tilde{b}, \tilde{c}$ refer to the complement $\mathfrak{m}$ in $\mathfrak{g}$ ($\mathfrak{g} = \mathfrak{g}_0 + \mathfrak{m}$), $\hat{i}, \hat{j}, \hat{k}$ to the subalgebra $\mathfrak{g}_0$ of $\mathfrak{g}$. Let us suppose a symmetry requirement

$$[\mathfrak{m}, \mathfrak{m}] \subset \mathfrak{g}_0. \quad (14.312)$$

From Eq. (14.312) it follows that

$$C_{\tilde{b}\tilde{c}}^{\tilde{a}} = f_{\tilde{b}\tilde{c}}^{\tilde{a}} = 0. \quad (14.313)$$

Eq. (4.51) looks like

$$\Phi_{\tilde{b}}^c C_{\tilde{i}\tilde{a}}^{\tilde{b}} = \Phi_{\tilde{a}}^b C_{\tilde{i}\tilde{b}}^c. \quad (14.314)$$

Eq. (4.55)

$$H_{\tilde{a}\tilde{b}}^c = C_{ab}^c \Phi_{\tilde{a}}^a \Phi_{\tilde{b}}^b - C_{\tilde{a}\tilde{b}}^c - \Phi_{\tilde{a}}^c C_{\tilde{a}\tilde{b}}^{\tilde{d}}. \quad (14.315)$$

Let us come back to Eq. (14.157).

For our inflation driving agent is a scalar field Ψ we do not carry any backreaction of Higgs' fields and we can consider some nonstandard scenarios of an evolution of those fields. Thus we can consider an evolution which is not a slow-roll evolution. Let us suppose that

$$\frac{d}{dt}(\Phi_{\tilde{m}}^a) \simeq 0. \quad (14.316)$$

In this way we get from (14.316)

$$\frac{d}{dt}(L_{\tilde{b}}^{d0})_{av} + \frac{e^{n\Psi_1}}{2r^2} l^{db} \left\{ \frac{\delta V'}{\delta \Phi_{\tilde{n}}^b} g_{\tilde{b}\tilde{n}} \right\}. \quad (14.317)$$

The next step in simplification is such that we collapse all the degrees of freedom of Higgs' fields into one

$$\Phi_{\tilde{m}}^a = \delta_{\tilde{m}}^a \Phi. \quad (14.318)$$

In this way constraints (14.314) are satisfied trivially and Eq. (14.315) is as follows:

$$H_{\tilde{a}\tilde{b}}^c = C_{\tilde{a}\tilde{b}}^c (\Phi^2 - 1 - \Phi). \quad (14.319)$$

In this way $V = V'$ (no constraints) and

$$V = \frac{A}{\alpha_s^2} (\alpha_s^2 \Phi^2 - 1 - \alpha_s \Phi)^2 \quad (14.320)$$

where

$$A = \left[l_{ab} \left(2g^{[\tilde{m}\tilde{n}]} g^{[\tilde{a}\tilde{b}]} C_{\tilde{m}\tilde{n}}^c C_{\tilde{a}\tilde{b}}^b - C_{\tilde{m}\tilde{n}}^b g^{\tilde{a}\tilde{n}} g^{\tilde{b}\tilde{m}} \tilde{L}_{\tilde{a}\tilde{b}}^a \right) \right]_{av} \quad (14.321)$$

and

$$L_{\tilde{a}\tilde{b}}^a = \tilde{L}_{\tilde{a}\tilde{b}}^a \left(\alpha_s \Phi^2 - \frac{1}{\alpha_s} - \Phi \right) \quad (14.322)$$

$$l_{dc} g_{\tilde{m}\tilde{b}} g^{\tilde{c}\tilde{m}} \tilde{L}_{\tilde{c}\tilde{a}}^d + l_{cd} g_{\tilde{a}\tilde{m}} g^{\tilde{m}\tilde{c}} \tilde{L}_{\tilde{b}\tilde{c}}^d = 2l_{cd} g_{\tilde{a}\tilde{m}} g^{\tilde{m}\tilde{c}} C_{\tilde{b}\tilde{c}}^d. \quad (14.323)$$

We can get the following identity

$$l_{dc} g^{\tilde{c}\tilde{q}} g^{\tilde{a}\tilde{p}} \tilde{L}_{\tilde{c}\tilde{a}}^d C_{\tilde{p}\tilde{q}}^c + l_{cd} g^{\tilde{p}\tilde{c}} g^{\tilde{q}\tilde{b}} \tilde{L}_{\tilde{b}\tilde{c}}^d C_{\tilde{p}\tilde{q}}^c = 2l_{cd} g^{\tilde{p}\tilde{c}} g^{\tilde{q}\tilde{b}} C_{\tilde{b}\tilde{c}}^d C_{\tilde{p}\tilde{q}}^c. \quad (14.324)$$

Using Eq. (14.317) we get the following equation for Φ :

$$\frac{d^2 \Phi}{dt^2} + \frac{e^{n\Psi_1}}{2r^2} \frac{\delta V}{d\Phi} = 0 \quad (14.325)$$

if the following constraints are satisfied

$$\int \sqrt{|\tilde{g}|} d^n x \left(g_{\tilde{q}\tilde{c}} \tilde{l}^{\tilde{p}\tilde{c}} + l^{\tilde{c}\tilde{n}} g_{\tilde{c}\tilde{n}} \delta_{\tilde{q}}^{\tilde{p}} \right) = 2V_2 \delta_{\tilde{c}}^{\tilde{e}} \delta_{\tilde{q}}^{\tilde{p}}. \quad (14.326)$$

Thus from Eq. (14.325) one gets a first integral of motion

$$\frac{\dot{\Phi}^2}{2} + \frac{e^{n\Psi_1}}{2r^2} V = \frac{B}{2} = \text{const.} \quad (14.327)$$

From Eq. (14.327) one gets

$$\int_{\Phi_0}^{\Phi} \frac{d\Phi}{\sqrt{-\frac{e^{n\Psi_1}}{r^2 \alpha_s^2} A (\alpha_s^2 \Phi^2 - \alpha_s \Phi - 1)^2 + B}} = \pm (t - t_0) \quad (14.328)$$

and finally

$$\int_{\varphi_0}^{\varphi} \frac{d\varphi}{\sqrt{(a - \varphi^2 + \varphi + 1)(a + \varphi^2 - \varphi - 1)}} = \pm \sqrt{\frac{A e^{n\Psi_1}}{r^2}} (t - t_0) \quad (14.329)$$

where

$$a = \alpha_s \sqrt{\frac{B r^2}{A e^{n\Psi_1}}} \quad (14.330)$$

$$\Phi = \frac{1}{\alpha_s} \varphi. \quad (14.331)$$

In order to find an integral on the left hand side of Eq. (14.329) and its inverse function we use a uniformization theory of algebraic curves. Let

$$f(\varphi) = (a - \varphi^2 + \varphi + 1)(a + \varphi^2 - \varphi - 1) = -\varphi^4 + 2\varphi^3 + \varphi^2 - 2\varphi + (a^2 - 1). \quad (14.332)$$

One easily notices that

$$\varphi_0 = \frac{1 + \sqrt{5 + 4a}}{2} \quad (14.333)$$

is a root of the polynomial $f(\varphi)$ (a real root).

In this way one gets

$$\varphi = \frac{f'(\varphi_0)}{4 \left(P(z) - \frac{1}{24} f''(\varphi_0) \right)} + \varphi_0 \quad (14.334)$$

where

$$P(z) = P(z; g_2, g_3) \quad (14.335)$$

is a P Weierstrass function with invariants:

$$g_2 = \frac{5 - 3a^2}{3} \quad (14.336)$$

$$g_3 = \frac{a^2}{12} + \frac{10}{27}. \quad (14.337)$$

One easily finds

$$f'(\varphi_0) = \frac{1}{2}(5 + 3a)\sqrt{5 + 4a} \quad (14.338)$$

$$f''(\varphi_0) = -2(5 + 6a) \quad (14.339)$$

and

$$z = \int_{\varphi_0}^{\varphi} [f(x)]^{-\frac{1}{2}} dx. \quad (14.340)$$

In this way one gets

$$\Phi(t) = \frac{(5 + 3a)\sqrt{5 + 4a}}{8\alpha_s \left(P(z) + \frac{1}{12}(5 + 6a) \right)} + \frac{1 + \sqrt{5 + 4a}}{2\alpha_s} \quad (14.341)$$

where

$$z = \pm \sqrt{\frac{Ae^{n\Psi_1}}{r^2}}(t - t_0). \quad (14.342)$$

Let us come back to the potential V Eq. (14.320) and let us look for its critical points. One gets

$$\frac{dV}{d\Phi} = \frac{2A}{\alpha_s} (\alpha_s^2 \Phi^2 - \alpha_s \Phi - 1) (2\alpha_s \Phi - 1) \quad (14.343)$$

and from

$$\frac{dV}{d\Phi} = 0 \quad (14.344)$$

easily finds

$$\Phi_1 = \frac{1}{2\alpha_s} \quad (14.345a)$$

$$\Phi_2 = \frac{1 - \sqrt{5}}{2\alpha_s} \quad (14.345b)$$

$$\Phi_3 = \frac{1 + \sqrt{5}}{2\alpha_s} \quad (14.345c)$$

One obtains

$$V(\Phi_1) = \frac{25A}{16\alpha_s^2} \quad (14.346a)$$

$$V(\Phi_2) = 0 \quad (14.346b)$$

$$V(\Phi_3) = 0. \quad (14.346c)$$

It is easy to see that Φ_1 is a maximum and Φ_2 and Φ_3 are minima of the potential V .

However, in our simplified model of an evolution of Higgs' fields we have not a second (local) minimum. Thus in order to mimic a real situation we start an evolution of a Higgs field from the value

$$\Phi = 0. \quad (14.347)$$

In this case

$$V(0) = \frac{A}{\alpha_s^2}. \quad (14.348)$$

Thus

$$\Phi(t_{\text{initial}}) = 0 \quad (14.349)$$

$$\Phi(t_{\text{end}}) = \frac{1 + \sqrt{5}}{2\alpha_s}. \quad (14.350)$$

Coming back to Eq. (14.329) we get

$$\int_0^{\frac{1+\sqrt{5}}{2}} \frac{d\varphi}{\sqrt{(a - \varphi^2 + \varphi + 1)(a + \varphi^2 - \varphi - 1)}} = \sqrt{\frac{Ae^{n\Psi_1}}{r^2}} (t_{\text{end}} - t_{\text{initial}}). \quad (14.351)$$

Thus the amount of inflation obtained in our evolution is equal to

$$\overline{N}_0 = H_0 \sqrt{\frac{r^2}{Ae^{n\Psi_1}}} \int_0^{\frac{1+\sqrt{5}}{2}} \frac{d\varphi}{\sqrt{(a - \varphi^2 + \varphi + 1)(a + \varphi^2 - \varphi - 1)}}. \quad (14.352)$$

Supposing simply

$$a < \frac{5}{4} \quad (14.353)$$

one gets

$$\bar{N}_0 = H_0 \sqrt{\frac{2r^2}{5Aae^{n\Psi_1}}} \left(\int_0^{\varphi_1} \frac{d\theta}{\sqrt{1 - \left(\frac{4a+5}{10}\right) \sin^2 \theta}} - \int_0^{\varphi_2} \frac{d\theta}{\sqrt{1 - \left(\frac{4a+5}{10}\right) \sin^2 \theta}} \right) \quad (14.354)$$

where

$$\cos^2 \varphi_1 = \frac{1}{5 + 4a} \quad (14.355)$$

$$\cos^2 \varphi_2 = \frac{5}{5 + 4a} \quad (14.356)$$

or

$$\bar{N}_0 = H_0 \sqrt{\frac{2r^2}{5Aae^{n\Psi_1}}} \int_{\varphi_2}^{\varphi_1} \frac{d\theta}{\sqrt{1 - \left(\frac{4a+5}{10}\right) \sin^2 \theta}} \quad (14.357)$$

or

$$\bar{N}_0 = \frac{H_0}{\sqrt{5\alpha_s}} \sqrt[4]{\frac{4r^2}{ABe^{n\Psi_1}}} \int_{\varphi_2}^{\varphi_1} \frac{d\theta}{\sqrt{1 - \left(\frac{4a+5}{10}\right) \sin^2 \theta}}. \quad (14.358)$$

In order to get 60-fold inflation

$$\bar{N}_0 \simeq 60 \quad (14.359)$$

we should play with parameters in our theory. Let us notice that in our simple model the constant α_1 equals

$$\alpha_1 = m_{\tilde{A}}^2 \left(\frac{m_{\tilde{A}}}{m_{\text{pl}}} \right)^2 \left(\frac{A}{\alpha_s^6} \right) = \frac{1}{r^2} \left(\frac{l_{\text{pl}}}{r} \right)^2 \left(\frac{A}{\alpha_s^6} \right) \quad (14.360)$$

(see Eqs (14.32–33a).

Let us consider a slow-roll dynamic of Higgs' field in this simplified model. From Eq. (14.165) one gets

$$\frac{d}{dt} \left(\Phi_{\tilde{p}}^f \right) = - \frac{e^{n\Psi_1}}{12V_2 r^2 H_0} \left(l^{fb} g_{\tilde{p}\tilde{n}} + l^{bf} g_{\tilde{n}\tilde{p}} \right) \frac{\delta V'}{\delta \Phi_{\tilde{n}}^b}. \quad (14.361)$$

Using our simplifications from Eqs (14.309–319) and Eqs (14.300–306) one gets

$$\frac{d\Phi}{dt} = - \frac{e^{n\Psi_1}}{12r^2 H_0} \frac{dV}{d\Phi} \quad (14.362)$$

supposing a condition

$$\delta_{\tilde{p}}^f = l^{f\tilde{n}} g_{\tilde{p}\tilde{n}} + l^{\tilde{n}f} g_{\tilde{n}\tilde{p}} \quad (14.363)$$

or

$$\frac{d\Phi}{dt} + \frac{e^{n\Psi_1} A}{6\alpha_s r^2 H_0} \left(\alpha_s^2 \Phi^2 - \alpha_s \Phi - 1 \right) (2\alpha_s \Phi - 1) = 0. \quad (14.364)$$

Changing the dependent variable into

$$\varphi = \alpha_s \Phi \quad (14.365)$$

one gets

$$\frac{d\varphi}{dt} + b(\varphi^2 - \varphi - 1)(2\varphi - 1) = 0 \quad (14.366)$$

where

$$b = \frac{e^{n\Psi_1} A}{6r^2 H_0} > 0. \quad (14.367)$$

From Eq. (14.366) one finds

$$\frac{\varphi^2 - \varphi - 1}{(\varphi - \frac{1}{2})^2} = e^{-5b(t-t_0)} \quad (14.368)$$

and finally

$$\Phi(t) = \frac{1}{2\alpha_s} \left[1 \pm \sqrt{\frac{5}{1 - e^{-5b(t-t_0)}}} \right]. \quad (14.369)$$

One easily notices that for $(t - t_0) \rightarrow \infty$

$$\Phi(t) \rightarrow \frac{1}{\alpha_s} \left(\frac{1 \pm \sqrt{5}}{2} \right). \quad (14.370)$$

It simply means that a slow-roll inflation never ends.

If we suppose that an inflation starts at $\varphi = 3$, i.e. for

$$t_{\text{initial}} = t_0 + \frac{1}{5b} \ln \frac{5}{4}, \quad (14.371)$$

it is evident that to complete it we need an eternity.

Summing up we get in our simplified model a finite amount of an inflation for a nonstandard dynamic of a Higgs field and an infinite amount of inflation for a slow-roll dynamic. The truth probably is in a middle.

Probably it would be necessary to consider a full equation for a Higgs field in a simplified model

$$\frac{d^2\Phi}{dt^2} - 3H_0 \frac{d\Phi}{dt} - 3bH_0 (\alpha_s^2 \Phi^2 - \alpha_s \Phi - 1)(2\alpha_s \Phi - 1) = 0 \quad (14.372)$$

or

$$\frac{d^2\varphi}{dt^2} - 3H_0 \frac{d\varphi}{dt} - 3b\alpha_s H_0 (\varphi^2 - \varphi - 1)(2\varphi - 1) = 0. \quad (14.372a)$$

Changing an independent variable from t to $\tau = H_0 t$ one gets

$$\frac{d^2\varphi}{d\tau^2} - 3 \frac{d\varphi}{d\tau} - 3 \frac{b\alpha_s}{H_0} (\varphi^2 - \varphi - 1)(2\varphi - 1) = 0. \quad (14.372b)$$

Let

$$\frac{3b\alpha_s}{H_0} = \bar{a} \quad (14.373)$$

and let us change a dependent variable φ into

$$z = 2\varphi - 1 \quad (14.374)$$

$$\varphi = \frac{z+1}{2}. \quad (14.375)$$

One gets

$$\frac{d^2 z}{d\tau^2} - 3\frac{dz}{d\tau} - \frac{\bar{a}}{4}z^3 - \frac{(15\bar{a})}{2}z = 0. \quad (14.376)$$

Now let us take a special value for $\bar{a} = -\frac{4}{15}$ and let us change z into r :

$$z = i\sqrt{5}r. \quad (14.377)$$

One gets

$$\frac{d^2 r}{d\tau^2} + 3\frac{dr}{d\tau} - 2r^3 + 2r = 0. \quad (14.378)$$

This equation can be solved in general ([112])

$$r(\tau) = iC_1 e^\tau \operatorname{sn}(C_1 e^\tau + C_2, -1). \quad (14.379)$$

Thus

$$z(\tau) = -\sqrt{5}C_1 e^\tau \operatorname{sn}(C_1 e^\tau + C_2, -1). \quad (14.380)$$

$\operatorname{sn}(u, -1)$ is a Jacobi elliptic function with a modulus $K^2 = -1$, C_1 and C_2 are constants. In this way

$$\Phi(t) = \frac{1}{2\alpha_s} \left(1 - \sqrt{5}C_1 e^{H_0 t} \operatorname{sn}(C_1 e^{H_0 t} + C_2, -1) \right). \quad (14.381)$$

However, we are forced to put $\bar{a} = -\frac{4}{15}$. Let us remind that

$$\bar{a} = \frac{e^{n\Psi_1}}{2r^2 H_0^2} A \quad (14.382)$$

and we are supposing $A > 0$. However (see Eq. (14.321) for a definition of A), it is possible to consider $A < 0$ only for a pleasure to play. Thus we use formula (14.381) to consider a dynamics of a Higgs field. Let us start an inflation for $t = 0$ with $\Phi = 0$.

$$\Phi(0) = 0. \quad (14.383)$$

One finds

$$\frac{1}{C_1 \sqrt{5}} = \operatorname{sn}(C_1 + C_2, -1) \quad (14.384)$$

(i.e. $t_{\text{initial}} = 0$).

Let

$$\Phi(t_{\text{end}}) = \frac{1}{\alpha_s} \left(\frac{1 + \sqrt{5}}{2} \right) \quad (14.385)$$

(a true minimum—a “true” vacuum). One gets

$$\frac{1}{2} = -C_1 e^{H_0 t_{\text{end}}} \operatorname{sn} (C_1 e^{H_0 t_{\text{end}}} + C_2, -1). \quad (14.386)$$

Let us calculate the derivative of $\Phi(t)$. One gets

$$\begin{aligned} \frac{d\Phi}{dt} = & -\frac{\sqrt{5}}{2\alpha_s} C_1 H_0 e^{H_0 t} \left(\operatorname{sn} (C_1 e^{H_0 t} + C_2, -1) \right. \\ & \left. + \operatorname{cn} (C_1 e^{H_0 t} + C_2, -1) \cdot \operatorname{dn} (C_1 e^{H_0 t} + C_2, -1) \right). \end{aligned} \quad (14.387)$$

Let us suppose a slow movement of Φ such that

$$\frac{d\Phi}{dt}(0) = 0. \quad (14.388)$$

From (14.388) one gets

$$\operatorname{sc}(C_1 + C_2, -1) = -\operatorname{dn}(C_1 + C_2, -1). \quad (14.389)$$

This gives us an equation for a sum of integration constants. Thus we can get from Eq. (14.389) and Eq. (14.384) integration constants C_1 and C_2 and from Eq. (14.386) t_{end} , which can give us an amount of inflation

$$\bar{N}_0 = H_0 t_{\text{end}}. \quad (14.390)$$

In some sense Eq. (14.386) is an equation for an amount of inflation

$$\frac{1}{2} = -C_1 e^{\bar{N}_0} \operatorname{sn} (C_1 e^{\bar{N}_0} + C_2, -1). \quad (14.391)$$

Using Eq. (14.389) and Eq. (14.384) one easily gets

$$C_1 = -\sqrt{\frac{5 - \sqrt{5}}{10}} \simeq -0.5256. \quad (14.392)$$

One gets

$$C_1 + C_2 = \frac{1}{\sqrt{2}} (F(90^\circ \setminus 45^\circ) - F(32^\circ \setminus 45^\circ)) = 0.9431 \quad (14.393)$$

where F is an elliptic integral of the first kind and

$$\arcsin \left(\sqrt{\frac{5 - \sqrt{5}}{10}} \right) \simeq \arcsin(0.5256) \simeq 32^\circ. \quad (14.394)$$

Let us come back to Eq. (14.391) and let us denote

$$e^{\bar{N}_0} = x_0. \quad (14.395)$$

One gets using (14.393) and (14.392)

$$C_2 \simeq 1.4687 \quad (14.396)$$

and finally

$$1.4866x_0 - 5.0354 = F(\varphi/45^\circ), \quad (14.397)$$

where

$$\cos \varphi = \frac{1.9026}{x_0}. \quad (14.398)$$

From Eq. (14.397) one finds

$$\frac{2.8284}{\cos \varphi} - 5.0354 = \int_0^\varphi \frac{d\theta}{\sqrt{1 - \frac{1}{2} \sin^2 \theta}}. \quad (14.399)$$

We come back to this equation later.

Thus we get the following results. The Higgs field evolves from a “false” vacuum value ($\Phi = 0$) to the “true” vacuum value ($\Phi = \Phi_0$) completing a second order phase transition. The potential of selfinteracting Ψ field changes and a new equilibrium Ψ_0 is attained. However, the field Ψ evolves from Ψ_0 to new value a little different from Ψ_0 and afterwards a radiation era starts. The change of the value of a scalar field Ψ is small in terms of a variable y (see Eq. (14.94) for a definition) only from $\sqrt{\frac{n}{n+2}}$ to 1 (n is a natural number $n = \dim H$, and H is at least $G2$, $\dim G2 = 14$). Let us consider an evolution of a field Ψ in this epoch. From Eq. (14.84) we get

$$\ddot{\varphi} + \omega_0^2 \varphi = 0 \quad (14.400)$$

where

$$\omega_0^2 = \frac{M_{\text{pl}}^2}{2M}(n+2) \left(\frac{n}{n+2} \right)^{\frac{n}{2}} |\gamma| \left(\frac{|\gamma|}{\beta} \right)^{\frac{n}{2}} \quad (14.401)$$

$$\Psi = \Psi_0 + \varphi \quad (14.402)$$

and we linearize Eq. (14.84) around Ψ_0 neglecting a term with a first derivative of φ . Thus a scalar field Ψ undergoes small oscillations around the equilibrium Ψ_0 . These oscillations cannot be too long for a field should change from

$$\Psi_0 = \ln \left(\sqrt{\frac{n|\gamma|}{(n+2)\beta}} \right) \quad (14.403)$$

to

$$\bar{\Psi} = \ln \left(\sqrt{\frac{|\gamma|}{\beta}} \right). \quad (14.404)$$

In terms of a field φ from $\varphi = 0$ to $\varphi = \frac{1}{2} \ln(1 + \frac{2}{n}) \simeq \frac{1}{n}$. Thus we have

$$\varphi = \varphi_0 \sin(\omega_0 t). \quad (14.405)$$

For $t = 0$, $\varphi = 0$ the time Δt to go from $\varphi = 0$ to $\varphi = \frac{1}{n}$ is simply equal to

$$\Delta t = \frac{\left| \arcsin\left(\frac{1}{\varphi_0 n}\right) \right|}{\omega_0} \leq \frac{\pi}{2\omega_0}. \quad (14.406)$$

Thus the full amount of an inflation in this epoch is equal to

$$N_1 = \Delta t H_1 = \frac{H_1}{\omega_0} \left| \arcsin\left(\frac{1}{\varphi_0 n}\right) \right| \leq \frac{\pi H_1}{2\omega_0} = \frac{\pi M_{\text{pl}}}{2\sqrt{6\bar{M}}(n+2)}. \quad (14.407)$$

In our notation

$$t_1 = t_r + \Delta t \quad (14.408)$$

(see Eq. (14.114)).

After a time t_1 the Universe goes to the phase of radiation domination with a strong interaction between a radiation and a scalar field Ψ up to a minimal temperature where an ordinary (a dust) matter appears. This matter evolves afterwards independently of a radiation and of a scalar field Ψ . The scalar field now is evolving as a quintessence.

In order to get some comparison let us consider an evolution of a scalar field Ψ during the second de Sitter phase in a slow-roll approximation. Using Eq. (14.84) one gets

$$\frac{dy}{dt} = \frac{\bar{B} y^{\frac{(n-2)}{2}} (y^2 - (\frac{n}{n+2}))}{\sqrt{1-y^2}} \quad (14.409)$$

or

$$\int \frac{\sqrt{1-y^2} dy}{y^{\frac{(n-2)}{2}} (y^2 - (\frac{n}{n+2}))} = \bar{B}(t - t_0) \quad (14.410)$$

where

$$\bar{B} = 2(n+2)|\gamma| \left(\frac{n}{n+2} \right)^{\frac{n}{2}} \left(\frac{|\gamma|}{\beta} \right)^{\frac{n}{2}} \quad (14.411)$$

and

$$y = \sqrt{\frac{\beta}{|\gamma|}} \cdot e^{\Psi}. \quad (14.412)$$

Let

$$I = \int \frac{\sqrt{1-y^2} dy}{y^{(\frac{n-2}{2})(y^2 - (\frac{n}{n+2}))}} \simeq \int \frac{\sqrt{1-y^2} dy}{(y^2 - (\frac{n}{n+2}))} \quad (14.413)$$

for

$$\frac{n}{n+2} \leq y^2 \leq 1 \quad (14.414)$$

$$n > \dim G2 = 14. \quad (14.415)$$

For the integral I one finds:

$$\begin{aligned} I = & -\arcsin(y) + \frac{1}{\sqrt{n(n+2)}} \sqrt{1-y^2} \\ & + \left(\sqrt{\frac{2}{n}} + \sqrt{\frac{n}{2}} \right) \frac{1}{n+2} \cdot \ln \left[\frac{2 \left[\left(1 + y \sqrt{\frac{n}{n+2}} \right) \left(y + \sqrt{\frac{n}{n+2}} \right) + \sqrt{\frac{2}{n+2}} \sqrt{1-y^2} \right]}{\left(y + \sqrt{\frac{n}{n+2}} \right)} \right] \\ & - \frac{1}{n+2} \left(\sqrt{\frac{2}{n}} + \sqrt{\frac{n}{2}} \right) \cdot \ln \left[\frac{2 \left[\left(1 - y \sqrt{\frac{n}{n+2}} \right) \left(y - \sqrt{\frac{n}{n+2}} \right) + \sqrt{\frac{2}{n+2}} \sqrt{1-y^2} \right]}{\left(y - \sqrt{\frac{n}{n+2}} \right)} \right]. \end{aligned} \quad (14.416)$$

In order to find an amount of an inflation let us find limits of I for $y = 1$ and $y = \sqrt{\frac{n}{n+2}}$. One finds

$$\begin{aligned} \lim_{y \rightarrow 1} I = & -\frac{\pi}{2} + \left(\sqrt{\frac{2}{n}} + \sqrt{\frac{n}{2}} \right) \frac{\ln \left(2 \left(1 + \sqrt{\frac{n}{n+2}} \right) \right)}{(n+2)} \\ & - \left(\sqrt{\frac{2}{n}} + \sqrt{\frac{n}{2}} \right) \frac{\ln \left(2 \left(1 - \sqrt{\frac{n}{n+2}} \right) \right)}{(n+2)} \end{aligned} \quad (14.417)$$

$$\lim_{y \rightarrow \sqrt{\frac{n}{n+2}}} I = -\infty. \quad (14.418)$$

Thus we see that a slow-roll approximation offers an infinite in time evolution of a field y from $\sqrt{\frac{n}{n+2}}$ to 1. It is similar to an evolution of a Higgs field in some slow-roll approximation schemes. The inflation never ends. Let us come back to the Eq. (14.135) and let us calculate a Hubble constant and a deceleration parameter for this model

$$H = \frac{\dot{R}}{R} = \left(\frac{2|\gamma|^{n+4}}{3\bar{M}\beta^n(n-1)} \right) \frac{(t-t_1)^3}{\left(1 + \frac{|\gamma|^{n+4}}{3\bar{M}\beta^n(n-1)}(t-t_1)^4 \right)} \quad (14.419)$$

$$q = -\frac{\ddot{R}R}{\dot{R}^2} = -\frac{2}{3} \left(\frac{1}{(t-t_1)^4} + \frac{|\gamma|^{n+4}}{9\bar{M}\beta^n} \right). \quad (14.420)$$

Let us compare Eq. (14.419) with the similar equation for radiation dominated Universe in General Relativity

$$H_{\text{GR}} = \frac{2}{3(t - t_1)} \quad (14.421)$$

and let us calculate a speed up factor

$$\Delta_1 = \frac{H}{H_{\text{GR}}} = \frac{|\gamma|^{n+4}}{\overline{M}\beta^n(n-1)} \frac{(t - t_1)^4}{\left(1 + \frac{|\gamma|^{n+4}}{3\overline{M}\beta^n(n-1)}(t - t_1)^4\right)} \quad (14.422)$$

and let us do the same for a model Eq. (14.215) (i.e. radiation dominated with a quintessence). Using Eq. (14.216) one gets

$$\Delta_2 = \frac{H}{H_{\text{GR}}} = \frac{3\sqrt{3}m_{\text{pl}}\rho_Q(t - t_1)}{2B \operatorname{tgh}\left(\frac{2\sqrt{3}\rho_Q}{B}m_{\text{pl}}(t - t_1)\right)}. \quad (14.423)$$

Let us notice that for a $(t - t_1) \sim 0$

$$\Delta_1 \simeq 0 \quad (14.424)$$

and

$$\lim_{t \rightarrow \infty} \Delta_1 = +\infty. \quad (14.425)$$

Thus at early stages model Eq. (14.135) is slower than that in General Relativity. For large time the behaviour depends on details of the theory. In the case of model Eq. (14.215) we have

$$\lim_{t \rightarrow t_1} \Delta_2 = \frac{3}{4} \quad (14.426)$$

and

$$\lim_{t \rightarrow \infty} \Delta_2 = +\infty. \quad (14.427)$$

Thus in the first case Δ is monotonically going from 0 to $+\infty$ and in the second case from $\frac{3}{4}$ to infinity. According to modern ideas an expansion rate in early Universe has an important influence on a production of light elements, the so called primordial abundance of light elements (see Ref. [113]). If at a beginning of primordial nucleosynthesis the Universe expansion rate is slower than in GR, then we have ^4He underproduction. This can be balanced by considering a larger ratio of a number density of barions to number density of photons (remember the barion number is a conserved quantity)

$$\eta = \frac{n_B}{n_\gamma} \quad (14.428)$$

where n_B, n_γ —barion and photon density numbers for a high redshift $z = 10^{10}$. In contrast to the latest if an expansion rate became faster than in GR during nucleosynthesis

process, those bigger η (traditionally one uses the so called) η_{10}

$$\eta_{10} = 10^{10} \eta \quad (14.429)$$

do not result in excessive burning of deuterium because this happens in a shorter time. The standard model of big-bang nucleosynthesis (SBBN) demands

$$3 \leq \eta_{10} \leq 5.6. \quad (14.430)$$

It seems in the light of observational data from cosmic microwave background (CMB) (BOOMERANG and MAXIMA) and from the Lyman α -forest that $\eta_{10} = 8.8 \pm 1.4$, significantly higher than the SBBM value (for CMB) and 8.2 ± 2 or 12.3 ± 2 for Lyman α -forest. Thus the second model could help in principle to explain the data without modification of nuclear reaction rates due to neutrino degeneracy or introducing new decaying particles. As the Universe expands, cools and becomes more dilute, the nuclear reactions cease to create and destroy nuclei. The abundances of the light nuclei formed during this epoch are determined by the competition between a time available (an expansion rate) and a density of reactans: neutrons and protons.

The abundances of D, ^3He , ^7Li are limited by an expansion rate and are determined by the competition between the nuclear production/destruction rates and an (universal) expansion of the Universe. The SBBN theory is based on a flat radiation dominated Universe model in General Relativity and found in a laboratory nuclear reaction rates. Thus it is strongly constrained. Any significant discrepancy between observed and calculated value of η_{10} (known as baryometry) could be very dangerous. Thus changing the ratio of an universal expansion relative to the model of General Relativity can give some margin in a primordial alchemy.

In our theory after two inflationary phases we have in principle two radiation dominated phases described by Eq. (14.135) and Eq. (14.215) with speed up factors (14.422) and (14.423). However, the important point is to match those models. We get

$$\begin{aligned} R(t_d) &= R_0 \exp(H_0 t_r + H_1(t_1 - t_r)) \left(1 + \frac{\bar{r}}{(n-1)} \eta_{\min}^2\right)^{\frac{1}{2}} \\ &= \sqrt[4]{\frac{B}{\rho_Q}} \left(\sinh \left(\frac{2\sqrt{3}\sqrt{\rho_Q}}{B} m_{\text{pl}} \left(t_1 - \bar{t}_0 + \frac{\sqrt{2M}\beta^{\frac{n}{4}}}{|\gamma|^{\frac{n+4}{4}}} \eta_{\min}^{\frac{1}{2}} \right) \right) \right)^{\frac{1}{2}} \end{aligned} \quad (14.431)$$

and

$$\begin{aligned} B &= \rho_r(t_d) R^3(t_d) = \rho_r \left(t_1 + \frac{\sqrt{2M}\beta^{\frac{n}{4}}}{|\gamma|^{\frac{n+4}{4}}} \eta_{\min}^{\frac{1}{2}} \right) R^3 \left(t_1 + \frac{\sqrt{2M}\beta^{\frac{n}{4}}}{|\gamma|^{\frac{n+4}{4}}} \eta_{\min}^{\frac{1}{2}} \right) \\ &= \rho_{\min} R_0^3 \exp(3H_0 t_r + 3H_1(t_1 - t_r)) \left(1 + \frac{\bar{r}}{(n+1)} \eta_{\min}^2\right)^{\frac{3}{2}} \end{aligned} \quad (14.432)$$

where $\rho_{\min}$ is given by

$$\rho_{\min} = M_{\text{pl}}^2 \frac{\beta^{n+1}}{|\gamma|^n} (1 + \eta_{\min})^{2(n+1)} W_4(1 + \eta_{\min}), \quad (14.433)$$

$W_4(y)$ is given by formula (14.115) and $\eta_{\min}$ by formula (14.122).

The time $t_r = t_{\text{end}}$ in any inflation model for an evolution of the Higgs field in the first de Sitter phase. The time $t_1 = t_r + \Delta t$ can be calculated from Eq. (14.406)

$$\Delta t = \frac{\arcsin(\frac{1}{\varphi_0 n})}{\omega_0} \quad (14.434)$$

and is bounded by

$$\Delta t \leq \frac{1}{\omega_0} = \frac{1}{M_{\text{pl}} |\gamma|^{\frac{1}{2}}} \sqrt{\frac{2\overline{M}}{(n+2)}} \left(\frac{n+2}{n} \right)^{\frac{n}{4}} \left(\frac{\beta}{|\gamma|} \right)^{\frac{n}{4}}. \quad (14.435)$$

From Eq. (14.431) we can get the constant $\bar{t}_0$

$$\bar{t}_0 = t_1 + \frac{\sqrt{2\overline{M}} \beta^{\frac{n}{4}}}{|\gamma|^{\frac{n+4}{4}}} \eta_{\min}^{\frac{1}{2}} - T \quad (14.436)$$

where

$$T = \frac{B \operatorname{arsinh} \left[R_0^2 \sqrt{\frac{\rho_Q}{B}} \exp(2H_0 t_r + 2H_1(t_1 - t_r)) \left(1 + \frac{\bar{r}}{(n-1)} \eta_{\min}^2 \right) \right]}{2\sqrt{3} \sqrt{\rho_Q} M_{\text{pl}}} \quad (14.437)$$

and B is given by Eq. (14.432).

Let us notice that for $t = t_d$ the Universe undergoes a phase transition due to a change in an equation of state for a scalar field Ψ from $p_\Psi = \frac{4}{3}\rho_\Psi$ in a first radiation dominated phase to $p_\Psi = -\rho_\Psi$ in a second radiation dominated phase (a quintessence phase).

A matter which appears in the second phase does not play any rôle in an evolution of the Universe. For $t = t_d$ we have a discontinuity in Hubble constants (parameters) for both phases

$$H(t_d) = \frac{2|\gamma|^{n+4}}{3\overline{M}(n-1)\beta^n} \cdot \frac{(t_d - t_1)^{\frac{3}{2}}}{\left(1 + \frac{\bar{r}}{(n-1)} \eta_{\min}^2 \right)} \quad (14.438)$$

in the first phase and

$$\overline{H}(t_d) = \sqrt{3} m_{\text{pl}} \left(\frac{\sqrt{\rho_Q}}{B} \right) \operatorname{ctgh} \left(\frac{2\sqrt{3}}{B} \sqrt{\rho_Q} m_{\text{pl}} T \right) \quad (14.439)$$

for the second one,

$$H(t_d) \neq \overline{H}(t_d) \quad (14.440)$$

$$\rho_Q = \frac{1}{2}\lambda_{c0}(\Psi_0) = \frac{|\gamma|^{\frac{n}{2}+1}n^{\frac{n}{2}}}{\beta^{\frac{n}{2}}(n+2)^{\frac{n}{2}+1}}. \quad (14.441)$$

Let us notice that a speed up factor Δ_1 changes from

$$\Delta_1(t_1) = 0 \quad (14.442)$$

to

$$\Delta_1(t_d) = \frac{|\gamma|^{n+4}}{\overline{M}(n-1)\beta^n} \cdot \frac{(t_d - t_1)^4}{\left(1 + \frac{\bar{r}}{(n-1)}\eta_{\min}^2\right)} \quad (14.443)$$

and an expansion rate can be slower than in General Relativity.

The speed up factor Δ_2 has a more complicated behaviour than Δ_1 . It is natural to expect a rapid speed up of a rate of expansion resulting in non-equilibrium nuclear synthesis of light elements. Thus the presented scenario offers interesting possibilities for an evolution of early Universe. Recently some papers have appeared on scalar-tensor theories of gravitation exploiting the idea of a speed up factor in primordial nucleosynthesis [114].

Finally let us come back to the model (14.221), (14.224). We cannot invert analytically the formula (14.224). Moreover taking under consideration Eqs (14.242–245) and making some natural simplifications we come to the following approximative formulae for $R(t)$.

For $x < 1.1969$ (Eq. (14.243))

$$R(t) = \sqrt[3]{\frac{A}{\rho_Q}} (0.44 + 0.085N) \quad (14.444)$$

where

$$N = \exp\left(0.374\frac{\sqrt{\rho_Q}}{M_{\text{pl}}}(t - t_0) + 157.93\right). \quad (14.445)$$

One can calculate a Hubble parameter and a deceleration parameter and gets:

$$H = \frac{\dot{R}}{R} = \frac{\sqrt{\rho_Q}}{M_{\text{pl}}} \frac{3.17N}{44 + 8.5N} \quad (14.446)$$

$$-q = \frac{\ddot{R}R}{\dot{R}^2} = \frac{5.21}{N} + 1. \quad (14.447)$$

We have $H > 0$, and $q < 0$. Thus the model expands and accelerates. First of all we consider

$$\rho_m = \frac{A}{R^3} e^{aQ}$$

and we take boundary for R . In this way one gets

$$\rho_m = a_i \cdot \rho_Q e^{aQ}, \quad i = 1, 2, 3, 4, \quad (14.448)$$

$$\begin{aligned}
a_1 &= 2.015 \\
a_2 &= 0.034 \\
a_3 &= 1060 \\
a_4 &= 8
\end{aligned} \tag{14.449}$$

If we reconsider a contemporary gravitational constant G_N as $G_N e^{-(n+2)\Psi_0}$ we would get interesting ratios. Thus we get

$$\frac{\rho_m}{\rho_Q} = 0.034 \div 2.015 \quad \text{or} \quad 8 \div 1060 \tag{14.450}$$

which for the first is an excellent agreement with recent data concerning an acceleration of the Universe. Solving Eq. (14.445) for N if $R = 1.33\sqrt{\frac{A}{\rho_Q}}$ gives us

$$N = 10.47 \tag{14.451}$$

$$H = \frac{\sqrt{\rho_Q}}{M_{\text{pl}}} \cdot 0.25 \tag{14.452}$$

$$-q = 1.5. \tag{14.453}$$

Using Eq. (14.445) and Eq. (14.452) we can estimate a time of our contemporary epoch

$$t = -\frac{1}{h} \cdot 10^{15} \text{yr} - t_0 \tag{14.454}$$

where h is a dimensionless Hubble parameter, $H = h \cdot H_0$ (we take for a contemporary Hubble parameter $H_0 = 100 \frac{\text{km/s}}{\text{Mps}}$), $0.7 < h < 1$, which seems to be too much.

We can also estimate a density of a quintessence

$$\rho_Q = \frac{H^2}{G_N} \cdot 0.571 = h^2 \cdot 0.08 \cdot 10^{-23} \frac{\text{kg}}{\text{m}^3} = 8h^2 \cdot 10^{-29} \frac{\text{g}}{\text{cm}^3}. \tag{14.455}$$

Let us come back to the Eqs (14.334) and (14.341). In Eq. (14.341) a Higgs field is given by a P Weierstrass function. This function can be expressed by Jacobi elliptic function

$$P(z) = e_3 + (e_1 - e_2) \text{ns}^2 \left(z(e_1 - e_2)^{\frac{1}{2}} \right) \tag{14.456}$$

with

$$K^2 = \frac{e_2 - e_3}{e_1 - e_3} \tag{14.457}$$

where e_1, e_2, e_3 are roots of the polynomial

$$4x^3 - g_2x - g_3. \tag{14.458}$$

Thus we should solve a cubic equation

$$x^3 - \frac{1}{12} (5 - 3a^2) - \frac{1}{12} \left(\frac{a^2}{4} + \frac{10}{9} \right) = 0. \tag{14.459}$$

We want Eq. (14.459) to have all real roots. Thus we need

$$p = -\frac{1}{12} (5 - 3a^2) < 0 \quad (14.460)$$

and

$$D = \frac{p^3}{27} + \frac{q^2}{4} < 0 \quad (14.461)$$

where

$$q = -\frac{1}{12} \left(\frac{a^2}{4} + \frac{10}{9} \right). \quad (14.462)$$

Both conditions (14.460) and (14.461) are satisfied if

$$0 < a < 0.3115. \quad (14.463)$$

In this case we get

$$e_1 = \frac{1}{3} \sqrt{5 - 3a^2} \cos \frac{\varphi}{3} \quad (14.464)$$

$$e_2 = \frac{1}{3} \sqrt{5 - 3a^2} \cos \frac{\varphi + 2\pi}{3} \quad (14.465)$$

$$e_3 = \frac{1}{3} \sqrt{5 - 3a^2} \cos \frac{\varphi + 4\pi}{3} \quad (14.466)$$

where

$$\cos \varphi = -\frac{(9a^2 + 40)}{12(5 - 3a^2)^{3/2}}. \quad (14.467)$$

From (14.464–466) one gets

$$K^2 = \frac{\sin \frac{\varphi}{3}}{\sin \frac{\varphi + 2\pi}{3}} \quad (14.468)$$

$$e_1 > e_2 > e_3 \quad (14.469)$$

Thus

$$\Phi(t) = \frac{(5 + 3a)\sqrt{5 + 4a}}{8\alpha_s(A_1 \operatorname{ns}^2(u) + A_2)} + \frac{(1 + \sqrt{5 + 4a})}{2\alpha_s} \quad (14.470)$$

where

$$u = \sqrt[4]{\frac{Ae^{n\Psi_1}(5Ae^{n\Psi_1} - 3\alpha_s^2 r^2)}{r^4}} \sqrt{\sin \frac{\varphi + \pi}{3}} (t - t_0) \quad (14.471)$$

$$A_1 = e_1 - e_2 = \sqrt{\frac{5}{3} - a^2} \sin \frac{\varphi + \pi}{3} \quad (14.472)$$

$$A_2 = e_3 + \frac{1}{12}(5 + 6a) = \sqrt{\frac{5}{3} - a^2} \cos \frac{\varphi + 4\pi}{3} + \frac{1}{12}(5 + 6a) \quad (14.473)$$

and

$$\cos \varphi = -\frac{\sqrt{A}e^{\frac{n}{2}\Psi_1} (9\alpha_s^2 Br^2 + 40Ae^{n\Psi_1})}{12 (5Ae^{n\Psi_1} - 3\alpha_s^2 Br^2)^{3/2}} \quad (14.474)$$

$$0 < \sqrt{\frac{\alpha_s^2 Br^2}{Ae^{n\Psi_1}}} < 0.3115 \quad (14.475)$$

and a is given by Eq. (14.330).

The function $\text{ns}(u, K)$ is an elliptic Jacobi function with a modulus K given by Eq. (14.468)

$$\text{ns}(u, K) = \frac{1}{\text{sn}(u, K)} \quad (14.476)$$

$$\text{sn}(u, K) = \text{sinam}(u, K). \quad (14.477)$$

In order to justify our treatment of Eqs (14.287) and (14.290) we consider the full field equations

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = 2\bar{\Lambda}u_\mu u_\nu - \bar{\Lambda}g_{\mu\nu} \quad (14.478)$$

(where $\bar{\Lambda}$ is given by Eq. (14.293)).

However, in this case we consider $\bar{\Lambda}$ arbitrary. Let us consider static and spherically symmetric metric

$$ds^2 = e^v dt^2 - e^\lambda dr^2 - r^2 (d\theta^2 + \sin^2 \theta d\varphi^2) \quad (14.479)$$

where $v = v(r)$ and $\lambda = \lambda(r)$.

From Eqs (14.478–479) one gets

$$r^2 \bar{\Lambda} = e^{-\lambda}(rv' + 1) - 1 \quad (14.480)$$

$$r^2 \bar{\Lambda} = e^{-\lambda}(r\lambda' - 1) + 1 \quad (14.481)$$

$$\bar{\Lambda}' = -\bar{\Lambda}v' \quad (14.482)$$

where $'$ means derivation with respect to r . Summing up (14.480) and (14.481) one gets

$$e^{-\lambda}(\lambda' + v') = 2\bar{\Lambda}r. \quad (14.483)$$

Moreover in our case $\bar{\Lambda}$ is very small. Thus we get approximately

$$\frac{d\bar{\Lambda}}{dr} \simeq 0 \quad (14.484)$$

and

$$\lambda' \cong -v'. \quad (14.485)$$

In this way we go to the solution (14.294) for a small $\overline{\lambda}$ which is our case. In this way $\lambda = -v$ and

$$B(r) = e^v = e^{-\lambda}. \quad (14.486)$$

Let us check a consistency of our solution. In order to do this we consider Eqs (14.480–482) in full. One gets from Eq. (14.482)

$$\overline{\lambda} = e^{-\mu_0} e^{-v}. \quad (14.487)$$

For $\overline{\lambda}$ is very small we should suppose that μ_0 is positive and very large ($\mu_0 \rightarrow \infty$). From Eqs (14.480–481) one gets

$$\frac{d}{dr}(\lambda + v) = 2e^{-\mu_0} \cdot r \cdot e^{(\lambda-v)}. \quad (14.488)$$

Supposing that

$$\lambda(r_0) + v(r_0) = 0 \quad (14.489)$$

for an established $r_0 > 0$ (r_0 is greater than a Schwarzschild radius of the mass M_0 from the solution (14.294)). Taking sufficiently big μ_0 we get approximately

$$\frac{d}{dr}(\lambda + v) \simeq 0 \quad (14.490)$$

and of course

$$\lambda \simeq -v. \quad (14.491)$$

Let us come back to the Eq. (14.399) and consider it for

$$\varphi = k\pi + \frac{\pi}{2} - \varepsilon, \quad k = 0, \pm 1, \pm 2, \dots, \quad (14.492)$$

where ε is considered to be small. One gets

$$\frac{2.8284(-1)^k}{\sin \varepsilon} - 5.0354 = \int_0^{\frac{\pi}{2}-\varepsilon} \frac{d\theta}{\sqrt{1 - \frac{1}{2} \sin^2 \theta}} + 2kK\left(\frac{1}{2}\right), \quad (14.493)$$

where

$$K\left(\frac{1}{2}\right) = \int_0^{\frac{\pi}{2}} \frac{d\theta}{\sqrt{1 - \frac{1}{2} \sin^2 \theta}} = 1.8541 \quad (14.494)$$

is a full elliptic integral of the first kind for a modulus equal $\frac{1}{2}$. For a small ε we can write

$$\sin \varepsilon \simeq \varepsilon \quad (14.495)$$

$$\int_0^{\frac{\pi}{2}-\varepsilon} \frac{d\theta}{\sqrt{1 - \frac{1}{2} \sin^2 \theta}} \simeq K\left(\frac{1}{2}\right) - \sqrt{2}\varepsilon. \quad (14.496)$$

Thus one gets for $k = 2l$

$$\frac{2.8284}{\varepsilon} - 5.0354 = -\sqrt{2}\varepsilon + (4l + 1)1.8541. \quad (14.497)$$

It is easy to notice that we can realize a 60-fold inflation in our model for

$$x_0 = \frac{1.9026}{|\varepsilon|} \quad (14.497a)$$

can be arbitrary big for sufficiently big l .

Let us take $l = 25 \cdot 10^{24}$. In this case we have for ε an equation

$$\varepsilon^2 - ((4l + 1)1.3110 + 5.0354)\varepsilon + 2 = 0 \quad (14.498)$$

or

$$\varepsilon^2 - (10^{26} + 6.0354)\varepsilon + 2 = 0, \quad (14.498a)$$

i.e.

$$\varepsilon^2 - 10^{26}\varepsilon + 2 = 0, \quad (14.498b)$$

and finally

$$|\varepsilon| \simeq 10^{-26} \quad (14.499)$$

$$x_0 \simeq 1.9026 \cdot 10^{26} \quad (14.500)$$

$$\ln x_0 \simeq 60.51 \quad (14.501)$$

which gives us a 60-fold inflation.

Moreover the Eq. (14.399) has an infinite number of solutions for $0 \leq \psi \leq \frac{\pi}{2}$, $\varphi = k\pi + \psi$,

$$\frac{2.8284}{\cos \psi} - 5.0354 = \int_0^\psi \frac{d\theta}{\sqrt{1 - \frac{1}{2}\sin^2 \theta}} + 7.5164 \cdot l \quad (14.502)$$

with

$$x_0 = \frac{1.9026}{\cos \psi}, \quad l = 0, 1, 2, 3, \dots, \quad k = 2l. \quad (14.503)$$

One can find roots of Eq. (14.502) for

$l = 5$

$$\begin{aligned} x_0 &= 16.83 \\ \ln x_0 &= 2.82, \end{aligned} \quad (14.504a)$$

$l = 50$

$$\begin{aligned} x_0 &= 0.24 \cdot 10^3 \\ \ln x_0 &= 5.48 \end{aligned} \quad (14.504b)$$

and $\ln x_0$ for $l = 5 \cdot 10^{24}$. In the last case one finds using 70-digit arithmetics

$$\ln x_0 \simeq 58.479 \dots \quad (14.505)$$

which gives us an almost 60-fold inflation. In general an amount of inflation

$$\overline{N}_0 = \ln x_0 \quad (14.506)$$

is a function of $l = 0, 1, 2, \dots$ and probably can be connected to the Dirac large number hypothesis.

Finally let us take $l = 10^n$ where $n > 10$. In this case one can solve Eq. (14.498a) and get

$$|\varepsilon| \simeq \frac{2}{[(4 \cdot 10^n + 1)1.3110 + 5.0354]}. \quad (14.507)$$

Thus for x_0 we find

$$x_0 \simeq 5 \cdot 10^n \quad (14.508)$$

and

$$\overline{N}_0(n) \simeq 1.60 + 2.30n. \quad (14.509)$$

In this way, for large l , $\overline{N}_0$ is a linear function of a logarithm of l ,

$$\begin{aligned} \overline{N}_0(10) &\simeq 24.6, \quad \overline{N}_0(20) \simeq 47.6, \quad \overline{N}_0(24) \simeq 56.8, \\ \overline{N}_0(25) &\simeq 59.1, \quad \overline{N}_0(26) \simeq 61.4 \end{aligned} \quad (14.510)$$

or

$$\overline{N}_0(\log_{10} l) = 1.6 + \ln l. \quad (14.511)$$

It is easy to see that Eq. (14.511) is an excellent approximation even for $l = 50$.

Let us come back to the Eq. (14.186) in order to find a power spectrum for our simplified model of inflation. Using Eqs (14.318), (14.387), (14.392) and (14.396) one gets after some algebra

$$P_R(K) \cong 2.8944 \left(\frac{H_0}{2\pi} \right)^2 n_1 \alpha_s^2 f \left(\frac{K}{R_0 H_0} \right) \quad (14.512)$$

where

$$\begin{aligned} f(x) = x^{-2} &\left(\operatorname{sn}(1.4687 - 0.5256x, -1) \right. \\ &\left. + \operatorname{cn}(1.4687 - 0.5256x, -1) \operatorname{dn}(1.4687 - 0.5256x, -1) \right)^{-2}. \end{aligned} \quad (14.513)$$

Using some relations among elliptic functions one gets

$$f(x) = \frac{1}{x^2} \cdot \frac{2 \operatorname{dn}^4(u, \frac{1}{2})}{\left(\operatorname{sn}(u, \frac{1}{2}) \operatorname{dn}(u, \frac{1}{2}) + \sqrt{2} \operatorname{cn}(u, \frac{1}{2}) \right)^2} \quad (14.514)$$

where

$$u = 2.0770 - 0.7433x. \quad (14.515)$$

Let us consider a more general situation for the Eq. (14.388), i.e.

$$\frac{d\Phi}{dt}(0) = h \quad (14.516)$$

where $h \neq 0$. In this way one gets

$$h = -\frac{\sqrt{5}}{\alpha_s} C_1 H_0 (\text{sn}(C_1 + C_2, -1) + \text{cn}(C_1 + C_2, -1) \text{dn}(C_1 + C_2, -1)). \quad (14.517)$$

Using Eq. (14.384) one gets

$$C_2 = \frac{K(\frac{1}{2})}{\sqrt{2}} - \frac{\sqrt{10}}{\sqrt{\sqrt{(1 + \frac{\alpha_s h}{H_0})^4 + 4} - (1 + \frac{\alpha_s h}{H_0})^2}} - \frac{1}{\sqrt{2}} \int_0^{\arccos \sqrt{\frac{\sqrt{(1 + \frac{\alpha_s h}{H_0})^4 + 4} - (1 + \frac{\alpha_s h}{H_0})^2}}}{\sqrt{1 - \frac{1}{2} \sin^2 \varphi}} d\varphi \quad (14.518)$$

and

$$|C_1| = \frac{\sqrt{10}}{\sqrt{\sqrt{(1 + \frac{\alpha_s h}{H_0})^4 + 4} - (1 + \frac{\alpha_s h}{H_0})^2}}. \quad (14.519)$$

Let us come back to the Eq. (14.386) to find an amount of inflation for C_2 and C_1 given by Eqs (14.518–519). One gets

$$\sqrt{2}C_2 - K\left(\frac{1}{2}\right) = \int_0^{\arccos\left(\frac{1}{2C_1 e^{N_0}}\right)} \frac{d\theta}{\sqrt{1 - \frac{1}{2} \sin^2 \theta}} - \sqrt{2}C_1 e^{N_0}, \quad (14.520)$$

or using a natural substitution

$$\cos \varphi = \frac{1}{2C_1 e^{N_0}}, \quad (14.521)$$

$$\sqrt{2}C_2 - K\left(\frac{1}{2}\right) = \int_0^{\varphi} \frac{d\theta}{\sqrt{1 - \frac{1}{2} \sin^2 \theta}} - \frac{\sqrt{2}}{2 \cos \varphi}. \quad (14.522)$$

Taking as usual

$$\varphi = 2l\pi + \frac{\pi}{2} - \varepsilon, \quad l = 0, 1, 2, \dots, \quad (14.523)$$

one gets for small ε

$$\varepsilon \simeq \frac{1}{\sqrt{2}(4l+2)K(\frac{1}{2}) - 2C_2} \simeq \frac{1}{(4l+2)\sqrt{2}K(\frac{1}{2})} \quad (14.524)$$

(for large l) and eventually

$$\begin{aligned} e^{N_0} &\simeq \frac{4\sqrt{2}K(\frac{1}{2})}{C_1} l \\ N_0 &= \ln \left(\frac{4\sqrt{2}K(\frac{1}{2})}{C_1} \right) + \ln l. \end{aligned} \quad (14.525)$$

Let us calculate $\frac{d\Phi}{dt}$ for $t = t^* = \frac{1}{H_0} \ln \left(\frac{K}{R_0 H_0} \right)$. One gets

$$\begin{aligned} \frac{d\Phi}{dt}(t^*) &= -\frac{\sqrt{5}}{2\alpha_s} C_1 H_0 \left(\frac{K}{R_0 H_0} \right) \left(\operatorname{sn} \left(C_1 \left(\frac{K}{R_0 H_0} \right) + C_2, -1 \right) \right. \\ &\quad \left. + \operatorname{cn} \left(C_1 \left(\frac{K}{R_0 H_0} \right) + C_2, -1 \right) \operatorname{dn} \left(C_1 \left(\frac{K}{R_0 H_0} \right) + C_2, -1 \right) \right). \end{aligned} \quad (14.526)$$

Thus we can write down a $P_R(K)$ function.

$$P_R(K) = \left(\frac{H_0}{2\pi} \right)^2 \cdot \frac{4\alpha_s^2 n_1}{5C_1^2} f \left(\frac{K}{R_0 H_0} \right) \quad (14.527)$$

where

$$f(x) = \frac{1}{x^2} \left(\operatorname{sn}(C_1 x + C_2, -1) + \operatorname{cn}(C_1 x + C_2, -1) \operatorname{dn}(C_1 x + C_2, -1) \right)^{-2}. \quad (14.528)$$

Using some relation among elliptic functions one finds

$$P_R(K) = \left(\frac{H_0}{2\pi} \right)^2 \cdot \frac{4\alpha_s^2 n_1}{5C_1^2} g \left(\frac{K}{R_0 H_0} \right) \quad (14.529)$$

where

$$g(x) = \frac{1}{x^2} \cdot \frac{2}{\left(\operatorname{sd}(u, \frac{1}{2}) + \sqrt{2} \operatorname{cd}(u, \frac{1}{2}) \operatorname{nd}(u, \frac{1}{2}) \right)^2} \quad (14.530)$$

and

$$u = \sqrt{2}C_1 x + \sqrt{2}C_2. \quad (14.531)$$

Moreover we can reparametrize (14.530–531) in the following way

$$u = \sqrt{2}C_1 x + K(\frac{1}{2}) - C_1 \sqrt{2} - \int_0^{\arccos(\frac{\sqrt{5}}{C_1})} \frac{d\varphi}{\sqrt{1 - \frac{1}{2} \sin^2 \varphi}} \quad (14.532)$$

where

$$\frac{\alpha_s h}{H_0} = -1 \pm \frac{1}{|C_1|} \sqrt{\frac{C_1^4 - 25}{5}} \quad (14.533)$$

$$|C_1| > \sqrt{5}. \quad (14.534)$$

For large C_1 one gets

$$u \cong \sqrt{2}C_1(x-1) \quad (14.535)$$

$$\frac{\alpha_s h}{H_0} \simeq \frac{C_1}{\sqrt{5}}. \quad (14.536)$$

Using Eq. (14.525),

$$u = \frac{4\sqrt{2}K(\frac{1}{2})l}{e^{N_0}}(x-1). \quad (14.537)$$

If we take large C_1 (it means, a large h)

$$h = \frac{C_1 H_0}{\sqrt{5}\alpha_s} \quad (14.538)$$

and simultaneously sufficiently large l we can achieve a 60-fold inflation with a function $P_R(K)$ given by the formula (14.530). Large C_1 means here $C_1 \simeq 100$, large l means $l \simeq 10^{25}$.

Let us calculate the spectral index for our $P_R(K)$ function, i.e.

$$n_s(K) - 1 = \frac{d \ln P_R(K)}{d \ln K}. \quad (14.539)$$

One gets

$$n_s(K) - 1 = -2 - 2C_1 x \frac{\sqrt{2} \text{cd}(u, \frac{1}{2}) - \text{sd}^3(u, \frac{1}{2})}{\text{sd}(u, \frac{1}{2}) + \sqrt{2} \text{cd}(u, \frac{1}{2}) \text{nd}(u, \frac{1}{2})} \quad (14.540)$$

where

$$u = \sqrt{2}(C_1 x + C_2) = \frac{\sqrt{2}}{R_0 H_0} (C_1 K + C_2 R_0 H_0) \quad (14.541)$$

$$x = \frac{K}{R_0 H_0}$$

A very interesting characteristic of $P_R(K)$ is also $\frac{dn_s(K)}{d \ln K}$. One gets

$$\begin{aligned} \frac{dn_s(K)}{d \ln K} = & -2C_1 x \frac{\sqrt{2} \text{cd}(u, \frac{1}{2}) - \text{sd}^3(u, \frac{1}{2})}{\text{sd}(u, \frac{1}{2}) + \sqrt{2} \text{cd}(u, \frac{1}{2}) \text{nd}(u, \frac{1}{2})} \\ & - 2C_1^2 x^2 \text{sd}^2(u, \frac{1}{2}) - 2C_1^2 x^2 \frac{(\sqrt{2} \text{cd}(u, \frac{1}{2}) - \text{sd}^3(u, \frac{1}{2}))^2}{(\text{sd}(u, \frac{1}{2}) + \sqrt{2} \text{cd}(u, \frac{1}{2}) \text{nd}(u, \frac{1}{2}))^2} \end{aligned} \quad (14.542)$$

where u and x are given by Eq. (14.541).

Let us take for a trial $C_1 = -0.5256$ and $C_2 = 1.4687$. In this case one finds

$$n_s(K) - 1 = -2 + 1.0512x \frac{\sqrt{2} \text{cd}(u, \frac{1}{2}) - \text{sd}^3(u, \frac{1}{2})}{\text{sd}(u, \frac{1}{2}) + \sqrt{2} \text{cd}(u, \frac{1}{2}) \text{nd}(u, \frac{1}{2})} \quad (14.543)$$

$$\begin{aligned} \frac{dn_s(K)}{d \ln K} &= 1.0512x \frac{\sqrt{2} \text{cd}(u, \frac{1}{2}) - \text{sd}^3(u, \frac{1}{2})}{\text{sd}(u, \frac{1}{2}) + \sqrt{2} \text{cd}(u, \frac{1}{2}) \text{nd}(u, \frac{1}{2})} \\ &\quad - 0.5525x^2 \text{sd}^2(u, \frac{1}{2}) - 0.5525x^2 \frac{(\sqrt{2} \text{cd}(u, \frac{1}{2}) - \text{sd}^3(u, \frac{1}{2}))^2}{(\text{sd}(u, \frac{1}{2}) + \sqrt{2} \text{cd}(u, \frac{1}{2}) \text{nd}(u, \frac{1}{2}))^2} \end{aligned} \quad (14.544)$$

$$\begin{aligned} u &= 2.0771 - 0.7433x \\ x &= \frac{K}{R_0 H_0} \end{aligned} \quad (14.544a)$$

The interesting point is to find $n_s(K) \simeq 1$ (a flat power spectrum). One gets

$$\text{sd}(u, \frac{1}{2}) + \sqrt{2} \text{cd}(u, \frac{1}{2}) \text{nd}(u, \frac{1}{2}) = C_1 x \left(\text{sd}^3(u, \frac{1}{2}) - \sqrt{2} \text{cd}(u, \frac{1}{2}) \right) \quad (14.545)$$

and

$$\text{sd}(u, \frac{1}{2}) + \sqrt{2} \text{cd}(u, \frac{1}{2}) \text{nd}(u, \frac{1}{2}) \neq 0. \quad (14.546)$$

Using (14.545–546) one gets

$$\frac{dn_s}{d \ln K} = -2C_1^2 x^2 \text{sd}^2(u, \frac{1}{2}) \quad (14.547)$$

if Eqs (14.545–546) are satisfied.

In the case of special C_1 and C_2 one gets

$$\frac{dn_s}{d \ln K} = -0.5525x^2 \text{sd}^2(u, \frac{1}{2}) \quad (14.548)$$

where x, u are given by Eq. (14.544a).

Let us reparametrize the Eq. (14.545). One gets

$$\text{sd}(u, \frac{1}{2}) + \sqrt{2} \text{cd}(u, \frac{1}{2}) \text{nd}(u, \frac{1}{2}) = \frac{u - \sqrt{2}C_2}{\sqrt{2}} \left(\text{sd}^3(u, \frac{1}{2}) - \sqrt{2} \text{cd}(u, \frac{1}{2}) \right). \quad (14.549)$$

For sufficiently big u one gets (the equation has infinite number of roots)

$$u = u_1 + 2lK(\frac{1}{2}), \quad l = 0, \pm 1, \pm 2, \dots \quad (14.550)$$

where u_1 satisfies the equation

$$\text{sn}(u_1, \frac{1}{2}) = \frac{1}{3} \left(1 + \sqrt[3]{5} \left(\sqrt[3]{65 + \sqrt{20357}} - \sqrt[3]{\sqrt{20357} - 65} \right) \right)^{\frac{1}{2}} \simeq 0.65 \dots \quad (14.551)$$

Moreover for $x = \frac{K}{R_0 H_0} > 0$ we have the condition

$$\frac{u_1 + 2lK(\frac{1}{2}) - \sqrt{2}C_2}{\sqrt{2}C_1} > 0. \quad (14.552)$$

One finds

$$\text{sd}^2(u_1, \frac{1}{2}) = 0.53 \dots \quad (14.553)$$

Thus

$$\frac{dn_s}{d \ln K} (u_1 + 2lK(\frac{1}{2})) \simeq -0.53 \left(\frac{u_1 + 2lK(\frac{1}{2}) - \sqrt{2}C_2}{\sqrt{2}C_1} \right)^2, \quad (14.554)$$

$$l = 0, \pm 1, \pm 2, \dots$$

One gets

$$\arcsin 0.65 \simeq 40^\circ.54 \quad (14.555)$$

$$u_1 \cong \int_0^{40^\circ.54} \frac{d\theta}{\sqrt{1 - \frac{1}{2} \sin^2 \theta}} = F(40^\circ.54/45^\circ) \cong 0.73 \dots \quad (14.556)$$

Taking special value of C_1 and C_2 one gets

$$x \cong 1.81 - 4.68l \quad (14.557)$$

where $l = 0, \pm 1, \pm 2, \dots$. For $x > 0$ it is easy to see that l should be nonpositive. Practically l should be a negative integer

$$l < -3 \quad (14.558)$$

and

$$\frac{dn_s}{d \ln K} \cong -0.29(1.81 - 4.68l)^2. \quad (14.559)$$

Thus for

$$K_l = R_0 H_0 (1.81 - 4.68l)$$

we get

$$n_s(K_l) \cong 1. \quad (14.560)$$

The important range of $\ln K$, $\Delta \ln K$ is about 10.

Thus

$$n_s(K) \simeq 1 \pm 2.9(1.81 - 4.68l)^2. \quad (14.561)$$

Taking $l = -3$ one gets

$$-300 < n_s(K) < 300 \quad (14.562)$$

and

$$P_R(K) \sim K^{1-n_s(K)}$$

is not flat in the range considered.

Moreover we can improve the results considering Eq. (14.554) for large C_1 . For large C_1 one gets

$$\frac{C_2}{C_1} \simeq -1. \quad (14.563)$$

Thus we find

$$\frac{dn_s}{d \ln K} \simeq -1.06 \left(u_1 + 2lK\left(\frac{1}{2}\right) + C_1 \right)^2. \quad (14.564)$$

Taking large value of C_1 in such a way that

$$C_1 = -2lK\left(\frac{1}{2}\right) - u_1 + \varepsilon \quad (14.565)$$

where ε is a small number,

$$\varepsilon \simeq 10^{-n},$$

one gets

$$\frac{dn_s}{d \ln K} \simeq -1.06 \cdot 10^{-2n}. \quad (14.566)$$

The last condition means that we should take

$$h \cong \frac{H_0|C_1|}{\sqrt{5}\alpha_s} = \frac{H_0}{\sqrt{5}\alpha_s} \left(0.73 - 2lK\left(\frac{1}{2}\right) - 10^{-n} \right) \quad (14.567)$$

where l is an integer, $l \simeq -100$.

Thus for some special values of h we can get arbitrarily small $\frac{dn_s}{d \ln K}$ which means we can achieve a flat power spectrum in the range considered ($\Delta \ln K \sim 10$),

$$n_s(K) = 1 \pm 10^{-(2n-1)}, \quad (14.568)$$

n arbitrarily big.

Let us consider the value of K corresponding to our value of $n_s(K) = 1$. One gets

$$x = \frac{0.73 + 2lK\left(\frac{1}{2}\right) - \sqrt{2}C_2}{\sqrt{2}C_1}. \quad (14.569)$$

Using our assumption on a large C_1 and Eq. (14.565) one finds

$$x \cong \left(1 - \frac{1}{\sqrt{2}} \right) - \frac{\varepsilon}{2lK\left(\frac{1}{2}\right)} \quad (14.570)$$

or (for ε is small and l quite big)

$$x \simeq 0.293 \quad (14.571)$$

and

$$K \simeq 0.3R_0H_0 \quad (14.572)$$

with the range $\Delta \ln K \sim 10$. It means that

$$0.3e^{-10} \leq \frac{K}{R_0 H_0} \leq 0.3 \cdot e^{10} \quad (14.573)$$

which gives us a full cosmologically interesting region

$$10^{-4} \leq \frac{K}{R_0 H_0} \leq 10^4. \quad (14.574)$$

We consider some cosmological consequences of the Nonsymmetric Kaluza–Klein (Jordan–Thiry) Theory. Especially we use the scalar field Ψ appearing in order to get cosmological models with a quintessence and phase transitions. We consider a dynamics of Higgs' fields with various approximations and models of inflation.

Eventually we develop a toy model of this dynamics to obtain an amount of inflation and $P_R(K)$ function (a spectral function for fluctuations). We calculate a spectral index $n_s(K)$ and $\frac{dn_s}{d \ln K}$ for this model (see Ref. [115]).

Finally let us consider in this section two problems. First we match our solutions of an evolution of the Universe, i.e. both de Sitter phases, radiation dominated Universe, matter dominated Universe and K-essence Universe (kinetic energy dominated quintessence). In order to simplify calculations we match the first de Sitter phase to radiation dominated Universe using some issues from the second de Sitter phase. Secondly we write down solution to the evolution of Universe with a cosmological constant (from the second de Sitter phase), with a radiation and with a matter (a dust). All of these material ingredients are evolving independently in this case, similarly as a matter (dust) and a quintessence for the model (14.224).

Let us consider Eq. (14.50) and let us match it to Eq. (14.215). One gets

$$R_1 = \sqrt[4]{\frac{B}{\rho_Q}} \sqrt{\sinh \left(\frac{2\sqrt{3}\sqrt{\rho_Q} M_{\text{Pl}}}{B} t_{\text{end}} \right)} \quad (14.575)$$

$$R_1 = R_0 e^{H_0 t_{\text{end}}} \quad (14.576)$$

where

$$t_{\text{end}} = \frac{N_0}{H_0} \quad (14.577)$$

is the time of the end of the first de Sitter phase. (N_0 is the amount of an inflation.)

Simultaneously we need a local conservation of an energy, i.e.

$$\tilde{\rho}_r(R_1) = \frac{B}{R_1^4} \quad (14.578)$$

where

$$\tilde{\rho}_r(R_1) = 6 (H_0^2 - H_1^2) \quad (14.579)$$

(see the formula (19.57)). After some calculations one gets

$$B = \frac{2\sqrt{3}M_{\text{pl}}\sqrt{\rho_Q}t_{\text{end}}}{\text{arsinh}\left(\sqrt{\frac{\rho_Q}{\tilde{\rho}_r}}\right)}. \quad (14.580)$$

Now we need match a solution with a radiation dominated period to the solution with matter dominated period (with the same cosmological constant—quintessence energy density). One gets

$$\frac{A}{R_2^3} + \rho_Q = \frac{B}{R_2^4} + \rho_Q \quad (14.581)$$

and finally

$$A = \frac{B}{R_2} \quad (14.582)$$

where

$$R_2 = 0.7916 \sqrt[3]{\frac{A}{\rho_Q}} = \sqrt[4]{\frac{B}{\rho_0}} \sqrt{\sinh\left(\frac{2\sqrt{3}\sqrt{\rho_Q}M_{\text{pl}}}{B}t_1\right)}, \quad (14.583)$$

t_1 is a time to match (see (14.247)). Let us consider

$$N_1 = H_1(t_1 - t_{\text{end}}) \quad (*)$$

as an amount of an inflation during the second de Sitter phase. In this way one gets

$$t_1 = \frac{N_1}{H_1} + \frac{N_0}{H_0} \quad (14.584)$$

and finally

$$A = 2.015 \cdot \frac{M_{\text{pl}}^{3/4} \rho_Q^{5/8} \left(\frac{N_0}{H_0}\right)^{3/4}}{\text{arsinh}^{4/3} \sqrt{\frac{\rho_Q}{\tilde{\rho}_r}}} \sinh^{3/2} \left(\sqrt{\frac{\rho_Q}{\tilde{\rho}_r}} \left(\frac{N_1}{N_0} \frac{H_0}{H_1} + 1 \right) \right) \quad (14.585)$$

$$B = \frac{2\sqrt{3}M_{\text{pl}}\sqrt{\rho_Q}}{\text{arsinh}\left(\sqrt{\frac{\rho_Q}{\tilde{\rho}_r}}\right)} \frac{N_0}{H_0}. \quad (14.586)$$

However, we should match also a time. In this way we get from Eqs (14.444–445)

$$t_0 = \frac{418.48}{\sqrt{\rho_Q}} M_{\text{pl}} + \frac{N_1}{H_1} \quad (14.587)$$

or using the value $\rho_Q = \lambda_{c0} = 10^{-52} \frac{1}{\text{m}^2}$ we get

$$t_0 = 0.447 \cdot 10^{14} \text{yr} + \frac{N_1}{H_1}. \quad (14.588)$$

If R_0 is an initial value of “a radius” of the Universe we get

$$\begin{aligned}
R_0 = R_1 e^{-N_0} &= \sqrt[4]{\frac{2\sqrt{3} M_{\text{pl}} \sqrt{\rho_Q}}{\tilde{\rho}_r \operatorname{arsinh}\left(\sqrt{\frac{\rho_Q}{\tilde{\rho}_r}}\right)}} \cdot \exp\left(-\left(N_0 + \frac{n+2}{4} \Psi_0\right)\right) \\
&= \sqrt[4]{\frac{2\sqrt{3} M_{\text{pl}} \sqrt{\rho_Q}}{\tilde{\rho}_r \operatorname{arsinh}\left(\sqrt{\frac{\rho_Q}{\tilde{\rho}_r}}\right)}} \cdot \left(\frac{(n+2)\beta}{n|\gamma|}\right)^{(n+2)/8} e^{-N_0}.
\end{aligned} \tag{14.589}$$

The only one ingredient from the second de Sitter phase is using the formula (*). Now we should match the solution (14.444–445) with conditions (14.247) to the solution (14.281), (14.273), (14.279). We should match a field Ψ , a density of an energy, a radius and a time. One gets

$$R_3 = \bar{R}_0 \exp\left(\frac{n+1}{M_{\text{pl}}}(t_2 - \bar{t}_0)\right) = 3.07 \sqrt[3]{\frac{A}{\rho_Q}} \tag{14.590}$$

$$\Psi(t_2) = \frac{1}{2n} \ln\left(\frac{\delta}{3|\gamma|}\right) - \frac{1}{n} \sqrt{\frac{2\delta}{3|\bar{M}|}} M_{\text{pl}}(t_2 - \bar{t}_0) = \Psi_0 \tag{14.591}$$

$$\rho_Q + \frac{A}{R_3^3} = \rho_0 e^{(n+2)\Psi_0} + \rho_\Psi \tag{14.592}$$

where

$$\rho_\Psi = \frac{\delta M_{\text{pl}}^2}{3n^2} - \frac{\sqrt{|\gamma|\delta}}{2\sqrt{3}} \exp\left(-\sqrt{\frac{2\delta}{3\bar{M}}} M_{\text{pl}}(t_2 - \bar{t}_0)\right) \tag{14.593}$$

$$\delta = 2 \frac{n+1}{M_{\text{pl}}^2} - \frac{3\rho_0}{M_{\text{pl}}^2} \tag{14.594}$$

$$\bar{t}_0 \neq 0. \tag{14.594a}$$

One finds:

$$t_2 = \sqrt{\frac{3\bar{M}}{2\delta}} \frac{1}{M_{\text{pl}}} \ln\left(\sqrt{\frac{\delta(n+2)^n \beta^n}{3|\gamma|^{n+1} n^n}}\right) + \bar{t}_0 \tag{14.595}$$

and t_2 is large,

$$\rho_\Psi \cong \frac{\delta M_{\text{pl}}^2}{3n^2} = \frac{M_{\text{pl}}^2}{3n^2} \left(2 \frac{n+1}{M_{\text{pl}}^2} - \frac{3\rho_0}{M_{\text{pl}}^2}\right) \tag{14.596}$$

$$\rho_0 = \frac{n^2 \lambda_{c0} - 3(n+1)}{n^2 x_0^{n+2} - 1}. \tag{14.597}$$

From

$$\begin{aligned}\frac{A}{R_3^3} &= -\rho_Q + \rho_0 \left(x_0^{n+2} - \frac{1}{n^2} \right) + \frac{3(n+1)}{n^2} \\ R_3 &= 3.072 \sqrt[3]{\frac{A}{\rho_Q}}\end{aligned}\tag{14.598}$$

one obtains

$$\delta = \frac{1}{M_{\text{pl}}^2} \left(2(n+1) - 3 \frac{\lambda_{c0} n^2 - 3(n+1)}{n^2 x_0^{n+2} - 1} \right).\tag{14.599}$$

From Eqs (14.247), (14.444) and (14.445) we get

$$t_2 = \frac{N_1}{H_1} + \frac{5.38}{\sqrt{\rho_Q}} M_{\text{pl}}\tag{14.600}$$

$$\bar{t}_0 = \frac{N_1}{H_1} + \frac{5.38}{\sqrt{\rho_Q}} M_{\text{pl}} - \sqrt{\frac{3\bar{M}}{2\delta}} \frac{1}{M_{\text{pl}}} \ln \left(\sqrt{\frac{\delta(n+2)^n \beta^n}{3|\gamma|^{n+1} n^n}} \right)\tag{14.601}$$

$$M_{\text{pl}} \frac{5.38}{\sqrt{\rho_Q}} = 5.64 \cdot 10^{10} \text{ yr}.\tag{14.602}$$

Finally we find $\bar{R}_0$:

$$\bar{R}_0 = R_3 \exp \left(-\frac{n+1}{M_{\text{pl}}} (t_2 - \bar{t}_0) \right) = 3.072 \sqrt[3]{\frac{A}{\rho_Q}} \exp \left(-\frac{n+1}{M_{\text{pl}}} (t_2 - \bar{t}_0) \right).\tag{14.603}$$

In this way we match an evolution of the Universe from the very beginning up to K-essence dominance. We find all the constants. Let us notice that the solution (14.247), (14.444–445) has a quite known behaviour in the theory of ordinary nonlinear differential equations. The solution cannot be extended after some value of the dependent variable (and also independent). One says the solution dies. Due to this interesting behaviour we obtain physical consequences. Matter and a quintessence cannot evolve independently. We get also a ratio between a density of matter and a quintessence which agrees with observational data and helps (in principle) to calculate a time of our contemporary epoch.

If we use Eqs (14.444) and (14.445) we can calculate a time of our contemporary epoch, i.e. a time when the ratio of a density of a matter to a quintessence density is $\frac{3}{7}$ (this is this ratio measured now). One gets

$$\bar{t}_{\text{contemporary}} = 8.97 \frac{M_{\text{pl}}}{\sqrt{\rho_Q}} + \frac{N_1}{H_1} = 9.42 \cdot 10^{10} \text{ yr} + \frac{N_1}{H_1}.\tag{I}$$

In the same way using Eqs (14.444), (14.445) and (14.446) we calculate a Hubble parameter for our contemporary epoch.

$$H_{\text{contemporary}} = h_{\text{contemporary}} \frac{100 \frac{\text{km}}{\text{s}}}{\text{Mpc}} \quad (\text{II})$$

where

$$h_{\text{contemporary}} = 1.21. \quad (\text{IIa})$$

Both values are a little too big.

Moreover one can improve the results. However, the age of the Universe is a little longer,

$$t_{\text{age of the Universe}} = t_{\text{contemporary}} + t_1 = 9.42 \cdot 10^{10} \text{ yr} + 2 \frac{N_1}{H_1} + \frac{N_0}{H_0}. \quad (\text{Ia})$$

In general one gets

$$H = \frac{\sqrt{\rho_Q}}{M_{\text{pl}}} \frac{32.27 \sqrt[3]{\frac{\rho_Q}{\rho_m}} - 16.38}{14.96 \sqrt[3]{\frac{\rho_Q}{\rho_m}} - 0.03}$$

and

$$-q = \frac{5.21}{11.76 \sqrt[3]{\frac{\rho_Q}{\rho_m}} - 5.17} + 1$$

where $\frac{\rho_Q}{\rho_m}$ is a ratio of a quintessence energy density to a matter (dust) energy density, or

$$h = 0.9115 \frac{32.27 \sqrt[3]{\frac{\rho_Q}{\rho_m}} - 16.38}{14.96 \sqrt[3]{\frac{\rho_Q}{\rho_m}} - 0.03}.$$

Finally, let us express an age of the Universe in terms of the ratio $\frac{\rho_Q}{\rho_m}$. One gets

$$t \left(\frac{\rho_Q}{\rho_m} \right) = 2.667 \ln \left(31.87 \sqrt[3]{\frac{\rho_Q}{\rho_m}} - 14.03 \right) \frac{M_{\text{pl}}}{\rho_Q} + 2 \left(\frac{N_1}{H_1} \right) + \left(\frac{N_0}{H_0} \right)$$

or

$$t \left(\frac{\rho_Q}{\rho_m} \right) = 2.82 \cdot 10^{10} \text{ yr} \cdot \ln \left(31.87 \sqrt[3]{\frac{\rho_Q}{\rho_m}} - 14.03 \right) \frac{M_{\text{pl}}}{\rho_Q} + 2 \left(\frac{N_1}{H_1} \right) + \left(\frac{N_0}{H_0} \right).$$

For $\frac{\rho_m}{\rho_Q} = 0.0034$ one gets

$$t(294.12) = 14.91 \cdot 10^{10} \text{ yr} + 2 \left(\frac{N_1}{H_1} \right) + \left(\frac{N_0}{H_0} \right).$$

For $\frac{\rho_m}{\rho_Q} = 2.015$

$$t(0.496) = 6.76 \cdot 10^{10} \text{ yr} + 2 \left(\frac{N_1}{H_1} \right) + \left(\frac{N_0}{H_0} \right).$$

$$\Delta t = t(294.12) - t(0.496) = 8.15 \cdot 10^{10} \text{ yr}.$$

The last number is a duration time of the epoch for ρ_Q is equal to the contemporary value of a cosmological constant.

Let us come back to the Eq. (14.209) and consider it in the flat case $K = 0$. One gets

$$\frac{R dR}{\sqrt{\frac{\rho_Q}{3M_{\text{pl}}^2} \cdot R^4 + \frac{A}{3M_{\text{pl}}^2} \cdot R + \frac{B}{3M_{\text{pl}}^2}}} = \pm dt. \quad (14.604)$$

We change R into x :

$$R = \sqrt[3]{\frac{A}{\rho_Q}} x \quad (14.605)$$

and finally get

$$\pm \int \frac{x dx}{\sqrt{x^4 + x + a}} = \sqrt{\frac{\rho_Q}{3M_{\text{pl}}^2}} (t - t_0) \quad (14.606)$$

where

$$0 < a = \frac{B}{A} \sqrt[3]{\frac{\rho_Q}{A}}. \quad (14.607)$$

This is an evolution of a flat Universe with radiation, matter (dust) and a density of a quintessence (this is a cosmological constant λ_{c0}). The integral on LHS of Eq. (14.606) is an elliptic integral and can be calculated. The properties of the result of calculations strongly depend on the value of a . Let us notice that the integral we have calculated here has $a = 0$ (no radiation).

The evolution of a quintessence, a matter and a radiation is here independent. One gets the following results for

$$I = \int \frac{x dx}{\sqrt{x^4 + x + a}}. \quad (14.608)$$

In general there are two cases:

$$\text{I} \quad B_2 > 0 \quad (14.609)$$

$$\text{II} \quad B_2 < 0 \quad (14.610)$$

where

$$B_2 = \frac{\sqrt{12b_1^4 - a} - 2b_1^2 - \sqrt{4b_1^4 - a}}{2\sqrt{12b_1^4 - a}} \quad (14.611)$$

and $b_1 > 0$ is a solution of the cubic equation

$$y^3 - ay - \frac{1}{8} = 0 \quad (14.612)$$

$$y = 2b_1^4 \quad (14.613)$$

Eq. (14.612) can be solved by using the Cardano formulae in two cases

- 1) $D > 0$,
- 2) $D < 0$,

$$D = \frac{27 - 256a^3}{27 \cdot 256}. \quad (14.614)$$

In case 1) one gets

$$b_1 = \frac{1}{2} \sqrt[4]{\sqrt[3]{4 + \frac{1}{2}\sqrt{27 - 256a^3}} + \sqrt[3]{4 - \frac{1}{2}\sqrt{27 - 256a^3}}} > 0 \quad (14.615)$$

$$0 < a < \frac{3}{4\sqrt[3]{4}} = 0.472470393\dots$$

In case 2) one gets

$$b_1 = \sqrt[4]{\frac{a}{3}} \sqrt{\cos \frac{\varphi}{3}} \quad (14.616)$$

where

$$\cos \varphi = \frac{3}{16a} \sqrt{\frac{3}{a}} \quad (14.617)$$

$$a > \frac{3}{4\sqrt[3]{4}}, \quad \varphi \in \langle 0, \frac{\pi}{2} \rangle. \quad (14.618)$$

Condition I can be transformed into

$$4b_1^2 < a + 1 \quad (14.619)$$

for both cases 1) and 2). In case 1) this condition has no solution. It means we have always $B_2 < 0$. In case 2) we have always (14.619). It means

$$B_2 > 0 \quad (14.620)$$

for an equation

$$\frac{1}{\sqrt[4]{4}} \frac{\cos^2(\frac{\varphi}{3})}{\cos^{4/3}\varphi} = \frac{3}{4\sqrt[3]{4} \cos^{2/3}\varphi} + 1. \quad (14.621)$$

has no solution in $(0, \frac{\pi}{2})$ and in $(\frac{3\pi}{2}, 2\pi)$.

The fact that in case 1) we have always

$$4b_1^2 > a + 1 \quad (14.622)$$

is caused by a nonexistence of real solutions of an equation

$$4(a+1)^{1/2} = \sqrt[3]{4 + \frac{1}{2}\sqrt{27 - 256a^3}} + \sqrt[3]{4 - \frac{1}{2}\sqrt{27 - 256a^3}} \quad (14.623)$$

where

$$0 < a < \frac{3}{4\sqrt[3]{4}}. \quad (14.624)$$

Thus for sufficiently small a we always have case I (in a limit $a = 0$, no radiation also).

One can express b_1 and a in terms of φ :

$$b_1^4 = \frac{3}{4\sqrt[3]{4}} \frac{\cos^2(\frac{\varphi}{3})}{\cos^{2/3}\varphi} \quad (14.625)$$

$$a = \frac{3}{4\sqrt[3]{4}} \cos^{-2/3}\varphi. \quad (14.626)$$

Thus one obtains in case I

$$\begin{aligned} I = \frac{1}{2} \ln & \frac{e(\frac{x-\alpha}{x-\beta})^2 + f + 2\sqrt{((\frac{x-\alpha}{x-\beta})^2 + c)((\frac{x-\alpha}{x-\beta})^2 + d)(1+c)(1+d)}}{(\frac{x-\alpha}{x-\beta})^2 - 1} \\ & + K_1(b_1, a) \Pi \left(\operatorname{arctg} \left[P(b_1, a) \frac{x-\alpha}{x-\beta} \right], n_1, q_1 \right) \\ & + L_1(b_1, a) F \left(\operatorname{arctg} \left[P(b_1, a) \frac{x-\alpha}{x-\beta} \right], q_1 \right) \end{aligned} \quad (14.627)$$

$$P_1(b_1, a) = \left(\frac{\sqrt{12b_1^4 - a} - 2b_1^2 - \sqrt{4b_1^4 - a}}{\sqrt{12b_1^4 - a} + 2b_1^2 + \sqrt{4b_1^4 - a}} \right)^{1/2} \quad (14.628)$$

$$q_1 = \frac{\sqrt{6}|b_1|}{\left(3b_1^2 + \sqrt{12b_1^4 - a} \right)^{1/2}} \quad (14.629)$$

$$K_1(b_1, a) = \left(\frac{2b_1^2 + \sqrt{4b_1^4 - a} + \sqrt{12b_1^4 - a}}{\sqrt{12b_1^4 - a} - 2b_1^2 - \sqrt{4b_1^4 - a}} \right)^{3/2} \quad (14.630)$$

$$n_1 = \frac{2\sqrt{12b_1^4 - a}}{2b_1^2 + \sqrt{4b_1^4 - a} + \sqrt{12b_1^4 - a}} \quad (14.631)$$

$$L_1(b_1, a) = \left(M(b_1, a) + \frac{2b_1^2 + \sqrt{4b_1^4 - a} + \sqrt{12b_1^4 - a}}{2\sqrt{12b_1^4 - a}} \right) \frac{1}{(3b_1^2 + \sqrt{12b_1^4 - a})^{1/2}} \\ \times \frac{2(12b_1^4 - a)^{1/2} \left(\sqrt{4b_1^4 - a} - b_1^2 \right)^{1/2}}{\left(2b_1^2 + \sqrt{4b_1^4 - a} - \sqrt{12b_1^4 - a} \right) \left(2b_1^2 + \sqrt{4b_1^4 - a} + \sqrt{12b_1^4 - a} \right)^{1/2}} \quad (14.632)$$

$$M(b_1, a) = \frac{\beta}{\alpha - \beta} \\ = \left((6b_1^4 + a)\sqrt{4b_1^4 - a} + 2b_1^2a + \sqrt{192b_1^{12} - 112b_1^8a + 20b_1^4a + a^3} \right. \\ \left. + \sqrt{576b_1^{12} - 240b_1^8a + 28b_1^4a - a^3} + 6b_1^2\sqrt{48b_1^8 - 16b_1^4a + a^2} \right. \\ \left. + (a - 4b_1^4)\sqrt{12b_1^4 - a} \right)^{1/2} \\ \times \left(\left((12b_1^4 - a)\sqrt{4b_1^4 - a} + 4b_1^4\sqrt{12b_1^4 - a} - 4b_1^2a + 24b_1^6 + 4b_1^4\sqrt{12b_1^4 - a} \right. \right. \\ \left. \left. - \sqrt{192b_1^{12} - 112b_1^8a + 20b_1^4a + a^3} - \sqrt{576b_1^{12} - 240b_1^8a + 28b_1^4a - a^3} \right)^{1/2} \right. \\ \left. - \left(8b_1^2\sqrt{48b_1^8 - 16b_1^4a + a^2} + 2(6b_1^4 - a)\sqrt{4b_1^4 - a} - 4b_1^4\sqrt{12b_1^4 - a} \right. \right. \\ \left. \left. + 4b_1^6 + 6b_1^2a + \sqrt{192b_1^{12} - 112b_1^8a + 20b_1^4a + a^3} \right. \right. \\ \left. \left. + \sqrt{576b_1^{12} - 240b_1^8a + 28b_1^4a - a^3} \right)^{1/2} \right)^{-1}. \quad (14.633)$$

$$\alpha = \sqrt{\frac{\sqrt{48b_1^8 - 16b_1^4a + a^2} + 6b_1^2\sqrt{4b_1^4 - a} + 2b_1^2\sqrt{12b_1^4 - a} + a}{2b_1^2 + \sqrt{4b_1^4 - a} + \sqrt{12b_1^4 - a}}} > 0 \quad (14.634)$$

$$\beta = \sqrt{\frac{\sqrt{48b_1^8 - 16b_1^4a + a^2} + 6b_1^2\sqrt{4b_1^4 - a} - 2b_1^2\sqrt{12b_1^4 - a} + a}{2b_1^2 - \sqrt{4b_1^4 - a} - \sqrt{12b_1^4 - a}}} > 0 \quad (14.635)$$

$$c = \frac{(3b_1^2 + \sqrt{12b_1^4 - a})(2b_1^2 - \sqrt{4b_1^4 - a} - \sqrt{12b_1^4 - a})}{(\sqrt{4b_1^4 - a} + 2b_1^2)(\sqrt{12b_1^4 - a} - 3b_1^2)}$$

$$d = \frac{\sqrt{12b_1^4 - a} + 2b_1^2 + \sqrt{4b_1^4 - a}}{\sqrt{12b_1^4 - a} - 2b_1^2 - \sqrt{4b_1^4 - a}}, \quad |d| > 1 \quad (14.636)$$

$$e = \frac{2\sqrt{12b_1^4 - a} \left(\sqrt{12b_1^4 - a} + \sqrt{4b_1^4 - a} - 4b_1^2 \right)}{\left(\sqrt{4b_1^4 - a} + 2b_1^2 + \sqrt{12b_1^4 - a} \right) \left(\sqrt{12b_1^4 - a} - 3b_1^2 \right)} \quad (14.637)$$

$$\begin{aligned} f = & \frac{2\sqrt{12b_1^4 - a}}{\left(\sqrt{12b_1^4 - a} - 3b_1^2 \right) \left(\sqrt{12b_1^4 - a} - 2b_1^2 - \sqrt{4b_1^4 - a} \right)} \\ & \times \left(2b_1^2 + \sqrt{4b_1^4 - a} + \sqrt{12b_1^4 - a} \right)^{-2} \\ & \times \left(2(14b_1^4 - a)\sqrt{4b_1^4 - a} - 4b_1^2a + 2\sqrt{192b_1^{12} - 112b_1^8a + 20b_1^4a + a^3} \right. \\ & \left. + 2(b_1^4 - a)\sqrt{12b_1^4 - a} - 2\sqrt{576b_1^{12} - 240b_1^8a + 28b_1^4a - a^3} \right). \end{aligned} \quad (14.638)$$

Case II

$$\begin{aligned} I = & \frac{1}{2} \ln \frac{e\left(\frac{x-\alpha}{x-\beta}\right)^2 + f + 2\sqrt{\left(\left(\frac{x-\alpha}{x-\beta}\right)^2 + c\right)\left(\left(\frac{x-\alpha}{x-\beta}\right)^2 + d\right)(1+c)(1+d)}}{\left(\frac{x-\alpha}{x-\beta}\right)^2 - 1} \\ & + K_2(b_1, a) \Pi \left(\arccos \left[P_2(b_1, a) \frac{x - \alpha}{x - \beta} \right], n_2, q_2 \right) \\ & + L_2(b_1, a) F \left(\arccos \left[P_2(b_1, a) \frac{x - \alpha}{x - \beta} \right], q_2 \right) \end{aligned} \quad (14.639)$$

$$q_2 = \frac{1}{q_1} = \frac{\left(3b_1^2 + \sqrt{12b_1^4 - a} \right)^{1/2}}{\sqrt{6}|b_1|} \quad (14.640)$$

$$n_2 = \frac{2b_1^2 + \sqrt{4b_1^4 - a} + \sqrt{12b_1^4 - a}}{2\sqrt{12b_1^4 - a}} \quad (14.641)$$

$$K_2(b_1, a) = \frac{\left(2b_1^2 + \sqrt{4b_1^4 - a} - \sqrt{12b_1^4 - a} \right) \sqrt{12b_1^4 - a} \left(\sqrt{4b_1^4 - a} - b_1 \right)^{1/2}}{\left(\sqrt{12b_1^4 - a} + 2b_1^2 + \sqrt{4b_1^4 - a} \right) \left(\sqrt{12b_1^4 - a} - 3b_1^2 \right)^{3/2}} \quad (14.642)$$

$$P_2(b_1, a) = \left(\frac{2b_1^2 + \sqrt{4b_1^4 - a} - \sqrt{12b_1^4 - a}}{\sqrt{12b_1^4 - a} + 2b_1^2 + \sqrt{4b_1^4 - a}} \right)^{1/2} \quad (14.643)$$

$$\begin{aligned} L_2(b_1, a) = & M(b_1, a) \\ & - \frac{4(12b_1^4 - a) \left(\sqrt{4b_1^4 - a} - b_1^2 \right)^{1/2} \left(\sqrt{12b_1^4 - a} - 2b_1^2 - \sqrt{4b_1^4 - a} \right)}{\left(3b_1^2 - \sqrt{12b_1^4 - a} \right)^{3/2} \left(2b_1^2 + \sqrt{4b_1^4 - a} + \sqrt{12b_1^4 - a} \right)}. \end{aligned} \quad (14.644)$$

The function (14.606) cannot be inverted globally. It can be inverted only locally in some intervals where the solution lives. The solution dies on the end of an interval being reborn on beginning of a next interval. All the intervals can be obtained from the condition

$$\overline{W} > 0 \quad (14.645)$$

where

$$\overline{W} = \frac{e(\frac{x-\alpha}{x-\beta})^2 + f + 2\sqrt{((\frac{x-\alpha}{x-\beta})^2 + c)((\frac{x-\alpha}{x-\beta})^2 + d)(1+c)(1+d)}}{(\frac{x-\alpha}{x-\beta})^2 - 1} \quad (14.646)$$

and

$$\left| \frac{x-\alpha}{x-\beta} \right| \neq 1 \quad (14.647)$$

In the case I $c > 0, d > 0$.

In the case II $c > 0, d < 0$.

Thus one can get very interesting behaviour from the physical point of view, because every end of an interval of living (existence) of the solution of an evolution equation means a start of nontrivial interaction between a matter, a radiation and a quintessence. In all of these intervals one can find an approximate inverse function, i.e. one can write

$$R = F(t) \quad (14.648)$$

where

$$F(t_1) = R_1 < R < R_2 = F(t_2) \quad (14.649)$$

and

$$t_1 < t < t_2. \quad (14.650)$$

Moreover we do not develop this project here in details. F and II are usual elliptic integrals of the first and of the third kind given by the formulae (14.228) and (14.230).

For future convenience of this project we find roots of the polynomial

$$P(x) = x^4 + x + a, \quad a > 0. \quad (14.651)$$

First of all according to the Ferrari method we should solve a resolvent equation which is a cubic equation. One gets

$$z^3 - 4az - 1 = 0. \quad (14.652)$$

The discriminant of Eq. (14.652) reads

$$\overline{D} = \frac{27 - 256a^3}{108}$$

and we have two cases

$$\overline{D} > 0, \quad 0 < a < \frac{3}{4\sqrt[3]{4}}, \quad (14.653)$$

$$\overline{D} < 0, \quad a > \frac{3}{4\sqrt[3]{4}}, \quad (14.654)$$

In the case (14.653) one gets

$$z_1 = \sqrt[3]{\frac{\sqrt{27 - 256a^3} + 3\sqrt{3}}{6\sqrt{3}}} - \sqrt[3]{\frac{\sqrt{27 - 256a^3} - 3\sqrt{3}}{6\sqrt{3}}} \quad (14.655)$$

$$\begin{aligned} z_{2,3} = & \frac{1}{2} \left(\sqrt[3]{\frac{\sqrt{27 - 256a^3} - 3\sqrt{3}}{6\sqrt{3}}} - \sqrt[3]{\frac{\sqrt{27 - 256a^3} + 3\sqrt{3}}{6\sqrt{3}}} \right) \\ & \pm \frac{i\sqrt{3}}{2} \left(\sqrt[3]{\frac{\sqrt{27 - 256a^3} - 3\sqrt{3}}{6\sqrt{3}}} + \sqrt[3]{\frac{\sqrt{27 - 256a^3} + 3\sqrt{3}}{6\sqrt{3}}} \right) \end{aligned} \quad (14.656)$$

In the second case (14.654)

$$z_1 = 4\sqrt{\frac{a}{3}} \cos\left(\frac{\overline{\varphi}}{3}\right) \quad (14.657)$$

$$z_2 = 4\sqrt{\frac{a}{3}} \cos\left(\frac{\overline{\varphi}}{3} + \frac{2\pi}{3}\right) \quad (14.658)$$

$$z_3 = 4\sqrt{\frac{a}{3}} \cos\left(\frac{\overline{\varphi}}{3} + \frac{4\pi}{3}\right) \quad (14.659)$$

where $\cos \overline{\varphi} = \sqrt{\frac{3}{16a}}$.

We can also distinguish the special case $\overline{D} = 0$ ($a = \frac{3}{4\sqrt[3]{4}}$):

$$z_1 = \sqrt[3]{4} \quad (14.660)$$

$$z_{2,3} = -\frac{1}{2}\sqrt[3]{4} \quad (14.661)$$

In the first case we get one real and two complex conjugated roots, in the second three real roots, in the third case two different real roots (one of them is double).

According to the Ferrari method one gets

$$2x_1 = \sqrt{z_1} + \sqrt{z_2} + \sqrt{z_3} \quad (14.662a)$$

$$2x_2 = \sqrt{z_1} - \sqrt{z_2} - \sqrt{z_3} \quad (14.662b)$$

$$2x_3 = -\sqrt{z_1} + \sqrt{z_2} - \sqrt{z_3} \quad (14.662c)$$

$$2x_4 = -\sqrt{z_1} + \sqrt{z_2} + \sqrt{z_3} \quad (14.662d)$$

$$z_1 z_2 z_3 = 1. \quad (14.663)$$

In the future analysis of an evolution of our model of the Universe there are three important cases for z_1, z_2 and z_3 :

- A. All z_1, z_2, z_3 are real and positive. In this case all x_1, x_2, x_3 and x_4 are real.
- B. One root (e.g. z_1) is positive, two remaining (z_2, z_3) are negative. In this case we have two pairs of conjugate roots.
- C. One root (e.g. z_1) is positive, two remaining (z_2, z_3) are complex conjugate. In this case we have two real roots and one pair of complex conjugate roots.

This analysis is quite important because only for positive real roots the integral I can have singular points. It means in A and C case. This gives us some restrictions on the parameter a .

Let us notice that in the case of $a = 0$ (considered before) we have to do with the case C. Fortunately one real root is zero and the second real root is negative ($x_2 = -1$).

Let us notice that our case with $a = 0$ is in some sense exceptional from the point of view of a general theory. In that case some of the formulae are singular and because of this it was considered separately.

Let us notice that for sufficiently big a the polynomial (14.651) has no real roots. This is true for $a > \frac{3}{4\sqrt[3]{4}}$. In this case the integral (14.608) has no singular points. This is case II.

Let us notice a simple relation between D and $\overline{D}$, $\overline{D} = 64D$. This connects Eq. (14.652) to Eq. (14.612) and gives the same restriction for a .

15. Geodetic Equations on the Manifold P

Let us consider geodetic equation for a curve $\Gamma \subset P$ on the manifold P i.e. a nonsymmetrically metrized fibre bundle over $E \times G/G_0$ with respect to the connection $\omega^{\tilde{A}}_{\tilde{B}}$. Such equations have a usual interpretation as test particle equations of motion. One gets ($\nabla_u u = 0$, where u is tangent to Γ):

$$u^{\tilde{A}} \nabla_{\tilde{A}} u^{\tilde{B}} = 0. \quad (15.1)$$

Using formulae (15.1), (5.8), (5.9), (5.11) one gets:

$$\begin{aligned} \frac{\overline{D}u^\alpha}{d\tau} + \left(\frac{q^c}{m_0}\right) u^\beta \left[\ell_{cd} g^{\alpha\delta} H^d_{\beta\delta} - \frac{1}{2}(\ell_{cd} g^{\alpha\delta} - \ell_{dc} g^{\delta\alpha}) L^d_{\beta\delta} + \right. \\ \left. - \frac{\|q\|^2}{8m_0^2} \tilde{g}^{(\alpha\beta)} \left(\frac{1}{\rho^2}\right)_{,\beta} + \left(\frac{q^c}{m_0}\right) u^{\tilde{b}} \left[\ell_{cd} g^{\alpha\delta} \nabla_\delta^{\text{gauge}} \Phi^d_{\tilde{b}} + \right. \right. \\ \left. \left. - \frac{1}{2}(\ell_{cd} g^{\alpha\delta} - \ell_{dc} g^{\delta\alpha}) L^d_{\tilde{b}\delta} \right] \right] = 0, \quad (15.2) \end{aligned}$$

$$\begin{aligned} \frac{\tilde{D}u^{\tilde{a}}}{d\tau} + \frac{1}{r^2} \left(\frac{q^c}{m_0}\right) u^\beta \left[\ell_{cd} g^{\tilde{a}\tilde{d}} \nabla_\beta^{\text{gauge}} \Phi^d_{\tilde{d}} - \frac{1}{2}(\ell_{cd} g^{\tilde{a}\tilde{d}} - \ell_{dc} g^{\tilde{d}\tilde{a}}) L^d_{\beta\tilde{d}} \right] + \\ + \frac{1}{r^2} \left(\frac{q^c}{m_0}\right) u^{\tilde{b}} \left[\ell_{cd} g^{\tilde{a}\tilde{d}} (C^d_{ab} \Phi^a_{\tilde{d}} \Phi^b_{\tilde{b}} - \mu^d_i f^i_{\tilde{d}\tilde{b}} - \Phi^d_{\tilde{e}} f^{\tilde{e}}_{\tilde{d}\tilde{b}}) + \right. \\ \left. - \frac{1}{2}(\ell_{cd} g^{\tilde{a}\tilde{d}} - \ell_{dc} g^{\tilde{d}\tilde{a}}) L^d_{\tilde{b}\tilde{d}} \right] = 0, \quad (15.3) \end{aligned}$$

$$\frac{d}{d\tau} \left(\frac{q^b}{m_0} \right) = 0, \quad (15.4)$$

where $\overline{D}$ means a covariant derivative along a line with respect to the connection $\overline{\omega}^\alpha_\beta$ on E , $\tilde{D}$ means a covariant derivative along a line with respect to the connection $\tilde{\omega}^{\tilde{a}}_{\tilde{b}}$ on G/G_0 ($r = \text{const}$).

$$u^{\tilde{A}} = (u^\alpha, u^{\tilde{a}}, u^a) = (\text{hor}(u), \text{ver}(u)) \quad (15.5)$$

and

$$2\rho^2 u^a = \frac{q^a}{m_o} \quad (q = m_0 \hat{\mathcal{K}}(\text{ver } u(\tau))),$$

u^α — is a four-velocity of a test particle, q^a its colour (isotopic) charge, $u^{\tilde{a}}$ a charge associated with a Higgs' field. This charge transforms according to the properties of a complement $\mathfrak{m}$ with respect to G_0 and G . One has $\text{hor}(u) = (P_E(\text{hor}(u)), P_{M=G/G_0}(\text{hor}(u)))$ where P_E is a projection on E and $P_{M=G/G_0}$ on M in the Cartesian product $V = E \times M = E \times G/G_0$.

Eq. (15.2) describes a movement of a test particle in a gravitational, gauge and Higgs' field. We should consider (of course) physical world-lines on E i.e. time-like or null-like.

Eq. (15.3) is an equation for a charge associated with Higgs' field. This charge describes a coupling between a test particle and a Higgs' field.

Eq. (15.4) has a usual meaning (a constancy of a colour (isotopic) charge).

In order to understand the significance of a new term in equation of motion for a test particle with nonzero colour and Higgs' charges let us suppose that:

$$g_{\alpha\beta} = g_{\beta\alpha}, \quad \ell_{cd} = \ell_{dc} = h_{cd}, \quad g_{\tilde{a}\tilde{b}} = h^0_{\tilde{a}\tilde{b}} = g_{\tilde{b}\tilde{a}},$$

i.e. that we have to do with a symmetric case. We suppose also that $\rho = \text{const}$. One easily gets:

$$\frac{\overline{D}u^\alpha}{d\tau} + \left(\frac{q^c}{m_0}\right) h_{cd} g^{\alpha\delta} H^d_{\beta\delta} u^\beta + \left(\frac{q^c}{m_0}\right) h_{cd} g^{\alpha\delta} \overset{\text{gauge}}{\nabla}_\delta \Phi_b^d u^{\tilde{b}} = 0, \quad (15.6)$$

$$\begin{aligned} \frac{\tilde{D}u^{\tilde{a}}}{d\tau} + \left(\frac{q^c}{m_0}\right) h_{cd} h^{0\tilde{a}\tilde{d}} \overset{\text{gauge}}{D} \frac{\Phi_{\tilde{d}}^d}{d\tau} + \\ + \left(\frac{q^c}{m_0}\right) h_{cd} h^{0\tilde{a}\tilde{d}} (C_{ab}^d \Phi_d^a \Phi_b^{\tilde{b}} - \mu_i^d f_{\tilde{d}\tilde{b}}^{\hat{i}} - \Phi_{\tilde{e}}^d f_{\tilde{d}\tilde{b}}^{\tilde{e}}) u^{\tilde{b}} = 0. \end{aligned} \quad (15.7)$$

In Eq. (15.6) we have an additional term:

$$\left(\frac{q^c}{m_0}\right) h_{cd} g^{\alpha\delta} \overset{\text{gauge}}{\nabla}_\delta \Phi_b^d u^{\tilde{b}}, \quad (15.8)$$

which plays the role of Lorentz force term for a Higgs' field. The test particle couples Higgs' (its gauge derivative) via a Higgs' charge $u^{\tilde{a}}$. Propagation properties of the Higgs' charge are described by the Eq. (15.7). They are much more complex than that of a colour charge. In general this charge is not constant during a motion. We put:

$$\frac{\overset{\text{gauge}}{D}}{d\tau} \Phi_{\tilde{d}}^d = \overset{\text{gauge}}{\nabla}_\beta \Phi_{\tilde{d}}^d u^\beta. \quad (15.9)$$

It is interesting to consider Eqs. (15.6) and (15.7) in the case of the completely broken group G i.e.

$$\frac{\overline{D}u^\alpha}{d\tau} + \left(\frac{q^c}{m_0}\right)h_{cd}g^{\alpha\delta}H_{\beta\delta}^d u^\beta + \left(\frac{q^c}{m_0}\right)h_{cd}g^{\alpha\delta}\overset{\text{gauge}}{\nabla}_\delta\Phi_b^d u^b = 0, \quad (15.10)$$

$$\frac{du^b}{d\tau} + \frac{1}{r^2}\left(\frac{q^c}{m_0}\right)h_{cd}\tilde{h}^{pq}\frac{\overset{\text{gauge}}{D}\Phi_q^d}{d\tau} + \frac{1}{r^2}\left(\frac{q^c}{m_0}\right)h_{cd}\tilde{h}^{bq}(C_{ap}^d\Phi_q^a\Phi_k^p - \Phi_e f_{qk}^e)u^k = 0. \quad (15.11)$$

In this case $\tilde{h}_{ab}$ is a Killing–Cartan tensor on G , u^b is an Ad-type quantity with respect to G . Moreover u^b is not a Yang–Mills’ gauge independent charge. This indicates that a charge which couples a test particle to the Higgs’ field has completely different properties than that known for Yang–Mills’ field.

Moreover for a vacuum state of the Higgs’ potential one gets:

$$\frac{\overline{D}u^\alpha}{d\tau} + \left(\frac{q^c}{m_0}\right)h_{cd}g^{\alpha\delta}H_{\beta\delta}^d u^\beta = 0, \quad (15.12)$$

$$\frac{du^b}{d\tau} = 0 \quad (15.13)$$

and u^b decouples from equations of motion.

If the Higgs’ field is closed to the vacuum configuration the Eq. (15.13) is not satisfied, but instead of it

$$\frac{d}{d\tau}\left(u^b + \left(\frac{q^c}{m_0}\right)h_{cd}\tilde{h}^{bq}\Phi_q^d\right) \cong 0 \quad (15.14)$$

and we can define a “constant” Higgs’ field charge:

$$g^b = u^b + \left(\frac{q^c}{m_0}\right)h_{cd}\tilde{h}^{bq}\Phi_q^d \quad (15.15)$$

and substitute it to Eq. (15.10). Thus we get:

$$\begin{aligned} \frac{\overline{D}u^\alpha}{d\tau} + \left(\frac{q^c}{m_0}\right)h_{cd}g^{\alpha\delta}H_{\beta\delta}^d u^\beta + \left(\frac{q^c g^b}{m_0}\right)h_{cd}g^{\alpha\delta}\overset{\text{gauge}}{\nabla}_\delta\Phi_b^d + \\ - \left(\frac{q^c q^e}{m_0}\right)h_{ef}\tilde{h}^{bq}\Phi_q^f h_{cd}g^{\alpha\delta}\overset{\text{gauge}}{\nabla}_\delta\Phi_b^d = 0. \end{aligned} \quad (15.16)$$

The Eq. (15.2) has the following first integral of motion:

$$\varkappa(u(\tau), u(\tau)) = g_{(\alpha\beta)}u^\alpha u^\beta + r^2 g_{\tilde{a}\tilde{b}}u^{\tilde{a}}u^{\tilde{b}} + \rho^2 \ell_{ab}u^a u^b = \text{const}. \quad (15.17)$$

Introducing the following notation:

$$\tilde{u}^2 = -g_{\tilde{a}\tilde{b}}u^{\tilde{a}}u^{\tilde{b}} \quad (15.18)$$

one gets:

$$g_{(\alpha\beta)}u^\alpha u^\beta - r^2\tilde{u}^2 - \frac{\|q\|^2}{4\rho^2} = \text{const}. \quad (15.18')$$

However in general the length of the charge $u^{\tilde{b}}$ is not constant even in the symmetric case. For a constant field $\rho = \text{const}$, one gets:

$$g_{(\alpha\beta)}u^\alpha u^\beta = r^2\tilde{u}^2 + \text{const}. \quad (15.19)$$

Thus the interpretation of the Eq. (15.3) demands more investigations.

Finally let us consider Eqs. (15.2–3) in a more general case for the scalar field $\rho = \rho(x, y)$. In this case we get one more term in Eq. (15.3) involving the field ρ ($\rho \in \mathbb{C}^1(E \times M)$) i.e.:

$$-\frac{\|q\|^2}{8m_0^2 r^2} \tilde{g}^{(\tilde{a}\tilde{b})} \left(\frac{1}{\rho^2} \right)_{,\tilde{b}} \quad (15.20)$$

and the term in Eq. (15.2):

$$-\frac{\|q\|}{8m_0^2} \tilde{g}^{(\alpha\beta)} \left(\frac{1}{\rho^2} \right)_{,\beta} \quad (15.21)$$

depends on the point $y \in M$.

If we suppose that:

$$\rho(x, y) = \rho_0(x)\rho_1(y),$$

then we get

$$-\frac{\|q\|^2}{8m_0^2 r^2 \rho_0^2} \tilde{g}^{(\tilde{a}\tilde{b})} \left(\frac{1}{\rho_1^2} \right)_{,\tilde{b}} \quad (15.20a)$$

and

$$-\frac{\|q\|^2}{8m_0^2 \rho_1^2} \tilde{g}^{(\alpha\beta)} \left(\frac{1}{\rho_0^2} \right)_{,\beta}. \quad (15.21a)$$

In the case of the expansion of ρ into generalized harmonics one gets:

$$\frac{\|q\|^2}{4m_0^2 r^2 \rho^3} \tilde{g}^{(\tilde{a}\tilde{b})} \left(\sum_{k=1}^{\infty} \rho_k(x) \chi_{k,\tilde{b}}(y) \right) \quad (15.20b)$$

and

$$\frac{\|q\|^2}{4m_0^2 \rho^3} \tilde{g}^{(\alpha\beta)} \left(\rho_{0,\beta} + \sum_{k=1}^{\infty} \rho_{k,\beta} \chi_k(y) \right). \quad (15.21b)$$

The last two terms describe an interaction of a test particle with a tower of scalar fields $\rho_k(x)$, $k = 0, 1, 2, \dots$. The truncation procedure for the field ρ substitutes ρ_0 in place of ρ . We can give an interpretation of normal coordinates as velocities, gauge-independent charges and Higgs' charges for test particles. We can consider in our

theory geodetic deviation equations. In this way we can derive “tidal-forces” for the Higgs’ field additionally to gravitational and Yang–Mills’ ones.

Eventually let us remark that in the Kaluza–Klein Theory we can treat the Higgs’ field as a part of a multidimensional Yang–Mills’ field. However the Lorentz-like force term for the Higgs’ field seems to be more complex.

Let us consider a geodetic deviation equation in our theory

$$u^{\tilde{B}} \nabla_{\tilde{B}} v^{\tilde{A}} + R^{\tilde{A}}_{\tilde{C}\tilde{M}\tilde{B}} u^{\tilde{C}} \zeta^{\tilde{M}} u^{\tilde{B}} - Q^{\tilde{N}}_{\tilde{M}\tilde{B}} \nabla_{\tilde{N}} u^{\tilde{A}} \zeta^{\tilde{M}} u^{\tilde{B}} = 0 \quad (15.22)$$

and for $u^{\tilde{B}}$ Eq. (15.1) is satisfied, where

$$u^{\tilde{A}} = \frac{dx^{\tilde{A}}}{d\tau}, \quad v^{\tilde{A}} = \frac{d\zeta^{\tilde{A}}}{d\tau}.$$

In this way we consider a generalization of the geodetic deviation equation to the $(n + n_1 + 4)$ -dimensional case in non-Riemannian geometry.

$R^{\tilde{A}}_{\tilde{C}\tilde{M}\tilde{B}}$ is a curvature tensor for a connection $\omega^{\tilde{A}}_{\tilde{B}}$ on P and $Q^{\tilde{A}}_{\tilde{C}\tilde{B}}$ is a tensor of a torsion for $\omega^{\tilde{A}}_{\tilde{B}}$, $\nabla_{\tilde{N}}$ is a covariant derivative with respect to $\omega^{\tilde{A}}_{\tilde{B}}$.

One gets using formulae from sec. 5 and Eqs (14.56) and (14.57) from the fourth point of Ref. [18]

$$\begin{aligned} & u^B \overline{\nabla}_B v^A + \overline{R}^A_{BMN} u^B \zeta^M u^N - \overline{Q}^N_{MB} \overline{\nabla}_N u^A \zeta^M u^B \\ & + \left(\frac{q^b}{2m_0 \rho^2} \right) l_{bd} \gamma^{AB} (2H^d_{NB} - L^d_{NB}) v^N - \rho^2 v^n l_{dn} \gamma^{DA} L^d_{DB} u^B \\ & - \left(\frac{q^n}{2m_0 \rho^2} \right) \tilde{\gamma}^{(AB)} \rho_{,B} l_{bn} v^b \\ & + 2\rho^2 \left(l_{cd} \gamma^{AP} \left(2H^d_{P[M} - L^d_{P[M} \right) L^c_{N]B} + l_{db} \gamma^{DA} L^d_{DB} H^b_{NM} \right) u^B \zeta^M u^N \\ & + \frac{q^b}{2m_0 \rho^2} \left[2\overline{\nabla}_{[M} \left(\rho^2 l_{db} \gamma^{AB} \left(2H^d_{N]B} - L^d_{N]B} \right) \right) \right. \\ & + \rho^2 l_{db} \gamma^{AB} (2H^d_{CB} - L^d_{CB}) \overline{Q}^C_{MN} - \rho \tilde{\gamma}^{(AB)} \rho_{,B} l_{bc} H^c_{MN} \\ & + 2\rho l_{bd} \gamma^{AB} \left(2H^d_{[M|B|} - L^d_{[M|B|} \right) \gamma_{|D|N]} \tilde{\gamma}^{(DC)} \rho_{,C} \left. \right] \zeta^M u^N \\ & + \left\{ \rho l_{bd} \gamma^{AP} \gamma_{BD} \tilde{\gamma}^{(DZ)} \rho_{,Z} (H^d_{MP} - L^d_{MP}) - \overline{\nabla}_M (\rho^2 l_{db} \gamma^{DA} L^d_{DB}) \right. \\ & - \rho \tilde{\gamma}^{(AP)} \rho_{,P} l_{cd} L^c_{BM} \left. \right\} u^B \left(\zeta^M \frac{q^b}{2\rho^2 m_0} - \zeta^b u^M \right) \\ & + \left(2\rho^4 l_{d[b} l_{e|f]} \gamma^{DA} \gamma^{ZC} L^d_{DC} L^e_{ZB} + \tilde{\gamma}^{(AP)} \rho_{,P} \gamma_{BD} \tilde{\gamma}^{(DZ)} \rho_{,Z} l_{[bf]} \right) \end{aligned} \quad (15.23)$$

$$\begin{aligned}
& + \frac{\rho^2}{2} l_{dp} \gamma^{DA} L_{DB}^d C_{bf}^P \Big) u^B \left(\frac{\zeta^b q^f}{2m_0 \rho^2} \right) \\
& + \left\{ \bar{\nabla}_M \left(\rho \tilde{\gamma}^{(AB)} \rho_{,B} \right) l_{ba} + \tilde{\nabla}_a \left(\rho^2 l_{bd} \gamma^{AB} (2H_{MB}^d - L_{MB}^d) \right) \right. \\
& - \rho^4 l_{da} l_{bf} \gamma^{DA} \gamma^{CB} L_{DC}^d (2H_{MB}^d - L_{MB}^d) \\
& \left. - \tilde{\gamma}^{(AZ)} \rho_{,Z} \gamma_{DM} \tilde{\gamma}^{(DN)} \rho_{,N} l_{ba} \right\} \left(\frac{q^b}{2m_0 \rho^2} \right) \left(\zeta^a u^M - \zeta^M \frac{q^a}{2m_0 \rho^2} \right) \\
& + \left(\rho \tilde{\gamma}^{(AB)} \rho_{,B} l_{bk} C_{ac}^k + 2\rho^3 \tilde{\gamma}^{(CB)} \rho_{,B} \gamma^{DA} L_{DC}^d l_{d[a|b|c]} \right) \left(\frac{q^b \zeta^a q^c}{4\rho^4 m_0^2} \right) \\
& - (2\rho^2 l_{bd} \gamma^{NB} H_{CB}^d + \rho^2 (l_{bd} \gamma^{NB} + l_{db} \gamma^{BN}) L_{BC}^d) \\
& \quad \times \left(\bar{\nabla}_N u^A + \frac{1}{2} \left(l_{fd} \gamma^{AB} (2H_{NB}^d - L_{NB}^d) \left(\frac{q^f}{m_0} \right) \right) \right) \left(v^C \frac{q^b}{2\rho^2 m_0} - v^b u^C \right) \\
& - 2\mu \rho \tilde{\gamma}^{(NB)} \rho_{,B} k_{bc} \left(\bar{\nabla}_N u^A + \frac{1}{2} l_{df} \gamma^{AC} (2H_{NC}^d - L_{NC}^d) \left(\frac{q^f}{m_0} \right) \right) \left(\frac{q^c v^b}{2m_0 \rho} \right) \\
& + 2 (H_{MN}^n - L_{NM}^n) \left(\rho^2 l_{dn} \gamma^{DA} L_{DB}^d u^B + \frac{1}{2\rho^2} \tilde{\gamma}^{(AB)} \rho_{,B} l_{bn} \left(\frac{q^b}{m_0} \right) \right) v^M u^N \\
& - \left(\gamma_{ZM} \tilde{\gamma}^{(AC)} \rho_{,C} \rho l_{bm} \gamma^{DZ} L_{DB}^b u^B + \frac{1}{2\rho^2} \tilde{\gamma}^{(AB)} \rho_{,B} l_{bm} \left(\frac{q^b}{m_0} \right) \right) \\
& \quad \times \left(v^M \left(\frac{q^m}{2\rho^2 m_0} \right) - v^m u^M \right) \\
& - \frac{1}{2\rho^2} \tilde{Q}^n{}_{mb} \left(\rho^2 l_{dn} \gamma^{DA} L_{DB}^d u^B - \frac{1}{2\rho} \tilde{\gamma}^{(AB)} \rho_{,B} l_{pn} \left(\frac{q^p}{m_0} \right) \right) v^m \left(\frac{q^b}{m_0} \right) \\
& = 0
\end{aligned} \tag{15.23cont.}$$

and

$$\begin{aligned}
& \frac{dv^a}{d\tau} + \tilde{R}^a{}_{bcd} \left(\frac{q^b}{2\rho^2 m_0} \right) \zeta^c \left(\frac{q^d}{2\rho^2 m_0} \right) - \tilde{Q}^a{}_{mb} \tilde{\nabla}_n \left(\frac{q^a}{2\rho^2 m_0} \right) v^m \left(\frac{q^b}{2\rho^2 m_0} \right) \\
& + \left(2\rho^2 l_{bd} \gamma^{CB} (2H_{[N|B]}^d - L_{[N|B]}^d) L_{[C|M]}^a + \frac{1}{\rho} \delta^a{}_b \gamma_{DA} \tilde{\gamma}^{(DC)} \rho_{,C} \bar{Q}^A{}_{MN} \right. \\
& \left. - 2\delta^a{}_b \nabla_{[N} \left(\frac{1}{\rho} \gamma_{|D|M]} \tilde{\gamma}^{(DC)} \rho_{,C} \right) \right) \left(\frac{q^b \zeta^M u^N}{2\rho^2 m_0} \right) \\
& + \left(\rho \tilde{\gamma}^{(CB)} \rho_{,B} l_{bc} L_{CM}^a + \rho l_{bd} \tilde{\gamma}^{(BN)} (2H_{MB}^d - L_{MB}^d) \delta^a{}_c \right)
\end{aligned} \tag{15.24}$$

$$\begin{aligned}
& \times \left(\frac{q^b}{2\rho^2 m_0} \right) \left(\zeta^c u^M - \zeta^M \left(\frac{q^c}{2\rho^2 m_0} \right) \right) \\
& + \left(\tilde{\nabla}_b L_{BC}^a - \overline{\nabla}_C \left(\frac{1}{\rho} \gamma_{BD} \tilde{\gamma}^{(DN)} \rho_{,N} \right) \delta^a_b + \rho^2 l_{db} \gamma^{DM} L_{MC}^a L_{DB}^d \right. \\
& - \frac{1}{\rho^2} \gamma_{DC} \tilde{\gamma}^{(DE)} \rho_{,E} \gamma_{BN} \tilde{\gamma}^{(NM)} \rho_{,M} \delta^a_b \left. \right) u^B \left(\zeta^b u^C - \zeta^C \left(\frac{q^b}{2\rho^2 m_0} \right) \right) \\
& + \left(-\frac{1}{\rho} \gamma_{BD} \tilde{\gamma}^{(DC)} \rho_{,C} C^a_{bc} + 2\rho \gamma_{CN} \tilde{\gamma}^{(NM)} \rho_{,M} g^{DC} L_{DB}^d l_{d[b} \delta^a_{c]} \right) u^B \left(\frac{\zeta^b q^c}{2\rho^4 m_0} \right) \\
& - 2\gamma_{CD} \tilde{\gamma}^{(DM)} \rho_{,M} \tilde{\gamma}^{(CB)} \rho_{,B} l_{b[c} \delta^a_{d]} \left(\frac{q^b \zeta^d q^c}{4\rho^2 m_0^2} \right) \\
& + \left(2\overline{\nabla}_{[M} L_{|B|N]}^a + L_{BC}^a \overline{Q}^C_{MN} + \frac{2}{\rho} \gamma_{BD} \tilde{\gamma}^{(DC)} \rho_{,C} H_{MN}^a \right. \\
& \quad \left. + \frac{2}{\rho} \gamma_{C[M} L_{|B|N]}^a \tilde{\gamma}^{(DC)} \rho_{,C} \right) u^B \zeta^M u^N \\
& - \frac{1}{2\rho} \tilde{Q}^a_{mb} \gamma_{BD} \tilde{\gamma}^{(DC)} \rho_{,C} u^B v^m \left(\frac{q^b}{m_0} \right) \\
& - \tilde{Q}^N_{MZ} u^Z v^M \left(-\frac{1}{\rho^3} \rho_{,N} \left(\frac{q^a}{m_0} \right) + L_{BN}^a u^B + \frac{1}{2\rho^3} \gamma_{DN} \tilde{\gamma}^{(DC)} \rho_{,C} \left(\frac{q^c}{m_0} \right) \right) \\
& - \frac{1}{\rho} \gamma_{ZM} \tilde{\gamma}^{(ZP)} \rho_{,P} \left(\tilde{\nabla}_b \left(\frac{q^a}{2\rho^2 m_0} \right) + \frac{1}{\rho} \gamma_{BD} \tilde{\gamma}^{(DC)} \rho_{,C} u^B \delta^a_b \right) \\
& \quad \times \left(v^M \left(\frac{q^b}{2\rho^2 m_0} \right) - v^b u^M \right) \\
& - 2(H_{MN}^n - L_{MN}^n) \left(\tilde{\nabla}_n \left(\frac{q^a}{2\rho^2 m_0} \right) + \frac{1}{\rho} \gamma_{BD} \tilde{\gamma}^{(DC)} \rho_{,C} u^B \delta^a_n \right) v^M u^N \\
& - 2\mu \rho \tilde{\gamma}^{(NM)} \rho_{,M} k_{bc} \left(-\frac{1}{\rho^2} \left(\frac{q^a}{m_0} \right) \rho_{,N} + L_{CN}^a u^C \right. \\
& \quad \left. + \frac{1}{2\rho^3} \gamma_{DN} \tilde{\gamma}^{(DC)} \rho_{,C} \left(\frac{q^a}{m_0} \right) \right) \left(\frac{v^b q^c}{2\rho^2 m_0} \right) \\
& - (2\rho^3 l_{bd} \gamma^{NB} H_{CB}^d + \rho^2 (l_{bd} \gamma^{NB} + l_{db} \gamma^{BN}) L_{BC}^d) \\
& \quad \times \left(-\frac{1}{\rho^3} \left(\frac{q^a}{m_0} \right) \rho_{,N} + L_{CN}^a u^C + \frac{1}{2\rho^3} \gamma_{DN} \tilde{\gamma}^{(DC)} \rho_{,C} \left(\frac{q^a}{m_0} \right) \right) \\
& \quad \times \left(v^C \left(\frac{q^b}{2\rho^2 m_0} \right) - v^b u^C \right)
\end{aligned}$$

(15.24_{cont.})

$$\begin{aligned}
& + \frac{1}{2\rho^3} \gamma_{BD} \tilde{\gamma}^{(DC)} \rho_{,C} v^B \left(\frac{q^a}{m_0} \right) + L_{BN}^a v^B u^N \\
& + \frac{1}{\rho} \gamma_{DN} \tilde{\gamma}^{(DC)} \rho_{,C} v^a u^N \\
& = 0
\end{aligned} \tag{15.24cont.}$$

where in both Eqs (15.23–24) $\overline{Q}^A_{MN}$ is a torsion on $E \times M$ for a connection $\overline{\omega}^A_B$ defined here, $\overline{R}^A_{BCD}$ is a tensor of a curvature for this connection, $\overline{\nabla}_N$ is a covariant derivative for this connection. A field ρ is defined on $E \times M$, $\rho = \rho(x, y)$.

$\tilde{\nabla}_n$ is a covariant derivative with respect to a connection on H (a nonsymmetric connection on a group manifold H), $\tilde{Q}^a_{bc}$ is a torsion of this connection, $\tilde{R}^a_{bcd}$ a tensor of a curvature. H^a_{MN} is a curvature of a connection ω on fibre bundle defined over base manifold $E \times M$, L^a_{MN} is defined by Eq. (5.13). We have

$$u^{\tilde{A}} = (u^A, u^a) \tag{15.25}$$

$$v^{\tilde{A}} = (v^A, v^a). \tag{15.26}$$

$$\zeta^{\tilde{A}} = (\zeta^A, \zeta^a) \tag{15.27}$$

and of course

$$v^A = \frac{d\zeta^A}{d\tau} \tag{15.28}$$

$$v^a = \frac{d\zeta^a}{d\tau} \tag{15.29}$$

$\tilde{\gamma}^{(AB)}$ is defined by Eq. (5.12). In this way we have geodetic deviation equation on P . Moreover, we need a dimensional reduction procedure to obtain them in terms of Yang–Mills’ and Higgs’ fields. It is easy to see that

$$\frac{d}{d\tau} q^b = 0 \tag{15.30}$$

or

$$q^b = \text{const.} \tag{15.30a}$$

Let us consider Eq. (15.23) according to sections 5 and 6. In this way one gets for $A = \alpha$ (i.e. for space-time E):

$$\begin{aligned}
& u^\beta \overline{\nabla}_\beta v^\alpha + \overline{R}^\alpha_{\beta} u^\beta \zeta^\mu u^\nu - \overline{Q}^\nu_{\mu\beta} (\overline{\nabla}_\nu u^\alpha) \zeta^\mu u^\beta \\
& + \left(\frac{q^b}{2\rho^2 m_0} \right) l_{bd} g^{\alpha\beta} \left[(2H^d_{\nu\beta} - L^d_{\nu\beta}) v^\nu + \left(2 \overset{\text{gauge}}{\nabla}_\beta \Phi^d_{\tilde{n}} - L^d_{\beta\tilde{n}} \right) v^{\tilde{n}} \right] \\
& - \rho^2 v^n l_{dn} g^{\delta\alpha} \left(L^d_{\delta\beta} u^\beta + L^d_{d\tilde{b}} u^{\tilde{b}} \right) - \left(\frac{q^n}{2\rho^2 m_0} \right) \tilde{g}^{(\alpha\beta)} \rho_{,\beta} l_{bn} v^b
\end{aligned} \tag{15.31}$$

$$\begin{aligned}
& + 2\rho^2 \left[\left(l_{cd} g^{\alpha\omega} \left(2H_{\omega[\mu}^d - L_{\omega[\mu}^d \right) L_{\nu]\beta}^c + l_{db} g^{\delta\alpha} L_{\delta\beta}^d H_{\mu\nu}^b \right) \zeta^\mu u^\nu u^\beta \right. \\
& + \left(\frac{1}{2} l_{cd} g^{\alpha\omega} \left(2 \overset{\text{gauge}}{\nabla}_\omega \Phi_{\tilde{m}}^d - L_{\omega\tilde{m}}^d \right) L_{\nu\beta}^c - l_{db} g^{\delta\alpha} L_{\delta\beta}^d \overset{\text{gauge}}{\nabla}_\nu \Phi_{\tilde{m}}^b \right) \zeta^{\tilde{m}} u^\nu u^\beta \\
& + \left(\frac{1}{2} l_{cd} g^{\alpha\omega} \left(2H_{\omega\mu}^d - L_{\omega\mu}^d \right) L_{\tilde{n}\beta}^c + l_{db} g^{\delta\alpha} L_{\delta\beta}^d \overset{\text{gauge}}{\nabla}_\mu \Phi_{\tilde{n}}^b \right) \zeta^\mu u^{\tilde{n}} u^\beta \\
& + \left(\frac{1}{2} l_{cd} g^{\alpha\omega} \left(2H_{\omega\mu}^d - L_{\omega\mu}^d \right) L_{\nu\tilde{b}}^c + l_{db} g^{\delta\alpha} L_{\delta\tilde{b}}^d H_{\mu\nu}^b \right) \zeta^\mu u^\nu u^{\tilde{b}} \\
& + \left(\frac{1}{2} l_{cd} g^{\alpha\omega} \left(2 \overset{\text{gauge}}{\nabla}_\omega \Phi_{\tilde{m}}^d - L_{\omega\tilde{m}}^d \right) L_{\tilde{n}\tilde{b}}^c + l_{db} g^{\delta\alpha} L_{\delta\tilde{b}}^d H_{\tilde{m}\tilde{n}}^b \right) \zeta^{\tilde{m}} u^{\tilde{n}} u^{\tilde{b}} \\
& + \left(\frac{1}{2} l_{cd} g^{\alpha\omega} \left(2 \overset{\text{gauge}}{\nabla}_\omega \Phi_{\tilde{m}}^d - L_{\omega\tilde{m}}^d \right) L_{\nu\tilde{b}}^c - l_{db} g^{\delta\alpha} L_{\delta\tilde{b}}^d \overset{\text{gauge}}{\nabla}_\nu \Phi_{\tilde{m}}^b \right) \zeta^{\tilde{m}} u^\nu u^{\tilde{b}} \\
& + \left(\frac{1}{2} l_{cd} g^{\alpha\omega} \left(2H_{\omega\mu}^d - L_{\omega\mu}^d \right) L_{\tilde{n}\tilde{b}}^c + l_{db} g^{\delta\alpha} L_{\delta\tilde{b}}^d \overset{\text{gauge}}{\nabla}_\mu \Phi_{\tilde{n}}^b \right) \zeta^\mu u^{\tilde{n}} u^{\tilde{b}} \\
& + \left. \left(\frac{1}{2} l_{cd} g^{\alpha\omega} \left(2 \overset{\text{gauge}}{\nabla}_\omega \Phi_{\tilde{m}}^d - L_{\omega\tilde{m}}^d \right) L_{\tilde{n}\beta}^c + l_{db} g^{\delta\alpha} L_{\delta\beta}^d H_{\tilde{m}\tilde{n}}^b \right) \zeta^{\tilde{m}} u^{\tilde{n}} u^\beta \right] \\
& + \frac{q^b}{2\rho^2 m_0} \left\{ \left[2\overline{\nabla}_{[\mu} \left(\rho^2 l_{db} g^{\alpha\beta} \left(2H_{\nu]\beta}^d - L_{\nu]\beta}^d \right) \right) \right. \right. \\
& + \rho^2 l_{db} g^{\alpha\beta} \left(2H_{\gamma\beta}^d - L_{\gamma\beta}^d \right) \overline{Q}^\gamma{}_{\mu\nu} - \rho \tilde{g}^{(\alpha\beta)} \rho_{,\beta} l_{bc} H_{\mu\nu}^c \\
& + 2\rho l_{bd} g^{\alpha\beta} \left(2H_{[\mu|\beta]}^d - L_{[\mu|\beta]}^d \right) g_{|\delta|\nu]} \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} \left. \right] \zeta^\mu u^\nu \\
& + \left[2\tilde{\nabla}_{[\tilde{m}} \left(\rho^2 l_{db} g^{\alpha\beta} \left(-2 \overset{\text{gauge}}{\nabla}_\beta \Phi_{\tilde{n}}^d + L_{\beta\tilde{n}}^d \right) \right) \right. \\
& + \rho^2 l_{db} g^{\alpha\beta} \left(2H_{\gamma\beta}^d - L_{\gamma\beta}^d \right) Q_{\tilde{m}\tilde{n}}^\gamma \\
& - \rho l_{bd} g^{\alpha\beta} \left(2 \overset{\text{gauge}}{\nabla}_\beta \Phi_{\tilde{m}}^d - L_{\beta\tilde{m}}^d \right) g_{\tilde{d}\tilde{n}} \tilde{g}^{(\tilde{d}\tilde{c})} \rho_{,\tilde{c}} \left. \right] \zeta^{\tilde{m}} u^{\tilde{n}} \\
& + \left[\tilde{\nabla}_{\tilde{m}} \left(\rho^2 l_{db} g^{\alpha\beta} \left(2H_{\nu\beta}^d - L_{\nu\beta}^d \right) \right) \right. \\
& + \rho^2 l_{db} g^{\alpha\beta} \left[\left(2H_{\gamma\beta}^d - L_{\gamma\beta}^d \right) Q_{\tilde{m}\nu}^\gamma + \left(-2 \overset{\text{gauge}}{\nabla}_\beta \Phi_{\tilde{c}}^d + L_{\beta\tilde{c}}^d \right) Q_{\tilde{m}\nu}^{\tilde{c}} \right] \\
& + \rho \tilde{g}^{(\alpha\beta)} \rho_{,\beta} l_{bc} \overset{\text{gauge}}{\nabla}_\nu \Phi_{\tilde{m}}^c \\
& + \rho l_{bd} g^{\alpha\beta} \left(- \overset{\text{gauge}}{\nabla}_\beta \Phi_{\tilde{m}}^d + L_{\beta\tilde{m}}^d \right) g_{\delta\nu} \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} \left. \right] \zeta^{\tilde{m}} u^\nu
\end{aligned} \tag{15.31cont.}$$

$$\begin{aligned}
& + \left[\bar{\nabla}_\mu \left(\rho^2 l_{db} g^{\alpha\beta} \left(-2 \overset{\text{gauge}}{\nabla}_\beta \Phi_{\tilde{n}}^d + L_{\beta\tilde{n}}^d \right) \right) \right. \\
& \quad + \rho^2 l_{db} g^{\alpha\beta} \left[(2H_{\gamma\beta}^d - L_{\gamma\beta}^d) Q^\gamma_{\mu\tilde{n}} \right. \\
& \quad \left. + \left(-2 \overset{\text{gauge}}{\nabla}_\beta \Phi_{\tilde{c}}^d + L_{\beta\tilde{c}}^d \right) Q_{\mu\tilde{n}}^{\tilde{c}} \right] - \rho \tilde{g}^{(\alpha\beta)} \rho_{,\beta} l_{bc} \overset{\text{gauge}}{\nabla}_\mu \Phi_{\tilde{n}}^c \\
& \quad \left. + 2\rho l_{bd} g^{\alpha\beta} (2H_{\beta\mu}^d - L_{\beta\mu}^d) g_{\tilde{d}\tilde{n}} \tilde{g}^{(\tilde{n}\tilde{c})} \rho_{,c} \right] \zeta^\mu u^{\tilde{n}} \Big\} \\
& + \left\{ \rho l_{bd} g^{\alpha\omega} g_{\beta\delta} \tilde{g}^{(\delta\zeta)} \rho_{,\zeta} (2H_{\mu\omega}^d - L_{\mu\omega}^d) - \bar{\nabla}_\mu (\rho^2 l_{db} g^{\delta\alpha} L_{\delta\beta}^d) \right. \\
& \quad \left. - \rho \tilde{g}^{(\alpha\omega)} \rho_{,\omega} l_{cd} L_{\beta\mu}^c \right\} u^\beta \left(\zeta^\mu \frac{q^b}{2\rho^2 m_0} - \zeta^b u^\mu \right) \\
& + \left(2\rho^4 l_{d[b} l_{|e|f]} g^{\delta\alpha} \left(g^{\zeta\gamma} L_{\delta\gamma}^d L_{\zeta\beta}^e + \frac{1}{r^2} g^{\tilde{z}\tilde{c}} L_{\delta\tilde{c}}^d L_{\tilde{z}\beta}^e \right) \right. \\
& \quad \left. + \tilde{g}^{(\alpha\omega)} \rho_{,\omega} g_{\beta\delta} \tilde{g}^{(\delta\zeta)} \rho_{,\zeta} l_{[bf]} + \frac{\rho^2}{2} l_{dp} g^{\delta\alpha} L_{\delta\beta}^d C_{bf}^p \right) u^\beta \left(\frac{\zeta^b q^f}{2\rho^2 m_0} \right) \\
& + \left(2\rho^4 l_{d[b} l_{|e|f]} g^{\delta\alpha} \left(g^{\zeta\gamma} L_{\delta\gamma}^d L_{\zeta\tilde{b}}^e + \frac{1}{r^2} g^{\tilde{z}\tilde{c}} L_{\delta\tilde{v}}^d L_{\tilde{z}\tilde{b}}^e \right) \right. \\
& \quad \left. + \mu \tilde{g}^{(\alpha\omega)} \rho_{,\omega} g^{\tilde{b}\tilde{d}} \tilde{g}^{(\tilde{d}\tilde{z})} \rho_{,\tilde{z}} k_{bf} + \frac{\rho^2}{2} l_{dp} g^{\delta\alpha} L_{\delta\tilde{b}}^d C_{bf}^p \right) u^{\tilde{b}} \left(\frac{\zeta^b q^f}{2\rho^2 m_0} \right) \\
& + \left\{ \bar{\nabla}_\mu \left(\rho \tilde{g}^{(\alpha\beta)} \rho_{,\beta} \right) l_{ba} + \tilde{\nabla}_a (\rho^2 l_{bd} g^{\alpha\beta} (2H_{\mu\beta}^d - L_{\mu\beta}^d)) \right. \\
& \quad - \rho^4 l_{dc} l_{bf} g_{\delta\alpha} g^{\gamma\beta} L_{\delta\gamma}^d (2H_{\mu\beta}^f - L_{\mu\beta}^f) \\
& \quad - \frac{\rho^4}{r^2} l_{dc} l_{bf} g^{\delta\alpha} g^{\tilde{c}\tilde{b}} L_{\delta\tilde{c}}^d \left(2 \overset{\text{gauge}}{\nabla}_\mu \Phi_{\tilde{b}}^f - L_{\mu\tilde{b}}^f \right) \\
& \quad \left. - \tilde{g}^{(\alpha\zeta)} \rho_{,\zeta} g_{\delta\mu} \tilde{g}^{(\delta\nu)} \rho_{,\nu} l_{bc} \right\} \left(\frac{q^b}{2\rho^2 m_0} \right) \left(\zeta^a u^\mu - \zeta^\mu \frac{q^a}{2\rho^2 m_0} \right) \\
& + \left\{ \hat{\bar{\nabla}}_{\tilde{m}} \left(\rho \tilde{g}^{(\alpha\beta)} \rho_{,\beta} \right) l_{ba} - \tilde{\nabla}_a \left(\rho^2 l_{bd} g^{\alpha\beta} \left(2 \overset{\text{gauge}}{\nabla}_\beta \Phi_{\tilde{m}}^d - L_{\beta\tilde{m}}^d \right) \right) \right. \\
& \quad + \rho^4 l_{dc} l_{bf} g^{\delta\alpha} g_{\gamma\beta} L_{\delta\gamma}^d \left(2 \overset{\text{gauge}}{\nabla}_\beta \Phi_{\tilde{m}}^f - L_{\beta\tilde{m}}^f \right) \\
& \quad \left. - \frac{\rho^4}{r^2} l_{dc} l_{bf} g^{\delta\alpha} \tilde{g}^{(\tilde{c}\tilde{b})} L_{\delta\tilde{c}}^d \left(2H_{\tilde{m}\tilde{b}}^f - L_{\tilde{m}\tilde{b}}^f \right) \right\}
\end{aligned} \tag{15.31_{cont.}}$$

$$\begin{aligned}
& -r^2 \tilde{g}^{(\alpha\zeta)} \rho_{,\zeta} g_{\tilde{d}\tilde{m}} \tilde{g}^{(\tilde{d}\tilde{n})} \rho_{,\tilde{n}} l_{bc} \left\} \left(\frac{q^b}{2\rho^2 m_0} \right) \left(\zeta^a u^{\tilde{m}} - \zeta^{\tilde{m}} \frac{q^a}{2\rho^2 m_0} \right) \\
& + \left(\rho \tilde{g}^{(\alpha\beta)} \rho_{,\beta} l_{bk} C_{ac}^k + 2\rho^3 \tilde{g}^{(\gamma\beta)} \rho_{,\beta} g^{\delta\alpha} L_{\delta\gamma}^d l_{d[a} l_{b|c]} \right. \\
& \quad \left. + \frac{2\rho^3}{r^2} \tilde{g}^{(\alpha\beta)} \rho_{,\beta} g^{\tilde{d}\tilde{a}} L_{\tilde{d}\tilde{c}}^d l_{d[a} l_{b|c]} \right) \left(\frac{q^b \zeta^a q^c}{4\rho^2 m_0^2} \right) \\
& - \left(2\rho^2 l_{bd} g^{\alpha\beta} H_{\gamma\beta}^d + \rho^2 (l_{bd} g^{\nu\beta} + l_{db} g^{\beta\nu}) L_{\beta\gamma}^d \right. \\
& \quad \times \left(\bar{\nabla}_\nu u^\alpha + \frac{1}{2} l_{bf} g^{\alpha\beta} (2H_{\nu\beta}^d - L_{\nu\beta}^d) \left(\frac{q^f}{m_0} \right) \right) \left(u^\gamma \frac{q^b}{2\rho^2 m_0} - v^b u^\gamma \right) \\
& - \left(\frac{2}{r^2} \rho^2 l_{bd} \tilde{g}^{(\tilde{n}\tilde{b})} \bar{\nabla}_\gamma^{\text{gauge}} \Phi_{\tilde{b}}^d + \frac{\rho^2}{r^2} (l_{bd} g^{\tilde{n}\tilde{b}} + l_{db} g^{\tilde{b}\tilde{n}}) L_{\gamma\tilde{b}}^d \right. \\
& \quad \times \left(\hat{\bar{\nabla}}_{\tilde{n}} u^\alpha + \frac{1}{2} l_{bf} g^{\alpha\beta} \left(2 \bar{\nabla}_\beta^{\text{gauge}} \Phi_{\tilde{n}}^d - L_{\beta\tilde{n}}^d \right) \right) \left(\frac{q^f}{m_0} \right) \left(u^\gamma \frac{q^b}{2\rho^2 m_0} - v^b u^\gamma \right) \\
& - 2\mu \rho \tilde{g}^{(\nu\beta)} \rho_{,\beta} k_{bc} \left(\bar{\nabla}_\nu u^\alpha + \frac{1}{2} (l_{df} g^{\alpha\gamma} (2H_{\nu\gamma}^d - L_{\nu\gamma}^d) \left(\frac{q^f}{m_0} \right)) \right) \left(\frac{q^c v^b}{2\rho^2 m_0} \right) \\
& - \frac{2\mu\rho}{r^2} \tilde{g}^{(\tilde{n}\tilde{b})} \rho_{,\tilde{b}} k_{bc} \left(\hat{\bar{\nabla}}_{\tilde{n}} u^\alpha - \frac{1}{2} (l_{df} g^{\alpha\gamma} (2 \bar{\nabla}_\gamma^{\text{gauge}} \Phi_{\tilde{n}}^d - L_{\gamma\tilde{n}}^d) \left(\frac{q^f}{m_0} \right)) \right) \left(\frac{q^c v^b}{2\rho^2 m_0} \right) \\
& + (2H_{\mu\nu}^n - L_{\mu\nu}^n) \left(\rho^2 l_{dn} g^{\delta\alpha} (L_{\delta\beta}^d u^\beta + L_{\delta\tilde{b}}^d u^{\tilde{b}}) + \frac{1}{2\rho^2} \tilde{g}^{(\alpha\beta)} \rho_{,\beta} l_{bn} \left(\frac{q^b}{m_0} \right) \right) v^\mu u^\nu \\
& + (2H_{\tilde{m}\tilde{n}}^n - L_{\tilde{m}\tilde{n}}^n) \left(\rho^2 l_{dn} g^{\delta\alpha} (L_{\delta\beta}^d u^\beta + L_{\delta\tilde{b}}^d u^{\tilde{b}}) + \frac{1}{2\rho^2} \tilde{g}^{(\alpha\beta)} \rho_{,\beta} l_{bn} \left(\frac{q^b}{m_0} \right) \right) v^{\tilde{m}} u^{\tilde{n}} \\
& + 2 \left(\bar{\nabla}_\mu^{\text{gauge}} \Phi_{\tilde{n}}^n - L_{\mu\tilde{n}}^n \right) \left(\rho^2 l_{dn} g^{\delta\alpha} (L_{\delta\beta}^d u^\beta + L_{\delta\tilde{b}}^d u^{\tilde{b}}) + \frac{1}{2\rho^2} \tilde{g}^{(\alpha\beta)} \rho_{,\beta} l_{bn} \left(\frac{q^b}{m_0} \right) \right) v^\mu u^{\tilde{n}} \\
& - 2 \left(\bar{\nabla}_\nu^{\text{gauge}} \Phi_{\tilde{m}}^n - L_{\nu\tilde{m}}^n \right) \left(\rho^2 l_{dn} g^{\delta\alpha} (L_{\delta\beta}^d u^\beta + L_{\delta\tilde{b}}^d u^{\tilde{b}}) + \frac{1}{2\rho^2} \tilde{g}^{(\alpha\beta)} \rho_{,\beta} l_{bn} \left(\frac{q^b}{m_0} \right) \right) v^{\tilde{m}} u^\mu \\
& - g_{\zeta\mu} \tilde{g}^{(\alpha\gamma)} \rho_{,\gamma} \left(\rho l_{bm} \gamma^{\delta\zeta} (L_{\delta\beta}^b u^\beta + L_{\delta\tilde{b}}^b u^{\tilde{b}}) \right. \\
& \quad \left. + \frac{1}{2\rho^2} \tilde{g}^{(\zeta\beta)} \rho_{,\beta} l_{bm} \left(\frac{q^b}{m_0} \right) \right) \left(v^\mu \left(\frac{q^m}{2\rho^2 m_0} \right) - v^m u^\mu \right) \\
& - g_{\tilde{z}\tilde{m}} \tilde{g}^{(\alpha\gamma)} \rho_{,\gamma} \left(\frac{\rho}{r^2} l_{bm} g^{\tilde{d}\tilde{z}} (L_{\tilde{d}\beta}^b u^\beta + L_{\tilde{d}\tilde{b}}^b u^{\tilde{b}}) + \frac{1}{2r^2 \rho^2} \tilde{g}^{(\tilde{z}\tilde{b})} \rho_{,\tilde{b}} l_{bm} \left(\frac{q^b}{m_0} \right) \right)
\end{aligned} \tag{15.31cont.}$$

$$\begin{aligned}
& \times \left(v^{\tilde{z}} \left(\frac{q^m}{2\rho^2 m_0} \right) - v^m u^{\tilde{z}} \right) - \frac{1}{2\rho^2} \tilde{Q}^n{}_{mb} \left(\rho^2 l_{dn} g^{\delta\alpha} \left(L_{\delta\beta}^d u^\beta + L_{\delta\tilde{b}}^d u^{\tilde{b}} \right) \right. \\
& \left. - \frac{1}{2\rho} \tilde{g}^{(\alpha\beta)} l_{pn} \left(\frac{q^p}{m_0} \right) \right) v^m \left(\frac{q^b}{m_0} \right) \\
& = 0
\end{aligned} \tag{15.31}_{\text{cont.}}$$

where $H_{m\tilde{n}}^a$ are given by Eq. (4.55) and $\overset{\text{gauge}}{\nabla}$ means a derivative with respect to the connection $\tilde{\omega}_E$ on a bundle P (over a space-time E). $\overline{Q}^\nu{}_{\mu\beta}$ means a torsion on the space-time E with respect to a connection $\overline{\omega}^\alpha{}_\beta$ and $\overline{\nabla}_\mu$ a covariant derivative with respect to this connection. $\overline{R}^\alpha{}_{\beta\mu\nu}$ means a curvature tensor calculated for this connection. $\widehat{\nabla}_{\tilde{m}}$ means a covariant derivative with respect to a connection $\widehat{\omega}^{\tilde{m}}{}_{\tilde{n}}$ on M , $\tilde{\nabla}_a$ means a covariant derivative with respect to a connection $\tilde{\omega}_b^a$ on a group manifold H .

$H_{\mu\nu}^a$ is a curvature of a connection $\tilde{\omega}_E$ given by Eq. (4.53), $L_{\mu\nu}^a$ is given by Eq. (5.22), $L_{\beta\tilde{n}}^a, L_{\tilde{a}\tilde{b}}^a$ are given by Eq. (5.23) and Eq. (5.24), respectively.

Let us consider Eq. (15.23) for $A = \tilde{a}$ (i.e. for a manifold M). One gets

$$\begin{aligned}
& u^\beta \overline{\nabla}_\beta v^{\tilde{a}} + u^{\tilde{b}} \tilde{\nabla}_{\tilde{b}} v^{\tilde{a}} + \widehat{R}^{\tilde{a}}{}_{\tilde{b}\tilde{m}\tilde{n}} u^{\tilde{b}} \zeta^{\tilde{m}} u^{\tilde{n}} - \widehat{Q}^{\tilde{n}}{}_{\tilde{m}\tilde{b}} \widehat{\nabla}_{\tilde{n}} u^{\tilde{a}} \zeta^{\tilde{m}} u^{\tilde{b}} \\
& + \frac{1}{r^2} \left(\frac{q^b}{2\rho^2 m_0} \right) l_{bd} g^{\tilde{a}\tilde{b}} \left(2 \overset{\text{gauge}}{\nabla}_\mu \Phi_b^d - L_{\mu\tilde{b}}^d \right) v^\mu \\
& + \frac{1}{r^2} \left(\frac{q^b}{2\rho^2 m_0} \right) l_{bd} g^{\tilde{a}\tilde{b}} (2H_{\tilde{n}\tilde{b}}^b - L_{\tilde{n}\tilde{b}}^b) v^{\tilde{b}} \\
& - \frac{\rho^2}{r^2} v^n l_{dn} g^{\tilde{d}\tilde{a}} \left(L_{\tilde{d}\tilde{b}}^d u^{\tilde{b}} + L_{\tilde{d}\beta}^d u^\beta \right) - \left(\frac{q^n}{2\rho^2 m_0} \right) \tilde{g}^{(\tilde{a}\tilde{b})} \rho_{,\tilde{b}} l_{bn} v^b \\
& - \frac{2\rho^2}{r^2} \left(l_{cd} g^{\tilde{a}\tilde{p}} \left(2 \overset{\text{gauge}}{\nabla}_{[\mu} \Phi_{|\tilde{p}|}^d - L_{[\mu|\tilde{p}]}^d \right) L_{\nu]\beta}^d + l_{db} g^{\tilde{d}\tilde{a}} L_{\beta\tilde{d}}^d H_{\mu\nu}^b \right) u^\beta \zeta^\mu u^\nu \\
& + \frac{2\rho^2}{r^2} \left(l_{cd} g^{\tilde{a}\tilde{p}} \left(2H_{\tilde{p}[\tilde{m}}^d - L_{\tilde{p}[\tilde{m}}^d \right) L_{\tilde{N}]\beta}^c + l_{db} g^{\tilde{d}\tilde{a}} L_{\beta\tilde{d}}^d H_{\tilde{m}\tilde{n}}^b \right) u^\beta \zeta^{\tilde{m}} u^{\tilde{n}} \\
& + \frac{2\rho^2}{r^2} \left(l_{cd} g^{\tilde{a}\tilde{p}} \left(2H_{\tilde{p}[\tilde{m}}^d - L_{\tilde{p}[\tilde{m}}^d \right) L_{\tilde{n}]\tilde{b}}^c + l_{db} g^{\tilde{d}\tilde{a}} L_{\tilde{d}\tilde{b}}^d H_{\tilde{m}\tilde{n}}^b \right) u^{\tilde{b}} \zeta^{\tilde{m}} u^{\tilde{n}} \\
& - \frac{2\rho^2}{r^2} \left(\frac{1}{2} l_{cd} g^{\tilde{a}\tilde{p}} \left(2 \overset{\text{gauge}}{\nabla}_\mu \Phi_{\tilde{p}}^d - L_{\mu\tilde{p}}^d \right) L_{\tilde{n}\tilde{b}}^c - l_{db} g^{\tilde{d}\tilde{a}} L_{\tilde{d}\tilde{b}}^d \overset{\text{gauge}}{\nabla}_\mu \Phi_{\tilde{n}}^d \right) u^{\tilde{b}} \zeta^\mu u^{\tilde{n}} \\
& + \frac{2\rho^2}{r^2} \left(\frac{1}{2} l_{cd} g^{\tilde{a}\tilde{p}} \left(2H_{\tilde{p}\tilde{m}}^d - L_{\tilde{m}\tilde{p}}^d \right) L_{\nu\tilde{b}}^c - l_{db} g^{\tilde{d}\tilde{a}} L_{\tilde{d}\tilde{b}}^d \overset{\text{gauge}}{\nabla}_\nu \Phi_{\tilde{m}}^d \right) u^{\tilde{b}} \zeta^{\tilde{m}} u^\nu \\
& - \frac{2\rho^2}{r^2} \left(l_{cd} g^{\tilde{a}\tilde{p}} \left(2 \overset{\text{gauge}}{\nabla}_{[\mu} \Phi_{|\tilde{p}|}^d - L_{[\mu|\tilde{p}]}^d \right) L_{\nu]\tilde{b}}^c - l_{db} g^{\tilde{d}\tilde{a}} L_{\tilde{d}\tilde{b}}^d H_{\mu\nu}^d \right) u^{\tilde{b}} \zeta^\mu u^\nu
\end{aligned} \tag{15.32}$$

$$\begin{aligned}
& -\frac{2\rho^2}{r^2} \left(\frac{1}{2} l_{cd} g^{\tilde{a}\tilde{p}} \left(2 \nabla_\mu^{\text{gauge}} \Phi_{\tilde{p}}^d - L_{\tilde{p}\mu}^d \right) L_{\tilde{n}\beta}^c - l_{db} g^{\tilde{d}\tilde{a}} L_{\tilde{d}\beta}^d \nabla_\mu^{\text{gauge}} \Phi_{\tilde{n}}^b \right) u^\beta \zeta^\mu u^{\tilde{n}} \\
& + \frac{2\rho^2}{r^2} \left(\frac{1}{2} l_{cd} g^{\tilde{a}\tilde{p}} \left(2 H_{\tilde{p}\tilde{m}}^d - L_{\tilde{p}\tilde{m}}^d \right) L_{\nu\beta}^c - l_{db} g^{\tilde{d}\tilde{a}} L_{\tilde{d}\beta}^d \nabla_\nu^{\text{gauge}} \Phi_{\tilde{m}}^b \right) u^\beta \zeta^{\tilde{m}} u^\nu \\
& + \frac{q^b}{2m_0 r^2 \rho^2} \left[2 \overline{\nabla}_{[\mu} \left(\rho^2 l_{db} g^{\tilde{a}\tilde{b}} \left(2 \nabla_{\nu]}^{\text{gauge}} \Phi_{\tilde{b}}^d - L_{\nu]}^d \right) \right) \right. \\
& \quad \left. + \rho^2 l_{db} g^{\tilde{a}\tilde{b}} \left(2 \nabla_\gamma^{\text{gauge}} \Phi_{\tilde{b}}^d - L_{\gamma\tilde{b}}^d \right) \overline{Q}^\gamma_{\mu\nu} - \rho \tilde{g}^{(\tilde{a}\tilde{b})} \rho_{,\tilde{b}} l_{bc} H_{\mu\nu}^c \right] \zeta^\mu u^\nu \\
& + \frac{q^b}{2m_0 r^2 \rho^2} \left[\overline{\nabla}_\mu \left(\rho^2 l_{db} g^{\tilde{a}\tilde{b}} \left(2 H_{\tilde{n}\tilde{b}}^d - L_{\tilde{n}\tilde{b}}^d \right) \right) \right. \\
& \quad \left. - \rho \tilde{g}^{(\tilde{a}\tilde{b})} \rho_{,\tilde{b}} l_{bc} \nabla_\mu^{\text{gauge}} \Phi_{\tilde{n}}^c + \rho l_{bd} g^{\tilde{a}\tilde{b}} \left(2 \nabla_\mu^{\text{gauge}} \Phi_{\tilde{b}}^d - L_{\mu\tilde{b}}^d \right) g_{\tilde{d}\tilde{n}} \tilde{g}^{(\tilde{d}\tilde{c})} \rho_{\tilde{c}} \right] \zeta^\mu u^{\tilde{n}} \\
& + \frac{q^b}{2r^2 m_0 \rho^2} \left[2 \widehat{\nabla}_{[\tilde{m}} \left(\rho^2 l_{db} g^{\tilde{a}\tilde{b}} \left(2 H_{\tilde{n}\tilde{b}}^d - L_{\tilde{n}\tilde{b}}^d \right) \right) \right. \\
& \quad + \rho^2 l_{db} g^{\tilde{a}\tilde{b}} \left(2 H_{\tilde{c}\tilde{b}}^d - L_{\tilde{c}\tilde{b}}^d \right) \widehat{Q}^{\tilde{c}}_{\tilde{m}\tilde{n}} - \rho \tilde{g}^{(\tilde{a}\tilde{b})} \rho_{,\tilde{b}} l_{bc} H_{\tilde{m}\tilde{n}}^c \\
& \quad \left. + 2 \rho l_{bd} g^{\tilde{a}\tilde{b}} \left(2 H_{[\tilde{m}|\tilde{b}]}^d - L_{[\tilde{m}|\tilde{b}]}^d \right) g_{|\tilde{d}|\tilde{n}} \tilde{g}^{(\tilde{d}\tilde{c})} \rho_{,\tilde{c}} \right] \zeta^{\tilde{m}} u^{\tilde{n}} \tag{15.32cont.} \\
& + \frac{1}{r^2} \left\{ \rho l_{bd} g^{\tilde{a}\tilde{p}} g_{\beta\delta} \tilde{g}^{(\delta\zeta)} \rho_{,\zeta} \left(2 \nabla_\mu^{\text{gauge}} \Phi_{\tilde{p}}^d - L_{\mu\tilde{p}}^d \right) - \overline{\nabla}_\mu \left(\rho^2 l_{db} g^{\tilde{d}\tilde{a}} L_{\tilde{d}\beta}^d \right) \right. \\
& \quad \left. - \rho \tilde{g}^{(\tilde{a}\tilde{p})} \rho_{,\tilde{p}} l_{cd} L_{\beta\mu}^c \right\} u^\beta \left(\zeta^\mu \left(\frac{q^b}{2\rho^2 m_0} \right) - \zeta^b u^\mu \right) \\
& + \frac{1}{r^2} \left\{ \rho l_{bd} g^{\tilde{a}\tilde{p}} g_{\tilde{b}\tilde{d}} \tilde{g}^{(\tilde{d}\tilde{z})} \rho_{,\tilde{z}} \left(2 \nabla_\mu^{\text{gauge}} \Phi_{\tilde{p}}^d - L_{\mu\tilde{p}}^d \right) - \overline{\nabla}_\mu \left(\rho^2 l_{db} g^{\tilde{d}\tilde{a}} L_{\tilde{d}\tilde{b}}^d \right) \right. \\
& \quad \left. + \rho \tilde{g}^{(\tilde{a}\tilde{p})} \rho_{,\tilde{p}} l_{cd} L_{\mu\tilde{b}}^c \right\} u^{\tilde{b}} \left(\zeta^\mu \left(\frac{q^b}{2\rho^2 m_0} \right) - \zeta^b u^\mu \right) \\
& + \frac{1}{r^2} \left\{ \rho l_{bd} g^{\tilde{a}\tilde{p}} g_{\tilde{b}\tilde{d}} \tilde{g}^{(\tilde{d}\tilde{z})} \rho_{,\tilde{z}} \left(2 H_{\tilde{m}\tilde{p}}^d - L_{\tilde{m}\tilde{p}}^d \right) - \widehat{\nabla}_{\tilde{m}} \left(\rho^2 l_{db} g^{\tilde{d}\tilde{a}} L_{\tilde{d}\tilde{b}}^d \right) \right. \\
& \quad \left. - \rho \tilde{g}^{(\tilde{a}\tilde{p})} \rho_{,\tilde{p}} l_{cd} L_{\tilde{b}\tilde{m}}^c \right\} u^{\tilde{b}} \left(\zeta^{\tilde{m}} \left(\frac{q^b}{2\rho^2 m_0} \right) - \zeta^b u^{\tilde{m}} \right) \\
& + \frac{1}{r^2} \left(2 \rho^4 l_{d[b} l_{e|f]} g^{\tilde{d}\tilde{a}} \left(\frac{1}{r^2} g^{\tilde{z}\tilde{c}} L_{\tilde{d}\tilde{c}}^d L_{\tilde{z}\beta}^e + g^{\zeta\gamma} L_{\tilde{d}\gamma}^d L_{\zeta\beta}^e \right) \right. \\
& \quad \left. + \tilde{g}^{(\tilde{a}\tilde{p})} \rho_{,\tilde{p}} g_{\tilde{b}\tilde{d}} \tilde{g}^{(\tilde{d}\tilde{z})} \rho_{,\tilde{z}} l_{[bf]} + \frac{\rho^2}{2} l_{dp} g^{\tilde{d}\tilde{a}} L_{\tilde{d}\beta}^d C^p_{bf} \right) u^\beta \left(\frac{\zeta^b q^f}{2\rho^2 m_0} \right)
\end{aligned}$$

$$\begin{aligned}
& + \frac{1}{r^2} \left\{ \bar{\nabla}_\mu \left(\rho \tilde{g}^{(\tilde{a}\tilde{b})} \rho_{,\tilde{b}} \right) l_{ba} + \tilde{\nabla}_a \left(\rho^2 l_{bd} g^{\tilde{a}\tilde{b}} \left(2 \nabla_\mu^{\text{gauge}} \Phi_b^d - L_{\mu\tilde{b}}^d \right) \right) \right. \\
& \quad - \rho^4 l_{dc} l_{bf} g^{\tilde{d}\tilde{a}} \left(\frac{1}{r^2} g^{\tilde{c}\tilde{b}} L_{\tilde{d}\tilde{c}}^d \left(2 \nabla_\mu^{\text{gauge}} \Phi_b^f - L_{\mu\tilde{b}}^f \right) \right. \\
& \quad \left. \left. + g^{\gamma\beta} L_{\tilde{d}\gamma}^d \left(2 H_{\mu\beta}^f - L_{\mu\beta}^f \right) \right) \right. \\
& \quad \left. - \tilde{g}^{(\tilde{a}\tilde{z})} \rho_{,\tilde{z}} g_{\delta\mu} \tilde{g}^{(\delta\nu)} \rho_{,\nu} l_{bc} \right\} \left(\frac{q^b}{2\rho^2 m_0} \right) \left(\zeta^a u^\mu - \zeta^\mu \left(\frac{q^a}{2\rho^2 m_0} \right) \right) \\
& + \frac{1}{r^2} \left\{ \widehat{\nabla}_{\tilde{m}} \left(\rho^2 \tilde{g}^{(\tilde{a}\tilde{b})} \rho_{,\tilde{b}} \right) l_{ba} + \tilde{\nabla}_a \left(\rho^2 l_{bd} g^{\tilde{a}\tilde{b}} \left(2 H_{\tilde{m}\tilde{b}}^d - L_{\tilde{m}\tilde{b}}^d \right) \right) \right. \\
& \quad - \rho^2 l_{dc} l_{bf} g^{\tilde{d}\tilde{a}} \left(\frac{1}{r^2} g^{\tilde{c}\tilde{b}} L_{\tilde{d}\tilde{c}}^d \left(2 H_{\tilde{m}\tilde{b}}^f - L_{\tilde{m}\tilde{b}}^f \right) - g^{\gamma\beta} L_{\tilde{d}\gamma}^d \left(2 \nabla_\beta^{\text{gauge}} \Phi_{\tilde{m}}^f - L_{\beta\tilde{m}}^f \right) \right) \\
& \quad \left. - \tilde{g}^{(\tilde{a}\tilde{z})} \rho_{,\tilde{z}} g_{\tilde{d}\tilde{m}} \tilde{g}^{(\tilde{d}\tilde{n})} \rho_{,\tilde{n}} l_{bc} \right\} \left(\frac{q^b}{2\rho^2 m_0} \right) \left(\zeta^a u^{\tilde{m}} - \zeta^{\tilde{m}} \left(\frac{q^a}{2\rho^2 m_0} \right) \right) \\
& + \frac{1}{r^2} \left(\rho \tilde{g}^{(\tilde{a}\tilde{b})} \rho_{,\tilde{b}} l_{bk} C_{ac}^k \right. \\
& \quad \left. + 2\rho^3 g^{\tilde{d}\tilde{a}} \left(\frac{1}{r^2} L_{\tilde{d}\tilde{c}}^d \tilde{g}^{(\tilde{c}\tilde{b})} \rho_{,\tilde{b}} + L_{\tilde{d}\gamma}^d \tilde{g}^{(\gamma\beta)} \rho_{,\beta} \right) l_{d[a} l_{b|c]} \right) \left(\frac{q^b \zeta^a q^c}{4q^4 m_0^2} \right) \\
& - \left\{ 2\rho^2 l_{bd} \left[\frac{1}{r^2} \left(g^{\tilde{n}\tilde{b}} H_{\tilde{c}\tilde{b}}^d + \rho^2 \left(l_{bd} g^{\tilde{n}\tilde{b}} + l_{db} g^{\tilde{b}\tilde{n}} \right) L_{\tilde{b}\tilde{c}}^d \right) \right. \right. \\
& \quad \times \left(\widehat{\nabla}_{\tilde{n}} u^{\tilde{a}} + \frac{1}{r^2} l_{fd} g^{\tilde{a}\tilde{b}} \left(2 H_{\tilde{n}\tilde{b}}^d - L_{\tilde{n}\tilde{b}}^d \right) \left(\frac{q^f}{m_0} \right) \right) \\
& \quad \left. + \left(-g^{\nu\beta} \nabla_\beta^{\text{gauge}} \Phi_{\tilde{c}}^d + \rho^2 \left(l_{bd} g^{\nu\beta} + l_{db} g^{\beta\nu} \right) L_{\beta\tilde{c}}^d \right) \right. \\
& \quad \left. \times \left(\bar{\nabla}_\nu u^{\tilde{a}} + \frac{1}{r^2} l_{fd} g^{\tilde{a}\tilde{b}} \left(2 \nabla_\nu^{\text{gauge}} \Phi_b^d - L_{\nu\tilde{b}}^d \right) \left(\frac{q^f}{m_0} \right) \right) \right] \right\} \left(u^{\tilde{c}} \left(\frac{q^b}{2\rho^2 m_0} \right) - v^b u^{\tilde{c}} \right) \\
& - \left\{ 2\rho^2 l_{bd} \left[\frac{1}{r^2} \left(g^{\tilde{n}\tilde{b}} \nabla_\gamma^{\text{gauge}} \Phi_b^d + \rho^2 \left(l_{bd} g^{\tilde{n}\tilde{b}} + l_{db} g^{\tilde{b}\tilde{n}} \right) L_{\tilde{b}\beta}^d \right) \right. \right. \\
& \quad \times \left(\widehat{\nabla}_{\tilde{n}} u^{\tilde{a}} + \frac{1}{r^2} l_{fd} g^{\tilde{a}\tilde{b}} \left(2 H_{\tilde{n}\tilde{b}}^d - L_{\tilde{n}\tilde{b}}^d \right) \left(\frac{q^f}{m_0} \right) \right) \\
& \quad \left. + \left(g^{\nu\beta} H_{\beta\gamma}^d + \rho^2 \left(l_{bd} g^{\nu\beta} + l_{db} g^{\beta\nu} \right) L_{\beta\gamma}^d \right) \right. \\
& \quad \left. \times \left(\bar{\nabla}_\nu u^{\tilde{a}} + \frac{1}{r^2} l_{fd} g^{\tilde{a}\tilde{b}} \left(2 \nabla_\nu^{\text{gauge}} \Phi_b^d - L_{\nu\tilde{b}}^d \right) \left(\frac{q^f}{m_0} \right) \right) \right] \right\} \left(u^\gamma \left(\frac{q^b}{2\rho^2 m_0} \right) - v^b u^\gamma \right) \\
& - \mu \left[\tilde{g}^{(\tilde{n}\tilde{b})} \rho_{,\tilde{b}} k_{bc} \left(\widehat{\nabla}_{\tilde{n}} u^{\tilde{a}} + \frac{1}{2} l_{df} g^{\tilde{a}\tilde{c}} \left(2 H_{\tilde{n}\tilde{c}}^d - L_{\tilde{n}\tilde{c}}^d \right) \left(\frac{q^f}{m_0} \right) \right) \right.
\end{aligned}$$

(15.32_{cont.})

$$\begin{aligned}
& + \tilde{g}^{(\nu\beta)} \rho_{,\beta} k_{bc} \left(\bar{\nabla}_\nu u^{\tilde{a}} + \frac{1}{2} l_{df} g^{\tilde{a}\tilde{c}} \left(2 \overset{\text{gauge}}{\nabla}_\nu \Phi_{\tilde{c}}^d - L_{\nu\tilde{c}}^d \right) \left(\frac{q^f}{m_0} \right) \right) \left(\frac{q^c v^b}{m_0} \right) \\
& + \frac{2}{r^2} \left(\rho^2 l_{dn} g^{\tilde{d}\tilde{a}} \left(L_{\tilde{d}\tilde{b}}^d u^{\tilde{b}} + L_{\tilde{d}\beta}^d u^\beta \right) + \frac{1}{2\rho^2} \tilde{g}^{(\tilde{a}\tilde{b})} \rho_{,\tilde{b}} l_{bn} \left(\frac{q^b}{m_0} \right) \right) \\
& \times \left((H_{\mu\nu}^n - L_{\mu\nu}^n) v^\mu u^\nu + (H_{\tilde{m}\tilde{n}}^n - L_{\tilde{m}\tilde{n}}^n) v^{\tilde{m}} u^{\tilde{n}} \right. \\
& \left. + \left(\overset{\text{gauge}}{\nabla}_\mu \Phi_{\tilde{n}}^n - L_{\mu\tilde{n}}^n \right) v^\mu u^{\tilde{n}} - \left(\overset{\text{gauge}}{\nabla}_\nu \Phi_{\tilde{m}}^n - L_{\nu\tilde{m}}^n \right) v^{\tilde{m}} u^\nu \right) \\
& - \frac{1}{r^2} \left(g_{\zeta\mu} \tilde{g}^{(\tilde{a}\tilde{c})} \rho_{,\tilde{c}} \rho l_{bn} g^{\delta\zeta} \left(L_{\delta\beta}^b u^\beta + L_{\delta\tilde{b}}^b u^{\tilde{b}} \right) + \frac{1}{2\rho^2} \tilde{g}^{(\tilde{a}\tilde{b})} \rho_{,\tilde{b}} l_{bm} \left(\frac{q^b}{m_0} \right) \right) \\
& \times \left(v^\mu \left(\frac{q^m}{2\rho^2 m_0} \right) - v^m u^\mu \right) \\
& - \frac{1}{r^2} \left(g_{\tilde{z}\tilde{m}} \tilde{g}^{(\tilde{a}\tilde{c})} \rho_{,\tilde{c}} \rho l_{bn} g^{\tilde{d}\tilde{z}} \left(L_{\tilde{d}\beta}^d u^\beta + L_{\tilde{d}\tilde{b}}^d u^{\tilde{b}} \right) + \frac{1}{2\rho^2} \tilde{g}^{(\tilde{a}\tilde{b})} \rho_{,\tilde{b}} l_{bm} \left(\frac{q^b}{m_0} \right) \right) \\
& \times \left(v^{\tilde{m}} \left(\frac{q^m}{2\rho^2 m_0} \right) - v^m u^{\tilde{m}} \right) \\
& - \frac{1}{2r^2 \rho^2} \tilde{Q}^n_{mb} \left(\rho^2 l_{dn} g^{\tilde{d}\tilde{a}} \left(L_{\tilde{d}\beta}^d u^\beta + L_{\tilde{d}\tilde{b}}^d u^{\tilde{b}} \right) - \frac{1}{2\rho} \tilde{g}^{(\tilde{a}\tilde{b})} \rho_{,\tilde{b}} l_{pn} \left(\frac{q^p}{m_0} \right) \right) \\
& \times v^m \left(\frac{q^b}{m_0} \right) \\
& = 0
\end{aligned} \tag{15.32cont.}$$

where $\widehat{R}^{\tilde{a}}_{\tilde{b}\tilde{m}\tilde{n}}$ means a curvature tensor of a connection $\widehat{\omega}^{\tilde{a}}_{\tilde{b}}$ on M and $\widehat{Q}^{\tilde{n}}_{\tilde{m}\tilde{b}}$ its tensor of torsion. $\tilde{g}^{(\tilde{a}\tilde{b})}$ and $\tilde{g}^{(\alpha\beta)}$ are defined similarly to $\tilde{\gamma}^{(AB)}$. We have

$$\zeta^A = (\zeta^\alpha, \zeta^{\tilde{a}}) \tag{15.33}$$

$$v^A = (v^\alpha, v^{\tilde{a}}) \tag{15.34}$$

$$u^A = (u^\alpha, u^{\tilde{a}}). \tag{15.35}$$

Let us pass to the Eq. (15.24) and develop it according to our scheme. One gets:

$$\begin{aligned}
& \frac{dv^a}{d\tau} + \tilde{R}^a_{bcd} \left(\frac{q^b}{2\rho^2 m_0} \right) \zeta^c \left(\frac{q^d}{2\rho^2 m_0} \right) - \tilde{Q}^c_{mb} \tilde{\nabla}_c \left(\frac{q^a}{2\rho^2 m_0} \right) v^m \left(\frac{q^b}{2\rho^2 m_0} \right) \\
& + \left(2\rho^2 l_{bd} g^{\gamma\delta} \left(2H_{[\nu|\delta]}^d - L_{[\nu|\delta]}^d \right) L_{|\gamma|\mu]}^a + \frac{1}{\rho} \delta^a_b g_{\delta\alpha} \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} \overline{Q}^\alpha_{\mu\nu} \right.
\end{aligned} \tag{15.36}$$

$$\begin{aligned}
& - 2\delta^a{}_b \bar{\nabla}_{[\nu} \left(\frac{1}{\rho} g_{|\delta|\mu|} \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} \right) \\
& + \left(\frac{1}{r^2} g^{\tilde{c}\tilde{d}} \left(2 \overset{\text{gauge}}{\nabla}_{[\nu} \Phi_{|\tilde{d}|}^d - L_{[\nu|\tilde{d}|}^d \right) L_{|\tilde{d}|\mu]}^a \right) \left(\frac{q^b \zeta^\mu u^\nu}{2\rho^2 m_0} \right) \\
& + \left(2\rho^2 l_{bd} g^{\gamma\delta} \left(- \overset{\text{gauge}}{\nabla}_\delta \Phi_{\tilde{n}}^d + \frac{1}{2} L_{\delta\tilde{n}}^d \right) L_{\gamma\mu}^a - \delta^a{}_b \widehat{\nabla}_{\tilde{n}} \left(\frac{1}{\rho} g_{\delta\mu} \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} \right) \right. \\
& \quad \left. + \frac{1}{2r^2} \left(g^{\tilde{c}\tilde{d}} (2H_{\tilde{n}\tilde{d}}^d - L_{\tilde{n}\tilde{d}}^d) L_{\tilde{d}\mu}^a \right) \right) \left(\frac{q^b \zeta^\mu u^{\tilde{n}}}{2\rho^2 m_0} \right) \\
& + 2\rho^2 l_{bd} \left(g^{\gamma\delta} \left(- \overset{\text{gauge}}{\nabla}_\delta \Phi_{\tilde{n}}^d + \frac{1}{2} L_{\delta\tilde{n}}^d \right) L_{\gamma\tilde{m}}^a \right. \\
& \quad \left. + \frac{1}{\rho} \delta^a{}_b g_{\tilde{d}\tilde{a}} \tilde{g}^{(\tilde{d}\tilde{c})} \rho_{,\tilde{c}} \widehat{Q}^{\tilde{a}}{}_{\tilde{m}\tilde{n}} - 2\delta^a{}_b \widehat{\nabla}_{[\tilde{n}} \left(\frac{1}{\rho} g_{|\tilde{d}|\tilde{m}|} \tilde{g}^{(\tilde{d}\tilde{c})} \rho_{,\tilde{c}} \right) \right. \\
& \quad \left. + \frac{1}{r^2} g^{\tilde{c}\tilde{d}} \left(2H_{[\tilde{n}|\tilde{d}]}^d - L_{[\tilde{n}|\tilde{d}]}^d \right) L_{|\tilde{c}|\tilde{m}}^a \right) \left(\frac{q^b \zeta^{\tilde{m}} u^{\tilde{n}}}{2\rho^2 m_0} \right) \\
& + 2\rho^2 l_{bd} \left(\frac{1}{2} g^{\gamma\delta} (2H_{\nu\delta}^d - L_{\nu\delta}^d) L_{\gamma\tilde{m}}^a - \delta^a{}_b \bar{\nabla}_\nu \left(\frac{1}{\rho} g_{\tilde{d}\tilde{m}} \tilde{g}^{(\tilde{d}\tilde{c})} \rho_{,\tilde{c}} \right) \right. \\
& \quad \left. + \frac{1}{2r^2} g^{\tilde{c}\tilde{d}} \left(2 \overset{\text{gauge}}{\nabla}_\nu \Phi_{\tilde{d}}^d - L_{\nu\tilde{d}}^d \right) L_{\gamma\tilde{m}}^a \right) \left(\frac{q^b \zeta^{\tilde{m}} u^\nu}{2\rho^2 m_0} \right) \\
& + \left(\rho \tilde{g}^{(\gamma\beta)} \rho_{,\beta} l_{bc} L_{\gamma\mu}^a + \frac{1}{r^2} \rho \tilde{g}^{(\tilde{c}\tilde{b})} \rho_{,\tilde{b}} l_{bc} L_{b\mu}^a + \rho l_{bd} \tilde{g}^{(\beta\nu)} \rho_{,\nu} (2H_{\mu\beta}^d - L_{\mu\beta}^d) \delta^a{}_c \right. \\
& \quad \left. + \frac{1}{r^2} \rho l_{bd} \tilde{g}^{(\tilde{b}\tilde{n})} \rho_{,\tilde{n}} \left(2 \overset{\text{gauge}}{\nabla}_\mu \Phi_{\tilde{b}}^d - L_{\mu\tilde{b}}^d \right) \delta^a{}_c \right) \left(\frac{q^b}{2\rho^2 m_0} \right) \left(\zeta^c u^\mu - \zeta^\mu \left(\frac{q^c}{2\rho^2 m_0} \right) \right) \\
& + \left(\rho \tilde{g}^{(\gamma\beta)} \rho_{,\beta} l_{bc} L_{\gamma\tilde{m}}^a + \frac{1}{r^2} \rho \tilde{g}^{(\tilde{c}\tilde{b})} \rho_{,\tilde{b}} l_{bc} L_{b\tilde{m}}^a - \rho l_{bd} \tilde{g}^{(\beta\nu)} \rho_{,\nu} \left(2 \overset{\text{gauge}}{\nabla}_\beta \Phi_{\tilde{m}}^d - L_{\beta\tilde{m}}^d \right) \delta^a{}_c \right. \\
& \quad \left. + \frac{1}{r^2} \rho l_{bd} \tilde{g}^{(\tilde{b}\tilde{n})} \rho_{,\tilde{n}} (2H_{\tilde{m}\tilde{b}}^d - L_{\tilde{m}\tilde{b}}^d) \delta^a{}_c \right) \left(\frac{q^b}{2\rho^2 m_0} \right) \left(\zeta^c u^{\tilde{m}} - \zeta^{\tilde{m}} \left(\frac{q^c}{2\rho^2 m_0} \right) \right) \\
& + \left(\tilde{\nabla}_b L_{\beta\gamma}^a - \bar{\nabla}_\gamma \left(\frac{1}{\rho} g_{\beta\delta} \tilde{g}^{(\delta\nu)} \rho_{,\nu} \right) \delta^a{}_b + \rho^2 l_{db} g^{\delta\mu} L_{\mu\gamma}^a L_{\delta\beta}^d \right. \\
& \quad \left. - \frac{1}{\rho^2} g_{\delta\gamma} \tilde{g}^{(\delta\beta)} \rho_{,\beta} g_{\beta\nu} \tilde{g}^{(\nu\mu)} \rho_{,\mu} \delta^a{}_b \right) u^\beta \left(\zeta^b u^\mu - \zeta^\mu \left(\frac{q^b}{2\rho^2 m_0} \right) \right) \\
& + \left(\tilde{\nabla}_b L_{\beta\tilde{c}}^a - \widehat{\nabla}_{\tilde{c}} \left(\frac{1}{\rho} g_{\beta\delta} \tilde{g}^{(\delta\nu)} \rho_{,\nu} \right) \delta^a{}_b + \frac{1}{r^2} l_{db} g^{\tilde{d}\tilde{m}} L_{\tilde{m}\tilde{c}}^a L_{\tilde{d}\beta}^d \right.
\end{aligned}$$

(15.36_{cont.})

$$\begin{aligned}
& + \rho^2 l_{db} g^{\delta\mu} L_{\mu\tilde{c}}^a L_{\delta\beta}^d - \frac{1}{\rho^2} g_{\tilde{d}\tilde{c}} \tilde{g}^{(\tilde{d}\tilde{b})} \rho_{,\tilde{b}} g_{\beta\nu} \tilde{g}^{(\nu\mu)} \rho_{,\mu} \delta^a_b \Big) u^\beta \left(\zeta^b u^{\tilde{c}} - \zeta^{\tilde{c}} \left(\frac{q^b}{2\rho^2 m_0} \right) \right) \\
& + \left(\tilde{\nabla}_b L_{\tilde{b}\tilde{c}}^a - \widehat{\nabla}_{\tilde{c}} \left(\frac{1}{\rho} g_{\tilde{b}\tilde{d}} \tilde{g}^{(\tilde{d}\tilde{n})} \rho_{,\tilde{n}} \right) \delta^a_b + \rho^2 l_{db} g^{\delta\mu} L_{\mu\tilde{c}}^a L_{\delta\tilde{b}}^d \right. \\
& \quad \left. + \rho^2 l_{db} g^{\tilde{d}\tilde{m}} L_{\tilde{m}\tilde{c}}^a L_{\tilde{d}\tilde{b}}^d - \frac{1}{\rho^2} g_{\tilde{d}\tilde{c}} \tilde{g}^{(\tilde{d}\tilde{e})} \rho_{,\tilde{e}} g_{\tilde{b}\tilde{n}} \tilde{g}^{(\tilde{n}\tilde{m})} \rho_{,\tilde{m}} \delta^a_b \right) u^{\tilde{b}} \left(\zeta^b u^{\tilde{c}} - \zeta^{\tilde{c}} \left(\frac{q^b}{2\rho^2 m_0} \right) \right) \\
& + \left(\tilde{\nabla}_b L_{\tilde{b}\gamma}^a - \overline{\nabla}_\gamma \left(\frac{1}{\rho} g_{\tilde{b}\tilde{d}} \tilde{g}^{(\tilde{d}\tilde{n})} \rho_{,\tilde{n}} \right) \delta^a_b + \rho^2 l_{db} g^{\delta\mu} L_{\mu\gamma}^a L_{\delta\tilde{b}}^d \right. \\
& \quad \left. + \rho^2 l_{db} g^{\tilde{d}\tilde{m}} L_{\tilde{m}\gamma}^a L_{\tilde{d}\tilde{b}}^d - \frac{1}{\rho^2} g_{\delta\gamma} \tilde{g}^{(\delta\varepsilon)} \rho_{,\varepsilon} g_{\tilde{b}\tilde{n}} \tilde{g}^{(\tilde{n}\tilde{m})} \rho_{,\tilde{m}} \delta^a_b \right) u^{\tilde{b}} \left(\zeta^b u^\gamma - \zeta^\gamma \left(\frac{q^b}{2\rho^2 m_0} \right) \right) \\
& + \left(-\frac{1}{\rho} g_{\beta\delta} \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} C^a_{bc} \right. \\
& \quad \left. + 2\rho \left(g_{\gamma\nu} \tilde{g}^{(\nu\mu)} \rho_{,\mu} g^{\delta\gamma} L_{\delta\beta}^d + \frac{1}{r^2} g_{\tilde{c}\tilde{n}} \tilde{g}^{(\tilde{n}\tilde{m})} \rho_{,\tilde{m}} g^{\tilde{d}\tilde{c}} L_{\tilde{d}\beta}^d \right) l_{d[b}\delta^a_{c]} \right) \left(\frac{\zeta^b q^c}{2\rho^2 m_0} \right) u^\beta \\
& + \left(-\frac{1}{\rho} g_{\tilde{b}\tilde{d}} \tilde{g}^{(\tilde{d}\tilde{e})} \rho_{,\tilde{e}} C^a_{bc} \right. \\
& \quad \left. + 2\rho \left(g_{\gamma\nu} \tilde{g}^{(\nu\mu)} \rho_{,\mu} g^{\delta\gamma} L_{\delta\tilde{b}}^d + \frac{1}{r^2} g_{\tilde{c}\tilde{n}} \tilde{g}^{(\tilde{n}\tilde{m})} \rho_{,\tilde{m}} g^{\tilde{d}\tilde{c}} L_{\tilde{d}\tilde{b}}^d \right) l_{d[b}\delta^a_{c]} \right) \left(\frac{\zeta^b q^c}{2\rho^2 m_0} \right) u^{\tilde{b}} \\
& - 2 \left(g_{\gamma\delta} \tilde{g}^{(\delta\mu)} \rho_{,\mu} \tilde{g}^{(\gamma\beta)} \rho_{,\beta} + \frac{1}{r^2} g_{\tilde{c}\tilde{d}} \tilde{g}^{(\tilde{d}\tilde{m})} \rho_{,\tilde{m}} \tilde{g}^{(\tilde{c}\tilde{b})} \rho_{,\tilde{b}} \right) l_{b[c}\delta^a_{d]} \left(\frac{q^b \zeta^d q^c}{4\rho^4 m_0^2} \right) \\
& + \left(2\overline{\nabla}_{[\mu} L_{|\beta|\nu]}^a + L_{\beta\gamma}^a \overline{Q}^\gamma_{\mu\nu} + \frac{2}{\rho} g_{\beta\delta} \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} H_{\mu\nu}^a + \frac{2}{\rho} g_{\delta[\mu} L_{|\beta|\nu]}^a \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} \right) u^\beta \zeta^\mu u^\nu \\
& + \left(2\widehat{\nabla}_{[\tilde{m}} L_{|\tilde{b}|\tilde{n}]}^a + L_{\tilde{b}\tilde{c}}^a \widehat{Q}^{\tilde{c}}_{\tilde{m}\tilde{n}} + \frac{2}{\rho} g_{\tilde{b}\tilde{d}} \tilde{g}^{(\tilde{d}\tilde{e})} \rho_{,\tilde{e}} H_{\tilde{m}\tilde{n}}^a + \frac{2}{\rho} g_{\tilde{d}[\tilde{m}} L_{|\tilde{b}|\tilde{n}]}^a \tilde{g}^{(\tilde{d}\tilde{e})} \rho_{,\tilde{e}} \right) u^{\tilde{b}} \zeta^{\tilde{m}} u^{\tilde{n}} \\
& + \left(2\overline{\nabla}_{[\mu} L_{|\tilde{b}|\nu]}^a + L_{\tilde{b}\gamma}^a \overline{Q}^\gamma_{\mu\nu} + \frac{2}{\rho} g_{\tilde{b}\tilde{d}} \tilde{g}^{(\tilde{d}\tilde{e})} \rho_{,\tilde{e}} H_{\mu\nu}^a + \frac{2}{\rho} g_{\delta[\mu} L_{|\tilde{b}|\nu]}^a \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} \right) u^{\tilde{b}} \zeta^\mu u^\nu \\
& + \left(\widehat{\nabla}_{\tilde{m}} L_{\beta\nu}^a - \frac{2}{\rho} g_{\beta\delta} \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} \overset{\text{gauge}}{\nabla}_\nu \Phi_{\tilde{m}}^a + \frac{1}{\rho} g_{\tilde{d}\tilde{m}} L_{\beta\nu}^a \tilde{g}^{(\tilde{d}\tilde{e})} \rho_{,\tilde{e}} \right) u^\beta \zeta^{\tilde{m}} u^\nu \\
& + \left(\overline{\nabla}_\nu L_{\beta\tilde{n}}^a + \frac{2}{\rho} g_{\beta\delta} \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} \overset{\text{gauge}}{\nabla}_\mu \Phi_{\tilde{n}}^a + \frac{1}{\rho r^2} g_{\delta\mu} L_{\beta\tilde{n}}^a \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} \right) u^\beta \zeta^\mu u^{\tilde{n}} \\
& + \left(2\widehat{\nabla}_{[\tilde{m}} L_{|\beta|\tilde{n}]}^a + L_{\beta\tilde{c}}^a \widehat{Q}^{\tilde{c}}_{\tilde{m}\tilde{n}} + \frac{2}{\rho} g_{\beta\delta} \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} H_{\tilde{m}\tilde{n}}^a + \frac{2}{\rho} g_{\tilde{d}[\tilde{m}} L_{|\beta|\tilde{n}]}^a \tilde{g}^{(\tilde{d}\tilde{e})} \rho_{,\tilde{e}} \right) u^\beta \zeta^{\tilde{m}} u^{\tilde{n}} \\
& + \left(\overline{\nabla}_\mu L_{\tilde{b}\tilde{n}}^a + \frac{2}{\rho} g_{\tilde{b}\tilde{d}} \tilde{g}^{(\tilde{d}\tilde{e})} \rho_{,\tilde{e}} \overset{\text{gauge}}{\nabla}_\mu \Phi_{\tilde{n}}^a + \frac{1}{\rho} g_{\delta\mu} L_{\tilde{b}\tilde{n}}^a \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} \right) u^{\tilde{b}} \zeta^\mu u^{\tilde{n}}
\end{aligned}$$

(15.36cont.)

$$\begin{aligned}
& + \left(\widehat{\nabla}_{\tilde{m}} L_{\tilde{b}\nu}^a - \frac{2}{\rho} g_{\tilde{b}\tilde{d}} \tilde{g}^{(\tilde{d}\tilde{c})} \rho_{,\tilde{c}} \overset{\text{gauge}}{\nabla}_{\nu} \Phi_{\tilde{m}}^a + \frac{1}{\rho} g_{\tilde{d}\tilde{m}} L_{\tilde{b}\nu}^a \tilde{g}^{(\delta\tilde{c})} \rho_{,\tilde{c}} \right) u^{\tilde{b}} \zeta^{\tilde{m}} u^{\nu} \\
& - \frac{1}{2\rho} \tilde{Q}^a{}_{mb} v^m \left(\frac{q^b}{m_0} \right) \left(g_{\beta\delta} \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} u^{\beta} + g_{\tilde{b}\tilde{d}} \tilde{g}^{(\tilde{d}\tilde{c})} \rho_{,\tilde{c}} u^{\tilde{b}} \right) \\
& + \widehat{\tilde{Q}}^{\tilde{n}}{}_{\tilde{m}\tilde{z}} u^{\tilde{z}} v^{\tilde{m}} \left(\frac{\rho_{,\tilde{n}}}{\rho^3} \left(\frac{q^a}{m_0} \right) - \left(L_{\beta\tilde{n}}^a u^{\beta} + L_{\tilde{b}\tilde{n}}^a u^{\tilde{b}} \right) - \frac{1}{2\rho^3} g_{\tilde{d}\tilde{n}} \tilde{g}^{(\tilde{d}\tilde{c})} \rho_{,\tilde{c}} \left(\frac{q^a}{m_0} \right) \right) \\
& + \overline{Q}^{\nu}{}_{\mu\zeta} u^{\zeta} v^{\mu} \left(\frac{\rho_{,\nu}}{\rho^3} \left(\frac{q^a}{m_0} \right) - \left(L_{\beta\nu}^a u^{\beta} + L_{\tilde{b}\nu}^a u^{\tilde{b}} \right) - \frac{1}{2\rho^3} g_{\delta\nu} \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} \left(\frac{q^a}{m_0} \right) \right) \\
& - \frac{1}{\rho} g_{\zeta\mu} \tilde{g}^{(\zeta\omega)} \rho_{,\omega} \left(\tilde{\nabla}_b \left(\frac{q^a}{2\rho^2 m_0} \right) \right. \\
& \quad \left. + \frac{1}{\rho} \left(g_{\beta\delta} \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} u^{\beta} + g_{\tilde{b}\tilde{d}} \tilde{g}^{(\tilde{d}\tilde{c})} \rho_{,\tilde{c}} u^{\tilde{b}} \right) \delta^a{}_b \right) \left(v^{\mu} \left(\frac{q^b}{2\rho^2 m_0} \right) - v^b u^{\mu} \right) \\
& - \frac{1}{\rho} g_{\tilde{z}\tilde{m}} \tilde{g}^{(\tilde{z}\tilde{p})} \rho_{,\tilde{p}} \left(\tilde{\nabla}_b \left(\frac{q^a}{2\rho^2 m_0} \right) \right. \\
& \quad \left. + \frac{1}{\rho} \left(g_{\tilde{b}\tilde{d}} \tilde{g}^{(\tilde{d}\tilde{c})} \rho_{,\tilde{c}} u^{\tilde{b}} + g_{\beta\delta} \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} u^{\beta} \right) \delta^a{}_b \right) \left(v^{\tilde{m}} \left(\frac{q^b}{2\rho^2 m_0} \right) - v^b u^{\tilde{m}} \right) \\
& - 2 \left(H_{\mu\nu}^n - L_{\mu\nu}^n \right) \left(\tilde{\nabla}_n \left(\frac{q^a}{2\rho^2 m_0} \right) + \frac{1}{\rho} \left(g_{\beta\delta} \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} u^{\beta} + g_{\tilde{b}\tilde{d}} \tilde{g}^{(\tilde{d}\tilde{c})} \rho_{,\tilde{c}} u^{\tilde{b}} \right) \delta^a{}_n \right) v^{\mu} u^{\nu} \\
& - 2 \left(H_{\tilde{m}\tilde{n}}^n - L_{\tilde{m}\tilde{n}}^n \right) \left(\tilde{\nabla}_n \left(\frac{q^a}{2\rho^2 m_0} \right) + \frac{1}{\rho} \left(g_{\beta\delta} \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} u^{\beta} + g_{\tilde{b}\tilde{d}} \tilde{g}^{(\tilde{d}\tilde{c})} \rho_{,\tilde{c}} u^{\tilde{b}} \right) \delta^a{}_n \right) v^{\tilde{m}} u^{\tilde{n}} \\
& - 2 \left(\overset{\text{gauge}}{\nabla}_{\mu} \Phi_{\tilde{n}}^n - L_{\mu\tilde{n}}^n \right) \left(\tilde{\nabla}_n \left(\frac{q^a}{2\rho^2 m_0} \right) + \frac{1}{\rho} \left(g_{\beta\delta} \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} u^{\beta} + g_{\tilde{b}\tilde{d}} \tilde{g}^{(\tilde{d}\tilde{c})} \rho_{,\tilde{c}} u^{\tilde{b}} \right) \right) v^{\mu} u^{\tilde{n}} \\
& + 2 \left(\overset{\text{gauge}}{\nabla}_{\nu} \Phi_{\tilde{m}}^n - L_{\nu\tilde{m}}^n \right) \left(\tilde{\nabla}_n \left(\frac{q^a}{2\rho^2 m_0} \right) + \frac{1}{\rho} \left(g_{\beta\delta} \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} u^{\beta} + g_{\tilde{b}\tilde{d}} \tilde{g}^{(\tilde{d}\tilde{c})} \rho_{,\tilde{c}} u^{\tilde{b}} \right) \right) u^{\nu} v^{\tilde{m}} \\
& - \left\{ \left[2\mu\rho \tilde{g}^{(\nu\mu)} \rho_{,\mu} k_{bc} \left(-\frac{1}{\rho^2} \left(\frac{q^a}{m_0} \right) \rho_{,\nu} + (L_{\gamma\nu}^a u^{\gamma} + L_{\tilde{c}\nu}^a u^{\tilde{c}}) \right. \right. \right. \\
& \quad \left. \left. + \frac{1}{2\rho^3} g_{\delta\nu} \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} \left(\frac{q^a}{m_0} \right) \right) \right] \\
& \quad + \left[2\mu\rho \tilde{g}^{(\tilde{n}\tilde{m})} \rho_{,\tilde{m}} k_{bc} \left(-\frac{1}{\rho^2} \left(\frac{q^a}{m_0} \right) \rho_{,\tilde{n}} + (L_{\gamma\tilde{n}}^a u^{\gamma} + L_{\tilde{c}\tilde{n}}^a u^{\tilde{c}}) \right. \right. \\
& \quad \left. \left. + \frac{1}{2\rho^3} g_{\tilde{d}\tilde{n}} \tilde{g}^{(\tilde{d}\tilde{c})} \rho_{,\tilde{c}} \left(\frac{q^a}{m_0} \right) \right) \right] \right\} \left(\frac{v^b q^c}{2\rho^2 m_0} \right)
\end{aligned}$$

(15.36_{cont.})

$$\begin{aligned}
& - \left(2\rho^3 l_{bd} g^{\nu\beta} H_{\gamma\beta}^d + \rho^2 (l_{bd} g^{\nu\beta} + l_{db} g^{\beta\nu}) L_{\beta\gamma}^d \right) \\
& \quad \times \left(-\frac{1}{\rho^3} \left(\frac{q^a}{m_0} \right) \rho_{,\nu} + L_{\delta\nu}^a u^\delta + L_{\tilde{d}\nu}^a u^{\tilde{d}} + \frac{1}{2\rho^3} g_{\delta\nu} \tilde{g}^{(\delta\varepsilon)} \rho_{,\varepsilon} \left(\frac{q^a}{m_0} \right) \right) \\
& \quad \times \left(v^\gamma \left(\frac{q^b}{2\rho^2 m_0} \right) - v^b u^\gamma \right) \\
& - \left(2\rho^3 l_{bd} g^{\tilde{n}\tilde{b}} \tilde{\nabla}_\gamma^{\text{gauge}} \Phi_{\tilde{b}}^d + \rho^2 (l_{bd} g^{\tilde{n}\tilde{b}} + l_{db} g^{\tilde{b}\tilde{n}}) L_{\tilde{b}\gamma}^d \right) \\
& \quad \times \left(-\frac{1}{\rho^3} \left(\frac{q^a}{m_0} \right) \rho_{,\tilde{n}} + L_{\alpha\tilde{n}}^a u^\alpha + L_{\tilde{c}\tilde{n}}^a u^{\tilde{c}} + \frac{1}{2\rho^3} g_{\tilde{d}\tilde{n}} \tilde{g}^{(\tilde{d}\tilde{c})} \rho_{,\tilde{c}} \left(\frac{q^a}{m_0} \right) \right) \\
& \quad \times \left(v^\gamma \left(\frac{q^b}{m_0} \right) - v^b u^\gamma \right) \\
& - \left(2\rho^3 l_{bd} g^{\tilde{n}\tilde{b}} H_{\tilde{c}\tilde{b}}^d + \rho^2 (l_{bd} g^{\tilde{n}\tilde{b}} + l_{db} g^{\tilde{b}\tilde{n}}) L_{\tilde{b}\tilde{c}}^d \right) \\
& \quad \times \left(-\frac{1}{\rho^3} \left(\frac{q^a}{m_0} \right) \rho_{,\tilde{n}} + L_{\tilde{c}\tilde{n}}^a u^{\tilde{c}} + L_{\alpha\tilde{n}}^a u^\alpha + \frac{1}{2\rho^3} g_{\tilde{d}\tilde{n}} \tilde{g}^{(\tilde{d}\tilde{e})} \rho_{,\tilde{e}} \left(\frac{q^a}{m_0} \right) \right) \\
& \quad \times \left(v^{\tilde{c}} \left(\frac{q^b}{m_0} \right) - v^b u^{\tilde{c}} \right) \tag{15.36cont.} \\
& - \left(-2\rho^3 l_{bd} g^{\nu\beta} \tilde{\nabla}_\beta^{\text{gauge}} \Phi_{\tilde{c}}^d + \rho^2 (l_{bd} g^{\nu\beta} + l_{db} g^{\beta\nu}) L_{\beta\tilde{c}}^d \right) \\
& \quad \times \left(-\frac{1}{\rho^3} \left(\frac{q^a}{m_0} \right) \rho_{,\nu} + L_{\tilde{e}\nu}^a u^{\tilde{e}} + L_{\alpha\nu}^a u^\alpha + \frac{1}{2\rho^3} g_{\delta\nu} \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} \left(\frac{q^a}{m_0} \right) \right) \\
& \quad \times \left(v^{\tilde{c}} \left(\frac{q^b}{m_0} \right) - v^b u^{\tilde{c}} \right) \\
& + \frac{1}{2\rho^3} \left(\frac{q^a}{m_0} \right) \left(g_{\beta\delta} \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} v^\beta + g_{\tilde{b}\tilde{d}} \tilde{g}^{(\tilde{d}\tilde{c})} \rho_{,\tilde{c}} v^{\tilde{b}} \right) \\
& + L_{\beta\nu}^a v^\beta u^\nu + L_{\tilde{b}\tilde{n}}^a v^{\tilde{b}} u^{\tilde{n}} + L_{\tilde{b}\nu}^a v^{\tilde{b}} u^\nu + L_{\beta\tilde{n}}^a v^\beta u^{\tilde{n}} \\
& + \frac{1}{\rho} g_{\delta\nu} \tilde{g}^{(\delta\gamma)} \rho_{,\gamma} v^a u^\nu + \frac{1}{\rho} g_{\tilde{d}\tilde{n}} \tilde{g}^{(\tilde{d}\tilde{c})} \rho_{,\tilde{c}} v^a u^{\tilde{n}} \\
& = 0
\end{aligned}$$

where $\tilde{R}^a_{bcd}$ is a tensor of a curvature for a connection $\tilde{\omega}^a_b$ on a group H and $\tilde{Q}^a_{bc}$ is a tensor of a torsion for this connection. It is worth to notice that $Q^\gamma_{\tilde{m}\nu}$, $Q^\gamma_{\mu\tilde{n}}$ are equal to zero. Moreover for a convenience of a future consideration it is good to keep them there (i.e. in Eq. (15.32)).

Let us notice that u^α, u^β satisfy equations (15.2) and (15.3). In this way we get geodetic deviation equations in our theory. If we suppose that we are close to the

minimum of a Higgs' potential we should put in our equation $H_{\tilde{a}\tilde{b}}^n = 0$. What is the physical meaning of geodetic deviation equations? In order to find it let us consider a flow of geodesic $\Gamma(\sigma)$, $\sigma \in U \subset \mathbb{R}$ given by

$$x^{\tilde{A}} = x^{\tilde{A}}(\tau, \sigma), \quad (15.37)$$

$$\zeta^{\tilde{A}} = \left(\frac{\partial x^\alpha}{\partial \sigma}, \frac{\partial x^{\tilde{a}}}{\partial \sigma}, \frac{\partial x^a}{\partial \sigma} \right) \Big|_{\sigma=\sigma_0}, \quad (15.38)$$

$$\sigma_0 \in U.$$

σ is a parameter such that for every $\sigma_1 \neq \sigma_2$, $x^A(\tau, \sigma_1)$ and $x^A(\tau, \sigma_2)$ are different geodesics. One can say that we have a family of geodesic curves, $\Gamma(\sigma)$. The geodesic considered here is $\Gamma(\sigma_0)$, i.e. for $\sigma = \sigma_0$.

Thus

$$\frac{dx^a}{d\tau} = u^a = \frac{1}{2\rho^2} \left(\frac{q^a}{m_0} \right) \quad (15.39)$$

where $\rho = \rho(\tau, \sigma) = \rho(x(\tau, \sigma))$.

Thus one gets

$$x^a = \frac{1}{2} \left(\frac{q^a}{m_0} \right) (\sigma) \int_{\tau_0}^{\tau} \frac{d\tau}{\rho^2(\tau, \sigma)} + x_0^a(\sigma) \quad (15.40)$$

and

$$\zeta^a = \frac{\partial}{\partial \sigma} \left[\frac{q^a}{m_0} (\sigma) \int_{\tau_0}^{\tau} \frac{d\tau}{\rho^2(\tau, \sigma)} \right] \Big|_{\sigma=\sigma_0} + \frac{dx_0^a}{d\sigma} \quad (15.41)$$

$$v^a = \frac{d\zeta^a}{d\tau} = \frac{\partial}{\partial \tau} \left(\frac{q^a}{m_0} (\sigma) \frac{1}{\rho^2(\tau, \sigma)} \right) \Big|_{\sigma=\sigma_0}. \quad (15.42)$$

v^a is (of course) an Ad-type quantity. In this way we get

$$\frac{dv^a}{d\tau} = -\frac{\partial}{\partial \sigma} \left(\frac{q^a}{m_0} (\sigma) \frac{1}{\rho^3} \frac{\partial \rho}{\partial \tau} \right) \Big|_{\sigma=\sigma_0}. \quad (15.43)$$

In this way Eq. (15.22) together with Eq. (15.1) give us an interpretation of geodetic deviation equations in our theory. They are analogous to the deviation equations for charged particles moving in nonabelian Yang-Mills' field and Higgs' field and non-symmetric gravitational field as well.

Let us give some remarks on a physical interpretation of the vector $\zeta^{\tilde{A}} = (\zeta^\alpha, \zeta^{\tilde{a}}, \zeta^a)$. The vector $\zeta^{\tilde{A}}$ —“geodesic separation”—is a displacement (tangent vector) from point on a fiducial geodesic to a point on nearby geodesic characterized by the same value of an affine parameter τ . Thus $v^{\tilde{A}} = (v^\alpha, v^{\tilde{a}}, v^a)$ means a relative “velocity” and $u^{\tilde{B}} \nabla_{\tilde{B}} v^{\tilde{A}}$ —a relative “acceleration”—equals, according to Eq. (15.22) terms with a

curvature and torsion. Thus we get “tidal forces” in our theory $((n+n_1+4)$ -dimensional case), for charged (in a nonabelian gauge field sense and Higgs’ field) test particles. In this equation we get gravitational “tidal forces” from NGT, Yang-Mills’ “tidal forces”, Higgs’ field “tidal forces” and additional effects which can be treated as gravito-Yang-Mills-Higgs tidal forces. The scalar field ρ is also a source of additional “tidal forces”.

These new effects are “interference effects” between gravitational, nonabelian Yang-Mills’ interactions and Higgs’ field. We can project our equation on a space-time E (they are defined on a bundle manifold P), taking any local section e of the bundle P . In this way we get gauge dependent charges Q^a (Yang-Mills’ charged-colour charges) and gauge dependent $\bar{v}^a$. We can substitute $\bar{v}^a$ into Eq. (15.36). However, we should substitute in the place of $\frac{dv^a}{d\tau}$ the expression

$$\frac{d\bar{v}^a}{d\tau} - C^a_{bc} A^c_\mu u^\mu \bar{v}^b - C^a_{bc} \Phi^c_{\tilde{b}} \bar{v}^{\tilde{b}} \quad (15.44)$$

where

$$e^* \omega = A^a_\mu \bar{\theta}^\mu X_a + \Phi^a_{\tilde{b}} \tilde{\theta}^{\tilde{b}} X_a. \quad (15.45)$$

Finally, let us remark that Eq. (15.31) represents tidal gravito-Yang-Mills-Higgs forces and Eq. (15.36) is a relative change of $(\frac{q^a}{m_0})(\sigma)$ for different test particles via v^a (or $(\frac{Q^a}{m_0})(r)$ via $\bar{v}^a$).

Eq. (15.32) is a relative change of a Higgs charge $u^{\tilde{b}}$ for different test particles via $\bar{v}^{\tilde{b}}$. The interpretation of this equation is much more complex for the “Higgs charge” $u^{\tilde{b}}$ is not “conserved” during a motion of a test particle. The existence of a “constant Higgs charge” introduced here can help us in physical understanding of these interesting phenomena. Our gravito-Yang-Mills-Higgs-scalar tidal forces are a strict and natural generalization of gravito-electromagnetic-scalar tidal forces from the Nonsymmetric Kaluza–Klein (Jordan–Thiry) Theory (in an electromagnetic case) and gravito-Yang-Mills-scalar tidal forces from the Nonsymmetric Kaluza–Klein (Jordan–Thiry) Theory in a general nonabelian case (see Ref. [18]).

16. A Tower of Scalar Fields in the Nonsymmetric-Nonabelian Jordan–Thiry Theory. The Warp Factor

In section 5 we mention a more general case of the scalar field $\rho = \rho(x, y)$, $x \in E$, $y \in M$. Moreover it is more convenient to consider the field Ψ in place of ρ .

$$\rho = e^{-\Psi}. \quad (16.1)$$

In this case one finds:

$$\Psi(x, y) = \sum_{k=0}^{\infty} \Psi_k(x) \chi_k(y). \quad (16.2)$$

Let us find a condition for the function $\rho(x, y)$ such that $\Psi(x, y)$ is an $L^2(M, dm, \sqrt{|\widetilde{g}|})$ -valued function of $x \in E$. From (16.1) one finds that $\ln(\rho)$ is an $L^2(M, dm, \sqrt{|\widetilde{g}|})$ -valued function i.e.:

$$\int_M (\ln(\rho))^2 \sqrt{|\widetilde{g}|} dm(y) < \infty. \quad (16.2a)$$

Thus the sufficient condition for ρ is that ρ and ρ^{-1} are bounded on $M - A$, where $m(A) = 0$. It means that $\widetilde{\rho}(x), \widetilde{\rho}^{-1}(x) \in L^\infty(M, dm, \sqrt{|\widetilde{g}|})$ and $\|\widetilde{\rho}(x)\|_\infty = \text{ess sup}_{y \in M} |\rho(x, y)| < \infty$. The convergence $\widetilde{\rho}_n(x, \cdot) \rightarrow \widetilde{\rho}(x, \cdot)$ with respect to the norm $\|\cdot\|_\infty$ means a uniform limit. We have the same for $\widetilde{\rho}^{-1}(x)$. We can also consider a decomposition of $\rho(x, y)$ into a series of generalized harmonics on M i.e.

$$\rho(x, y) \sim \rho_0(x) + \sum_{k=1}^{\infty} \rho_k(x) \chi_k(y) \quad (16.2b)$$

such that

$$\begin{aligned} \rho_k(x) &= \frac{1}{V_2} \int_M \sqrt{|\widetilde{g}|} \rho(x, y) \chi_k(y) dm(y), \\ \rho_0(x) &= \frac{1}{V_2} \int_M \sqrt{|\widetilde{g}|} \rho(x, y) dm(y). \end{aligned} \quad (16.2c)$$

If $\sum_{k=0}^{\infty} \rho_k^2(x) < \infty$, $x \in E$ the series on the right-hand side of (16.2b) converges to the function $\rho(x, y)$ in $L^2(M, dm, \sqrt{|\widetilde{g}|})$ sense. The limit can be understood in a uniform sense if $\rho(x, y)$ is continuous and $V(x) = \text{Var}(\widetilde{\rho}(x)) < \infty$ in M for every $x \in E$.

Moreover it is interesting to know properties of $\tilde{\Psi}(x)$ if $\tilde{\rho}(x)$ is an $L^2(M, dm, \sqrt{|\tilde{g}|})$ -valued function. One gets:

$$\|\tilde{\rho}(x)\|^2 \leq V_2 + \sum_{n=1}^{\infty} \frac{2^n}{n!} (\|\tilde{\Psi}(x)\|_n)^n < \infty, \quad (16.2d)$$

where

$$\|\Psi(x)\|_n = \left(\int_M \sqrt{|\tilde{g}|} |\Psi(x, y)|^n dm(y) \right)^{1/n}$$

is an $L^n(M, dm, \sqrt{|\tilde{g}|})$ norm. Let us suppose that for every $n \in \mathbb{N}_1^\infty$, $\tilde{\Psi}(x)$ is an $L^n(M, dm, \sqrt{|\tilde{g}|})$ -valued function. For

$$V_2 = \int_M \sqrt{|\tilde{g}|} dm(y) < \infty$$

one gets

$$\|\tilde{\Psi}(x)\|_n \leq \|\tilde{\Psi}(x)\|_{n'} V_2^{1/n - 1/n'},$$

if $1 \leq n \leq n' \leq \infty$ and

$$\begin{aligned} L^\infty(M, dm, \sqrt{|\tilde{g}|}) &\subset L^{n'}(M, dm, \sqrt{|\tilde{g}|}) \subset L^n(M, dm, \sqrt{|\tilde{g}|}) \subset \\ &\subset L^1(M, dm, \sqrt{|\tilde{g}|}) = L(M, dm, \sqrt{|\tilde{g}|}). \end{aligned}$$

Thus if $n' \rightarrow \infty$ one obtains

$$\|\tilde{\Psi}(x)\|_n \leq \|\tilde{\Psi}(x)\|_\infty V_2^{1/n} = (\text{ess sup}_{y \in M} |\Psi(x, y)|) V_2^{1/n}. \quad (16.2e)$$

From (16.2d) and (16.2e) we have

$$\|\tilde{\rho}(x)\|^2 \leq V_2 e^{2\|\Psi(\tilde{x})\|_\infty}. \quad (16.2f)$$

Thus if $\tilde{\Psi}(x) \in L^\infty(M, dm, \sqrt{|\tilde{g}|})$ $\tilde{\rho}(x)$ is an $L^2(M, dm, \sqrt{|\tilde{g}|})$ -valued function. This means that $\Psi(x, y)$ is bounded on $M - A$ where $m(A) = 0$. In this way we get an interesting duality. The sufficient condition for $\tilde{\Psi}(x)$ to be an $L^2(M, dm, \sqrt{|\tilde{g}|})$ -valued function is $\tilde{\rho}(x), \tilde{\rho}^{-1}(x) \in L^\infty(M, dm, \sqrt{|\tilde{g}|})$ and the sufficient condition for $\tilde{\rho}(x)$ to be an $L^2(M, dm, \sqrt{|\tilde{g}|})$ is $\tilde{\Psi}(x) \in L^\infty(M, dm, \sqrt{|\tilde{g}|})$. In the second case we can try an expansion of $\Psi(x, y)$ into a series of generalized harmonics similarly as for ρ (i.e. Eqs. (16.2b–c)).

The $(n_1 + 4)$ -dimensional lagrangian for the scalar field Ψ looks like:

$$\mathcal{L}_{\text{scal}}(\Psi) = (\bar{M} \tilde{\gamma}^{(CM)} \Psi_{,C} \Psi_{,M} + n^2 \gamma^{[MN]} \gamma_{DM} \tilde{\gamma}^{(DC)} \Psi_{,N} \Psi_{,C}) \quad (16.3)$$

or

$$\begin{aligned} \mathcal{L}_{\text{scal}}(\Psi) &= (\bar{M} \tilde{g}^{(\gamma\mu)} \Psi_{,\gamma} \Psi_{,\mu} + n^2 g^{[\mu\nu]} g_{\delta\mu} \tilde{g}^{(\delta\gamma)} \Psi_{,\nu} \Psi_{,\gamma}) + \\ &+ \frac{1}{r^2} (\bar{M} \tilde{g}^{(\tilde{c}\tilde{m})} \Psi_{,\tilde{c}} \Psi_{,\tilde{m}} + n^2 g^{[\tilde{m}\tilde{n}]} g_{\tilde{d}\tilde{m}} \tilde{g}^{(\tilde{d}\tilde{c})} \Psi_{,\tilde{n}} \Psi_{,\tilde{c}}) = \tilde{\mathcal{L}}_{\text{scal}}(\Psi) + Q(\Psi). \end{aligned} \quad (16.4)$$

Moreover we should average over M and H .

Thus one gets using (16.2), (16.4) and (5.50):

$$\begin{aligned} \frac{1}{V_1 V_2} \int_M (\mathcal{L}_{\text{scal}}(\Psi) + Q(\Psi)) \sqrt{|\tilde{g}|} \sqrt{|\ell|} dm(y) d\mu_H(h) = \\ = \sum_{k=0}^{\infty} \mathcal{L}_{\text{scal}}(\Psi_k) + \frac{1}{2} \sum_{k,l=1}^{\infty} \widehat{M}_{kl} \Psi_k \Psi_l, \end{aligned} \quad (16.5)$$

where $\mathcal{L}_{\text{scal}}(\Psi_k)$ is given by the formula (7.6) i.e. is a usual lagrangian for the scalar field Ψ in our theory and

$$\frac{1}{2} \widehat{M}_{kl} = \frac{(-1)}{V_2 r^2} \int_M (\overline{M} \tilde{g}^{(\tilde{c}\tilde{m})} + n^2 g^{[\tilde{m}\tilde{n}]} g_{\tilde{d}\tilde{m}} \tilde{g}^{(\tilde{d}\tilde{c})}) \chi_{k,\tilde{c}} \chi_{l,\tilde{n}} \sqrt{|\tilde{g}|} dm(y) < \infty \quad (16.6)$$

and $\widehat{M}_{kl} = \widehat{M}_{lk}$, $k, l = 1, 2, \dots$

It is easy to see that we get a tower of fields Ψ_k such that according to (16.6) the field Ψ_0 is massless and remaining fields are massive with a scale of a mass $\frac{1}{r^2}$.

The infinite dimensional quadratic form

$$\frac{1}{2} \sum_{k,l=1}^{\infty} \widehat{M}_{kl} \Psi_k(x) \Psi_l(x)$$

is convergent in ℓ^2 for every x . This is easily satisfied if the derivatives of χ_k belong to $L^2(M, dm, \sqrt{|\tilde{g}|})$ and $g_{\tilde{a}\tilde{b}}$ are continuous functions on M , which is always satisfied for our case (M is compact without boundaries and $g_{\tilde{a}\tilde{b}}$ defined in section 4). For $L^2(M, dm, \sqrt{|\tilde{g}|})$ is unitary equivalent to $L^2(M, dm)$ we can consider χ_k eigenfunctions of Beltrami–Laplace operator on M .

In some cases we can diagonalize the infinite dimensional matrix $\widehat{M}_{kl}$ getting some new fields

$$\Psi'_{k'}(x) = \sum_{k=1}^{\infty} A_{k'k} \Psi_k(x), \quad (16.7)$$

such that

$$b_k \delta_{kl} = \sum_{k',l'=1}^{\infty} A_{kk'} A_{ll'} \widehat{M}_{k'l'}, \quad (16.8)$$

where $A = (A_{k'k})$ is a unitary operator in ℓ^2 space i.e.

$$\|Aa\| = \|a\| \quad (16.9)$$

and $A(\ell^2) = \ell^2$.

The diagonalization procedure can be achieved if $\widehat{M}_{kl}$ is a symmetric unbounded (Hermitian) operator in ℓ^2 . This is equivalent to

$$\widehat{M}_{kl} = \widehat{M}_{lk} \quad (16.10)$$

and $M(\ell^2) = \ell^2$. Thus we get for the lagrangian:

$$\mathcal{L}_{\text{scal}}(\Psi) = \sum_{k=0}^{\infty} \mathcal{L}_{\text{scal}}(\Psi'_k) - \sum_{k=1}^{\infty} \frac{b_k}{2} (\Psi'_k)^2, \quad (16.11)$$

$\left(\frac{b_k}{M}\right)$ have an interpretation as m_k^2 for Ψ'_k . If $\widehat{M}_{kl}$ is a negatively defined operator in ℓ^2 we have for every k $b_k \geq 0$ if M_{kl} is invertible $b_k > 0$ for every k . The latest condition means that every Ψ'_k is massive with $\frac{1}{r^2}$ as a scale of the mass. Let us write down a self-interaction term for Ψ'_k coming from cosmological terms. One gets:

$$V(\{\Psi'_k\}) = \frac{1}{V_2} \int_M \sqrt{|\tilde{g}|} dm(y) \left(\frac{\alpha_S^2}{\ell_{pl}^2} \tilde{R}(\tilde{\Gamma}) e^{(n+2)\Psi(x,y)} + \frac{e^{n\Psi(x,y)}}{r^2} \widehat{R}(\widehat{\Gamma}) \right), \quad (16.12)$$

where

$$\Psi(x, y) = \Psi_0(x) + \sum_{k=1}^{\infty} \Psi_k(x) \chi_k(y) \quad (16.13)$$

and

$$\Psi'_k = \sum_{k'=1}^{\infty} A_{kk'} \Psi_{k'}. \quad (16.14)$$

The field $\Psi_0(x)$ can get a mass from a cosmological background. If we suppose $\Psi = \Psi(x)$ it means really that $\Psi(x) = \Psi_0(x)$ i.e. a truncation. In this more general case for the field ρ (or Ψ) one can easily get similar formulae for the remaining part of the lagrangian involving this field i.e. averaging over the manifold M .

All the factors in front of lagrangian terms are exponential functions of Ψ . The sufficient condition to make all the integration over M convergent is to suppose that $\tilde{\Psi}(x) \in L^2(M, dm, \sqrt{|\tilde{g}|}) \cap L^\infty(M, dm, \sqrt{|\tilde{g}|})$ i.e. $\tilde{\Psi}(x)$ is L^∞ -valued function.

In the case of a completely broken group G we should put G for M .

Let us give a more rigorous justification for an intuitive procedure concerning diagonalization of the infinite matrix $\widehat{M}$.

Let us consider a bilinear form in $L^2(M, dm, \sqrt{|\tilde{g}|}) = L^2 \simeq L^2(M, dm)$

$$\widehat{M}(f, g) = \frac{1}{V_2 r^2} \int_M (\overline{M} \tilde{g}^{(\tilde{c}\tilde{m})} + n^2 g^{[\tilde{m}(\tilde{n})]} g_{\tilde{d}\tilde{n}} \tilde{g}^{(|\tilde{d}|\tilde{c})}) f_{,\tilde{c}} g_{,\tilde{m}} \sqrt{|\tilde{g}|} dm(y), \quad (16.15)$$

for $f, g \in L^2(M, dm, \sqrt{|\tilde{g}|}) \cap \mathbb{C}^1(M)$.

This form is always defined for f and g , because $g_{\tilde{a}\tilde{b}}$ are smooth functions on a compact manifold and $f_{,\tilde{c}}, g_{,\tilde{m}} \in L^2(M, dm, \sqrt{|\tilde{g}|})$ (if they exist). $\mathbb{C}^1(M)$ is linearly dense in $L^2(M, dm, \sqrt{|\tilde{g}|})$. Let us suppose that the tensor $P^{(\tilde{c}\tilde{m})}$, where

$$P^{\tilde{c}\tilde{m}} = (\overline{M}\tilde{g}^{\tilde{c}\tilde{m}} + n^2 g^{[\tilde{m}\tilde{n}]} g_{\tilde{d}\tilde{n}} \tilde{g}^{(\tilde{d}\tilde{c})})$$

is invertible and negatively defined.

One notices that $P^{\tilde{c}\tilde{m}}$ is G -invariant on M . The same is of course true for $P_{(\tilde{c}\tilde{m})}$ being an inverse tensor of $\tilde{P}^{(\tilde{c}\tilde{m})}$. Moreover every G -invariant metric is induced by a scalar product $\langle \cdot, \cdot \rangle$ on $\mathfrak{g}/\mathfrak{g}_0 = \mathfrak{m}$ which is invariant under the action of Ad_{G_0} on $\mathfrak{m}$. Thus P is induced by such a scalar product. The tensor $P_{\tilde{c}\tilde{m}}$ induces on M an additional Einstein–Kaufmann geometry compatible with it. We can repeat some consideration concerning symmetric and skewsymmetric forms on M from section 4. If $P_{[\tilde{d}\tilde{b}]}$ is not degenerate and $\widehat{\nabla}_{\tilde{a}} P_{[\tilde{d}\tilde{b}]} = 0$ we can get an almost complex structure on M induced by $P_{\tilde{c}\tilde{m}} = P_{(\tilde{c}\tilde{m})} + P_{[\tilde{c}\tilde{m}]}$ (if $\dim M$ is even).

Let us consider a linear functional on $L^2 \cap \mathbb{C}^1(M) = D$

$$F_f(g) = \widehat{M}(f, g), \quad g, f \in D \subset L^2, \quad (16.16)$$

$F_f \in (L^2)^* = L^2$ (Riesz theorem). Thus for every f we have a functional F .

Thus we can define a linear operator on L^2 such that

$$L(f) = F_f, \text{ or } (Lf, g) = \widehat{M}(f, g). \quad (16.17)$$

It is easy to see that if $f_1 = f_2$ then $F_{f_1} = F_{f_2}$. Thus L is well defined. Simultaneously we find that $\text{Ker } L = \text{constant functions on } M \cong \mathbb{R}^1$. This is a subspace of L^2 . Thus we can consider L a quotient space of $\mathcal{H}$ and denote it $\widehat{L}$, where

$$\mathcal{H} = L^2 / (\text{Ker } L). \quad (16.18)$$

Let $F_{f_1} = F_{f_2}$ then one gets

$$\bigwedge_{g \in L^2 - (\text{Ker } L)} \widehat{M}(f_1, g) = \widehat{M}(f_2, g). \quad (16.19)$$

Using (16.15) one gets that $f_{1,\tilde{c}} = f_{2,\tilde{c}}$ modulo a set of a zero measure. Thus $[f_1] = [f_2] \in \mathcal{H}$. One can easily check that $\widehat{L}$ is symmetric because $\widehat{M}(f, g) = \widehat{M}(g, f)$.

The operator $\widehat{L}$ is unbounded on D and it is negatively strictly defined i.e.

$$(-\widehat{L}(f), f) \geq C\|f\|^2, \quad (16.20)$$

C is a positive constant.

For this we can apply Friedrichs' theory for $\widehat{L}$. We denote the Friedrichs' space for $\widehat{L}$, $L_0^2(M)$ and the scalar product in $L_0^2(M)$ is given by

$$(f, g)_F = (-\widehat{L}f, g), \quad (16.21)$$

$(D/(\text{Ker } L))$ is dense in $L_0^2(M)$ in a sense of the norm $\|\cdot\|_F$. Let $\tilde{L} = \overline{\hat{L}}$ (a closure of $\hat{L}$) denotes a Hermitian extension of $\hat{L}$ in $L_0^2(M)$, which exists due to Friedrichs' theorem and it is invertible. Moreover $\tilde{L}^{-1}$ is compact.

For this we can proceed a spectral decomposition of $\tilde{L}$. Such an operator has a pure point unbounded spectrum on a negative part of a real axis and eigenvectors (eigenfunctions) span $L_0^2(M)$. One gets

$$\tilde{L} = - \sum_{k=1}^{\infty} \mu_k^2 P_k, \quad \mu_k \neq 0, \mu_k \rightarrow \infty, \quad (16.22)$$

where $P_k^2 = 1$, $P_k P_l = \delta_{kl} I$ are projective operators in $L_0^2(M)$. The sum is understood in a weak topology at a point.

The above considerations justify an intuition of a diagonalization procedure for an infinite matrix $\widehat{M}_{kl}$.

The existence of the tower of scalar fields in our theory can help in renormalization problems. The fields can work as regulator fields if we change some parameters in our theory.

The Friedrichs' theory gives us the following properties of eigenvectors for $\tilde{L}$,

$$\tilde{L} \zeta_k = -\mu_k^2 \zeta_k \quad k = 1, 2, \dots \quad (16.23)$$

They are normalized and orthogonal:

$$\int_M \zeta_k^2(y) \sqrt{|\tilde{g}|} dm(y) = 1, \quad (16.24)$$

$$\int_M \zeta_k(y) \zeta_l(y) \sqrt{|\tilde{g}|} dm(y) = 0 \quad \text{if } k \neq l. \quad (16.25)$$

If the spectrum is degenerated i.e. μ_k^2 is repeated q times (q is a natural number)

$$\mu_k^2 = \mu_{k+1}^2 = \dots = \mu_{k+q-1}^2, \quad (16.26)$$

then $\zeta_k, \zeta_{k+1}, \dots, \zeta_{k+q-1}$ form an algebraic basis of a finite dimensional linear space of functions $f \in L_0^2(M) \cap \mathbb{C}^2(M)$ satisfying Eq. (16.26). After the orthogonalization procedure of $\{\zeta_k, \zeta_{k+1}, \dots, \zeta_{k+q-1}\}$ (for example Schmidt procedure) we can define projectors:

$$P_k = P_{\zeta_k} \text{ for every } k = 1, 2, \dots$$

The orthonormal set of eigenfunctions ζ_k , $k = 1, 2, \dots$ is complete in $L_0^2(M)$. Thus

$$I = \sum_{k=1}^{\infty} P_k = \sum_{k=1}^{\infty} P_{\zeta_k}, \quad (16.27)$$

where I is an identity operator in $L_0^2(M)$. Thus we can expand the field Ψ into a complete set of functions ζ_k

$$\Psi(x, y) = \Psi_0(x) + \sum_{k=1}^{\infty} \Psi_k(x) \zeta_k(y), \quad (16.28)$$

such that

$$\sum_{k=1}^{\infty} \Psi_k^2(x) < \infty, \quad (16.29)$$

for $x \in E$.

Let us find conditions of a diagonalization of the infinite matrix $\widehat{M}_{kl}$. It means we are looking for conditions of the existence of the unitary transformation from the basis χ_k , $k = 1, 2, \dots$ to the basis ζ_k , $k = 1, 2, \dots$ in $\mathcal{H}$ (or in $L_0^2(M)$). Later we give those conditions.

For $\{\zeta_k\}$ is complete one always gets

$$\chi_k = \sum_{l=1}^{\infty} A_{kl} \zeta_l \quad \text{for every } k \quad (16.30)$$

and

$$\sum_{l=1}^{\infty} A_{kl}^2 < \infty. \quad (16.31)$$

Moreover one has

$$A_{kl} = (\chi_k, \zeta_l). \quad (16.32)$$

For the same reasons we have

$$\zeta_k = \sum_{l=1}^{\infty} B_{kl} \chi_l, \quad (16.33)$$

$$\sum_{l=1}^{\infty} B_{kl}^2 < \infty \quad (16.33')$$

and

$$B_{kl} = (\zeta_k, \chi_l). \quad (16.34)$$

One gets

$$A_{kl} = B_{lk} \quad (16.35)$$

and

$$\chi_k = \sum_{l=1}^{\infty} A_{kl} \sum_{p=1}^{\infty} B_{lp} \chi_p = \sum_{p=1}^{\infty} \left(\sum_{l=1}^{\infty} A_{kl} B_{lp} \right) \chi_p. \quad (16.36)$$

This is possible only if

$$\sum_{l=1}^{\infty} A_{kl} B_{lp} = \sum_{l=1}^{\infty} A_{kl} A_{pl} = \delta_{kp}. \quad (16.37)$$

Thus A_{kl} is an isometry in the ℓ^2 -space and B_{kl} either. Moreover A_{kl} and B_{kl} are invertible and A is an inverse operation of B . This means A is an unitary transformation. The transformation A diagonalizes the infinite dimensional matrix $\widehat{M}_{kl}$. The only one condition for the existence of A is the following. The set $\{\chi_k\}$ or $\{\eta_k\}$ are complete bases in $L_0^2(M)$.

Let us consider the operator $\tilde{L}$ in more details and find its shape for $f \in \mathbb{C}^2(M)/\text{Ker } L$. The operator $\tilde{L}$ can be considered a Gateaux derivative of the quadratic form $\widehat{M}$ in $L_0^2(M)$. One finds:

$$\tilde{L}f = \frac{1}{\sqrt{|\tilde{g}|}} (\sqrt{|\tilde{g}|} \tilde{p}^{\tilde{a}\tilde{b}} f_{,\tilde{a}})_{,\tilde{a}} = (\ln \sqrt{|\tilde{g}|})_{,\tilde{b}} \tilde{p}^{\tilde{a}\tilde{b}} f_{,\tilde{a}} + \tilde{p}^{\tilde{a}\tilde{b}}_{,\tilde{b}} f_{,\tilde{a}} + \tilde{p}^{\tilde{a}\tilde{b}} f_{,\tilde{a}\tilde{b}}, \quad (16.38)$$

where

$$\tilde{p}^{\tilde{c}\tilde{m}} = \tilde{p}^{\tilde{m}\tilde{c}} = \left[\tilde{g}^{(\tilde{c}\tilde{m})} + \frac{n^2}{2} (g^{[\tilde{m}\tilde{n}]} g_{\tilde{d}\tilde{n}} \tilde{g}^{(\tilde{d}\tilde{c})} + g^{[\tilde{c}\tilde{n}]} g_{\tilde{d}\tilde{n}} \tilde{g}^{(\tilde{d}\tilde{m})}) \right] \quad (16.39)$$

The operator $\tilde{L}$ slightly differs from the Beltrami–Laplace operator defined on $(M, \bar{p})$.

$$\bar{p} = \bar{p}_{\tilde{a}\tilde{b}} \theta^{\tilde{a}} \otimes \theta^{\tilde{b}}, \quad \bar{p}_{\tilde{a}\tilde{b}} \bar{p}^{\tilde{a}\tilde{c}} = \delta_{\tilde{b}}^{\tilde{c}}$$

because $\det(\bar{p}_{\tilde{a}\tilde{b}}) \neq \det(g_{\tilde{a}\tilde{b}}) = \tilde{g}$. It differs also from the Beltrami–Laplace operator on (M, h^0) .

Moreover $\tilde{L}$ is a left-invariant operator on $C^2(M)$ of the second order. Supposing the reductive decomposition of $\mathfrak{g} = \mathfrak{g}_0 + \mathfrak{m}$ we have a complete description of an algebra of G -invariant operators on M , $D(G/G_0)$ in terms of Lie algebra $\mathfrak{g}$ and $\mathfrak{g}_0$. The algebra is commutative if M is a symmetric space. Let M be a symmetric space (of compact type of course) and let $\text{rank}(M) = K$. The algebra $D(G/G_0)$ has finite numbers of generators i.e. $D_1, D_2, \dots, D_K$. In this case every $D \in D(G/G_0)$ is a symmetric polynomial of D_i , $i = 1, 2, \dots, K$, $D = W(D_1, D_2, \dots, D_K)$. Notice that degrees of D_i , $i = 1, 2, \dots, K$ $d_i = 1, 2, \dots, K$ are canonically established by G .

Thus $\Delta = W_1(D_1, \dots, D_K)$ and $\tilde{L} = W_2(D_1, \dots, D_K, \zeta)$ and $W_1 \neq W_2 W_1(\dots) = W_2(\dots, 0)$.

The existence of the unitary operator A in ℓ^2 -space means that $\widehat{L}$ and $\widehat{\Delta} = \Delta|_{\mathcal{H}}$ have the same Friedrichs' theory i.e. the same $L_0^2(M)$. For M is compact without boundaries and $h^0_{\tilde{a}\tilde{b}} \theta^{\tilde{a}} \otimes \theta^{\tilde{b}}$, $\tilde{g}_{\tilde{a}\tilde{b}} \theta^{\tilde{a}} \otimes \theta^{\tilde{b}}$, $\bar{p}_{\tilde{a}\tilde{b}} \theta^{\tilde{a}} \otimes \theta^{\tilde{b}}$ are smooth functions on M one gets:

$$\overline{m}_2(-\widehat{\Delta}f, f) \leq (-\widehat{L}f, f) \leq \overline{m}_1(-\widehat{\Delta}f, f), \quad (16.40)$$

$\overline{m}_1, \overline{m}_2$ are positive constants.

This means that $L_0^2(M)$ for $\widehat{\Delta}$ and $\widehat{L}$ are equivalent and the conditions we need to imposed for $g_{\tilde{a}\tilde{b}}$ and $p_{\tilde{a}\tilde{b}}$ are

$$\det(g_{\tilde{a}\tilde{b}}) \neq 0 \quad \text{and} \quad \det(\overline{p}_{\tilde{a}\tilde{b}}) \neq 0.$$

These are the only conditions for the existence of the unitary operator A in ℓ^2 -space.

Let us consider $M = S^2$ with the nonsymmetric tensor (6.58). One gets

$$\tilde{L}f = \left(\overline{M} + \frac{n^2\zeta^2}{(\zeta^2 + 1)} \right) \Delta f, \quad (16.41)$$

where Δ is an ordinary Beltrami–Laplace operator on S^2 in spherical coordinates. In this case we do not need any procedure described above, because eigenfunction for $\tilde{L}$ are simply spherical functions $Y_{\ell m}(\theta, \varphi)$ and

$$\tilde{L}Y_{\ell m} = - \left(\overline{M} + \frac{n^2\zeta^2}{(\zeta^2 + 1)} \right) \ell(\ell + 1) Y_{\ell m}. \quad (16.42)$$

The mass spectrum is degenerated for $m = -\ell, -(\ell - 1) \dots 0 \dots \ell - 1, \ell$, $\ell = 1, 2, 3, \dots$ and

$$\overline{m}(\ell, m) = \frac{1}{r} \sqrt{\left(1 + \frac{n^2\zeta^2}{\overline{M}(\zeta^2 + 1)} \right) \ell(\ell + 1)}, \quad \ell = 1, 2, 3. \quad (16.43)$$

Let us notice the following fact: if the homogeneous space $M = G/G_0$ is a two-point homogeneous space the operator $\tilde{L}$ is proportional to the Beltrami–Laplace operator on M . This fact is coming from left-invariancy of $\tilde{L}$ and from this fact that $\tilde{L}$ is of the second order differential operator for $f \in C^{(2)}(M)$. In general this is not true. Moreover for $\zeta = 0$

$$\tilde{L} = \overline{M} \Delta. \quad (16.44)$$

Let us notice that for S^2 (see Eq. (8.71)) we need $\zeta \rightarrow \infty$ in order to kill cosmological constant. Thus $\overline{m}(\ell, m) \simeq \frac{1}{r} \sqrt{2(\ell + 1)\ell}$. We have the following two-point homogeneous spaces $SU(p + 1)/S(U(1) \times U(p))$, $SO(n + 1)/SO(n)$, $(f_{u(-52)}, so(9))$ of compact type, which can serve as the manifold $M = G/G_0$. Only the first one is Hermitian (Kählerian) and in this case one gets

$$\tilde{L} = \left(\overline{M} + \frac{n^2\zeta^2}{(\zeta^2 + 1)} \right) \Delta \quad (16.45)$$

the same formula as for S^2 . Moreover for every two-point symmetric space $\tilde{L}$ is proportional to Δ (Beltrami–Laplace operator)

$$\tilde{L} = \mathfrak{g}(\zeta) \Delta, \quad (16.45a)$$

$$\mathfrak{g}(0) = \overline{M}. \quad (16.45b)$$

It is interesting to find $L_0^2(M)$ in terms of irreducible spaces of representations of the group G . In the case of $S^2 = \text{SO}(3)/\text{SO}(2)$ one simply gets

$$L_0^2(M) = \sum_{\ell=1}^{\infty} \oplus H^\ell, \quad (16.46)$$

where H^ℓ is space of an irreducible representation of the group $\text{SO}(3)$

$$\dim H^\ell = 2\ell + 1.$$

In this case a dynamical group of $\tilde{L}$ is $\text{SO}(3,1)$ (the so called spectrum generation group) $\text{SO}(3) \subset \text{SO}(3,1)$.

$\sum_{\ell=0}^{\infty} \oplus H^\ell = L^2(M) = L_0^2(M) \oplus R$ is a representation space of a unitary representation of $\text{SO}(3,1)$.

In general the situation is more complex.

Let G be a simple compact Lie group and let G_0 be its closed subgroup such that it is a semisimple or $G_0 = U(1) \otimes G_0$, where G_0 is semisimple.

Let $C = h_{ab}Y^aY^b$ be a Casimir operator of G and let C_0 be a Casimir operator of G_0 (or G'_0). Let us suppose a reductive decomposition of the Lie algebra $\mathfrak{g}$, $\mathfrak{g} = \mathfrak{g}_0 + \mathfrak{m}$ (or $\mathfrak{g}_0'$) and consider an operator

$$-D = C - C_0 = h_{\tilde{a}\tilde{b}}Y^{\tilde{a}}Y^{\tilde{b}}. \quad (16.47)$$

This operator acts in the complement $\mathfrak{m}$, which is diffeomorphic to $\text{Tan}_0(G/G_0)$, $0 = \varphi(\varepsilon)$.

Let us define a left-invariant differential operator Δ on M corresponding to D , via a pull-back of the left action of the group G on M . Let us find eigenfunctions of this operator and its eigenvalues. This can be done as follows. Let $H_{\bar{\lambda}_k}$ be invariant spaces of irreducible representations of G corresponding to the value of the Casimir operator, C , $\bar{\lambda}_k$ and $H_{\bar{\mu}_\ell}$ be invariant spaces of irreducible representations of G_0 corresponding to the value of the Casimir operator C_0 , $\bar{\mu}_\ell$. Both groups are compact and those representations are finite-dimensional and unitary. All invariant spaces $H_{\bar{\lambda}_k}$ and $H_{\bar{\lambda}_\ell}$ are Hilbert spaces. One gets

$$L^2(M) = \sum_{\bar{\lambda}_k} \oplus H_{\bar{\lambda}_k} = \sum_{\bar{\lambda}_k} \oplus \sum_{\bar{\mu}_\ell \in \bar{\lambda}_k} \oplus b(\bar{\lambda}_k, \bar{\mu}_\ell) H_{\bar{\mu}_\ell}, \quad (16.48)$$

where $\bar{\mu}_\ell \in \bar{\lambda}_k$, means that $H_{\bar{\lambda}_k}$ is decomposed into some irreducible representation spaces $\bar{\mu}_\ell$ of the subgroup of G , G_0 . $b(\bar{\lambda}_k, \bar{\mu}_\ell)$ means a multiplicity of $H_{\bar{\mu}_\ell}$ in $H_{\bar{\lambda}_k}$. Thus one gets

$$\Delta f_{k,\ell,\ell'} = \bar{\eta}(k, \ell) f_{k,\ell,\ell'}, \quad (16.49)$$

where

$$\bar{\eta}(k, \ell) = \bar{\lambda}_k - \bar{\mu}_\ell < 0, \quad (16.50)$$

$f_{k,\ell,\ell'} \in L^2(M)$ and $f_{k,\ell,\ell'} \in H_{\bar{\mu}_\ell}^{(\ell')}, \bar{\mu}_\ell \in \bar{\lambda}_k$.

$$H_{\bar{\mu}_\ell}^1 \oplus \dots \oplus H^{\ell'} \oplus \dots \oplus H_{\bar{\mu}_\ell}^{b(\bar{\lambda}_k, \bar{\mu}_\ell)} \subset H_{\bar{\lambda}_k}.$$

The dimension of the space corresponding to $\eta_{(k,\ell)}$ can be easily calculated.

$$\dim H_{\bar{\eta}(k,\ell)} = b(\bar{\lambda}_k, \bar{\mu}_\ell) \dim H_{\bar{\mu}_\ell}. \quad (16.51)$$

Thus $\bar{\eta}(k, \ell)$ is in general degenerated. The most important results is this that

$$L^2(M) = \sum_{\bar{\lambda}_k} \oplus H_{\bar{\lambda}_k}, \quad \bar{\lambda} \neq 0 \quad (16.52)$$

or

$$L^2(M) = L^2_0(M) \oplus H_0, \quad \dim H_0 = 1, \quad H_0 \simeq R', \quad (16.52a)$$

or

$$L^2(M) = L^2_0(M)/R'. \quad (16.52b)$$

In the case of a $U(1)$ factor we get similarly

$$\Delta f_{(k,\ell,\ell',m)} = \bar{\eta}(k, \ell) f_{(k,\ell,\ell',m)}, \quad (16.53)$$

where

$$f_{(k,\ell,\ell',m)} = f_{k,\ell,\ell'} \mathfrak{g}_m, \quad m = 0, \pm 1, \pm 2, \dots, \quad (16.54)$$

$\mathfrak{g}_m$ is a function of a one-dimensional irreducible representation of $U(1)$, $m = 0, \pm 1, \pm 2, \dots$. Thus the construction of eigenfunctions of Δ is known from representation theory of G and G_0 (G'_0) and the spectrum is known as well. The interesting problem which arises here is as follows. What is the group (noncompact in general) for which $L^2(M)$ ($L^2_0(M)$) is the invariant space of a unitary, irreducible representation. In other words what is an analogue of $SO(3,1)$ for Δ on S^2 in a general case. This group (if exists) we call dynamical group of Δ , $\widehat{G}$ or a spectrum generating group. We suppose that such a group is minimal for the above requirement. In this way G must be maximally compact subgroup of this group, $G \subset \widehat{G}$. Thus $L^2(M)$ is a space of a unitary representation of $\widehat{G}$ (which is up to now unknown) $T : \widehat{G} \rightarrow L(L^2(M), L^2(M))$ such that

$$T|_G = \sum_{\lambda_k} \oplus H_{\bar{\lambda}_k}, \quad (16.55)$$

where the sum is over all the irreducible representation of G with multiplicity equals one. Such an representation of $\widehat{G}$, T is called maximally degenerated (or most degenerated). Such a situation is possible only if G is a maximal compact subgroup of $\widehat{G}$.

Thus the dynamical group $\widehat{G}$ for Δ is defined as follows:

1. $G \subset \widehat{G}$ and is a maximal compact subgroup $\widehat{G}$ ($\widehat{G}$ is noncompact, of course)

2. $\widehat{G}/G$ is a symmetric irreducible space of noncompact type, such that for a given G its rank is minimal.
3. The maximally degenerated, unitary, irreducible, infinite dimensional representation of $\widehat{G}$, T restricted to G is equivalent to the simple sum of all irreducible representation of the group G (which are finite dimensional).

In our case G is simple and compact. Thus we have the following possibilities:

Space	Rank
$\mathrm{SL}(n, R)/\mathrm{SO}(n)$	$n - 1$
$(\mathfrak{e}_6(\mathfrak{6}), sp(4))$	6
$(\mathfrak{e}_6(-2\mathfrak{6}), \mathfrak{f}_4)$	2
$(\mathfrak{e}_7(\mathfrak{7}), su(8))$	7
$(\mathfrak{e}_8(\mathfrak{8}), so(16))$	8
$(f_4(-2\mathfrak{0}), so(9))$	1

Except the listed above there are also some additional.

$$\mathrm{SO}_0(p, 1)/\mathrm{SO}(p) \quad , \quad 1$$

$\mathrm{SO}_0(3, 1)$ is a Lorentz group and this is our example with S^2 .

$$\mathrm{SU}(p, 1)/\mathrm{U}(p) \quad , \quad 1$$

These are the only possibilities for which we have pairs $(\widehat{G}, G)$ suspected to be a dynamical for Δ on G/G_0 . Thus we need maximally degenerated representations of $\mathrm{SO}_0(p, 1)$, $\mathrm{SU}(p, 1)$, $\mathrm{SL}(n, R)$, $\mathfrak{e}_6(\mathfrak{6})$, $\mathfrak{e}_6(-2\mathfrak{6})$, $\mathfrak{e}_7(\mathfrak{7})$, $\mathfrak{e}_8(\mathfrak{8})$, $\mathfrak{e}_4(-2\mathfrak{0})$ and their decompositions, after a restrictions to the maximally compact subgroup, to the irreducible representations of those subgroups.

Let us consider $\mathrm{SO}_0(p, 1)$. The most degenerate discrete series consists of

$$\widehat{T}(L), \quad L = -\left\{\frac{1}{2}(p+1) - 4\right\}, -\left\{\frac{1}{2}(p+1) - 4\right\} + 1, -\left\{\frac{1}{2}(p+1) - 4\right\} + 2, \dots$$

in $L^2(H^{p,1}, \mu)$.

$H^{p,1}$ — means a hyperboloid in R^{p+1} , $\sum_{i=1}^p (x^i)^2 - (x^{p+1})^2 = 1$ and μ is a measure on $H^{p,1}$, quasiinvariant with respect to the action of the group $\mathrm{SO}_0(p)$ on $H^{p,1}$. In this case after a restriction of $\widehat{T}(L)$ to the subgroup $\mathrm{SO}(p)$ every representation (finite dimensional, unitary, irreducible) enters with a multiplicity equal one (see 2nd position of Ref. [62]).

$$\widehat{T}_{\mathfrak{g}}(L)|_G = \sum_{\ell[p/2]} \oplus \widehat{T}^{\ell[p/2]}, \quad (16.56)$$

where $\widehat{T}^{\ell[p/2]}$ are symmetric finite-dimensional representations of $\text{SO}(p)$ determined by the highest weight

$$\begin{aligned} m &= [\ell_{[p/2]}, 0, 0 \dots], \\ \ell_{[p/2]} &= L + 2n, n = 0, 1, 2, \dots \end{aligned}$$

Thus in the case of $G = \text{SO}(p)$, $\widehat{G} = \text{SO}(p, 1)$.

In the case of $SU(p, 1)$ the situation is even easier. Let us come back to the operator $\widetilde{L}$. It is a G -invariant operator on M of the second order.

Thus it is a linear combination of $\ell = \text{rank } M$, independent operators of on M , left-invariant. If the rank of M is 1 it is proportional to the Beltrami–Laplace operator on M . Thus we have

$$\widetilde{L} = \overline{M} \sum_{i=0}^{\ell-1} \mathfrak{g}_i(\zeta) \Delta_i, \quad (16.57)$$

such that $\Delta_0 = \Delta$ (Beltrami–Laplace operator on M) and

$$\begin{aligned} \mathfrak{g}_0(0) &= 1 \\ \mathfrak{g}_i(0) &= 0, \quad i = 1, 2, \dots, \ell - 1, \end{aligned} \quad (16.58)$$

Δ_i , $i = 1, 2, \dots, \ell - 1$ are remaining G -invariant differential operators on M . Moreover if they commute we can find the spectrum of $\widetilde{L}$ using eigenfunctions of Δ finding spectrum of masses for the tower of scalar fields Ψ_k . The matrix M_{ke} can be diagonalized in $L^2(M, dm)$ which is equivalent to $L^2(\sqrt{\widetilde{g}}, M, dm)$ and to Friedrichs' Hilbert space for Δ . Thus the operator has the same spectrum in $L_0^2(M)$, moreover the eigenfunctions differ in their form from eigenfunctions of Beltrami–Laplace operator due to different scalar product in $L_0^2(M)$.

$$m_k = \frac{1}{2} \sqrt{-2 \sum_{i=0}^{\ell-1} \mu_i \mathfrak{g}_i(\zeta)}. \quad (16.59)$$

The spectrum generating group $\widehat{G}$ is the same for $\widetilde{L}$ as for the Beltrami–Laplace operator on M .

$$\widehat{G} \supset G \supset G_0.$$

Let us remind to the reader that the maximally degenerated representation means that all the Casimir operators are polynomials of the Casimir operator of the lowest order. In our case of the Casimir of the 2nd order. Our group $\widehat{G}$ is a spectrum generated group for the tower of scalar field $\Psi_k(x)$, $(\Psi(x, y))$ on $M \times E$.

Let us come back to the Eqs (5.17–19) and let us release the condition that ρ is independent of y . Thus $\rho = \rho(x, y)$, $y \in G/G_0$. In Eq. (5.34) we get in a place of

$$\frac{\lambda^2}{4} \rho^{n-2} \left(\overline{M} \widetilde{g}^{(\gamma\mu)} \rho_{,\gamma} \rho_{,\mu} + n^2 g^{[\mu\nu]} g_{\delta\mu} \widetilde{g}^{(\delta\gamma)} \right) \rho_{,\nu} \cdot \rho_{,\gamma} \quad (16.60)$$

the formula

$$\frac{\lambda^2}{4} \rho^{n-2} \left(\overline{M} \tilde{\gamma}^{(CM)} \rho_{,C} \rho_{,M} + n^2 \gamma^{[MN]} \gamma_{DM} \tilde{\gamma}^{(DC)} \rho_{,N} \cdot \rho_{,C} \right). \quad (16.61)$$

This ρ has nothing to do with a density of energy considered below.

Now let us repeat the procedure from Section 7, i.e. the redefinition of $g_{\mu\nu}$ and ρ . We get the formula (7.5), but in a place of $\mathcal{L}_{\text{scal}}(\Psi)$ we get the formula (16.3). Simultaneously $\Psi = \Psi(x, y)$, $x \in E$, $y \in G/G_0$ and the metric on a space-time E depends on $y \in G/G_0$, i.e.

$$g_{\mu\nu} = g_{\mu\nu}(x, y), \quad x \in E, \quad y \in G/G_0. \quad (16.62)$$

Thus $g_{\mu\nu}$ is parametrized by a point of G/G_0 . Simultaneously we can interpret a dependence on higher dimensions as an existence of a tower of scalar fields Ψ_K (see Eqs (16.4–8), Eqs (16.11–14) and a discussion below).

The interesting point will be to find physical consequences of this dependence for $g_{\mu\nu}$. This can be achieved by considering cosmological solutions of the theory. Thus let us come to Eq. (7.5) supposing the lagrangian of matter fields is written as L_{matter} , $g_{\mu\nu} = g_{\nu\mu}$ and depends on $y \in G/G_0$. We get

$$\begin{aligned} L\sqrt{-g}\sqrt{|\tilde{g}|} &= \sqrt{-g}\sqrt{|\tilde{g}|} \left(\overline{R}(\tilde{\Gamma}) + \frac{\lambda^2}{4} e^{-(n+2)\Psi} L_{\text{matter}} \right. \\ &\quad \left. + \frac{\lambda^2}{4} \mathcal{L}_{\text{scal}}(\Psi) + \frac{1}{\lambda^2} e^{(n+2)\Psi} \tilde{R}(\tilde{\Gamma}) + \frac{1}{r^2} e^{n\Psi} \tilde{P} \right). \end{aligned} \quad (16.63)$$

According to the standard interpretation of the constant λ we have

$$\frac{\lambda^2}{4} \sim \frac{1}{m_{\text{pl}}^2} \quad (16.64)$$

and

$$\frac{1}{r^2} \sim m_{\tilde{A}}^2 \quad (16.65)$$

where $m_{\tilde{A}}$ is a scale of a mass of broken gauge bosons in our theory and m_{pl} is a Planck mass, $c = 1$, $\hbar = 1$.

Thus one gets form variation principle with respect to $g_{\mu\nu}$ and Ψ

$$\begin{aligned} \tilde{\tilde{R}}_{\mu\nu} - \frac{1}{2} \tilde{\tilde{R}} g_{\mu\nu} &= \frac{e^{-(n+2)\Psi}}{m_{\text{pl}}^2} T_{\mu\nu}^{\text{matter}} + \frac{1}{m_{\text{pl}}^2} T_{\mu\nu}^{\text{scal}} \\ &\quad + \left(\frac{m_{\text{pl}}^2}{8} e^{(n+2)\Psi} \tilde{R}(\tilde{\Gamma}) + m_{\tilde{A}}^2 e^{n\Psi} \tilde{P} \right) g_{\mu\nu} \end{aligned} \quad (16.66)$$

$$\begin{aligned} \overline{M} g^{\mu\nu} \tilde{\nabla}_{\mu} \tilde{\nabla}_{\nu} \Psi + \frac{1}{r^2} \tilde{L} \Psi + \frac{m_{\text{pl}}^4}{4} (n+2) e^{(n+2)\Psi} \tilde{R}(\tilde{\Gamma}) \\ + (m_{\tilde{A}} m_{\text{pl}})^2 n e^{n\Psi} \tilde{P} - (n+2) T e^{-(n+2)\Psi} = 0 \end{aligned} \quad (16.67)$$

where $\widetilde{\widetilde{R}}_{\mu\nu}$ and $\widetilde{\widetilde{R}}$ are a Ricci tensor and a scalar curvature of a Riemannian geometry induced by a metric $g_{\mu\nu}$ (which can depend on $y \in G/G_0$).

$\widetilde{L}$ is an operator on G/G_0 defined by Eq. (16.38), $\widetilde{\nabla}_\mu$ a covariant derivative with respect to a connection induced by $g_{\mu\nu}$ (a Riemannian one).

$$T^{\text{matter}}{}^{\mu\nu} = (p + \varrho)u^\mu u^\nu - pg^{\mu\nu}. \quad (16.68)$$

Let us consider a simple model with a metric

$$ds_4^2 = e^{2v(y)} dt^2 - d\vec{r}^2 \quad (16.69)$$

and let us suppose that $G/G_0 = S^2 = SO(3)/SO(2)$. Thus

$$ds_2^2 = r^2 (d\chi^2 + \sin^2 \chi d\lambda^2), \quad (16.70)$$

$$(\chi, \lambda) = y, \quad x \in \left\langle 0, \frac{\pi}{2} \right\rangle, \quad \lambda \in \langle 0, 2\pi \rangle.$$

In this case $v = v(\chi, \lambda)$ and

$$g_{[\tilde{a}, \tilde{b}]} = g_{[56]} = -\zeta \sin \chi, \quad (16.71)$$

$\widetilde{L}$ is defined by Eq. (16.41),

$$\Psi = \Psi(\vec{r}, t, \chi, \lambda), \quad \varrho = \varrho(\vec{r}, t, \chi, \lambda), \quad p = p(\vec{r}, t, \chi, \lambda). \quad (16.72)$$

Finally let us come to the toy model for which we suppose $\chi = \frac{\pi}{2}$ and we get effectively a 5-dimensional world $E \times S^1$, i.e.

$$ds_5^2 = e^{2v(\lambda)} dt^2 - d\vec{r}^2 - r^2 d\lambda^2. \quad (16.73)$$

One easily gets

$$R_{44} = (v'' + (v')^2) e^v \quad (16.74)$$

$$R_{55} = r^2 (v'' + (v')^2) e^{-v} \quad (16.75)$$

where $'$ means a derivative with respect to λ . For simplicity we suppose $\varrho = p = 0$ and $\Psi = \Psi(\lambda)$ (does not depend on $\vec{r}$ and t). In this way

$$\widetilde{L} = \left(\overline{M} + \frac{n^2 \zeta^2}{\zeta^2 + 1} \right) \frac{\partial^2}{\partial \lambda^2}. \quad (16.76)$$

From Eq. (16.63) one obtains

$$\left(\frac{d^2}{d\lambda^2} e^v \right) e^{-v} = \left(\frac{m_{\text{pl}}^2}{8} e^{(n+2)\Psi} \widetilde{R}(\widetilde{\Gamma}) + m_A^2 e^{n\Psi} \widetilde{P} \right). \quad (16.77)$$

$$m_{\tilde{A}}^2 \left(\overline{M} + \frac{n^2 \zeta^2}{\zeta^2 + 1} \right) \frac{d^2}{d\lambda^2} \Psi + \frac{m_{\text{pl}}^4}{4} (n+2) e^{(n+2)\Psi} \tilde{R}(\tilde{\Gamma}) + (m_{\tilde{A}} m_{\text{pl}})^2 n e^{n\Psi} \tilde{\underline{P}} = 0. \quad (16.78)$$

The last equation can be transformed into

$$\frac{d^2 \Psi}{d\lambda^2} = A e^{(n+2)\Psi} + B e^{n\Psi} \quad (16.79)$$

where

$$A = - \frac{m_{\text{pl}}^4 (n+2)}{4m_{\tilde{A}}^2 \left(\overline{M} + \frac{n^2 \zeta^2}{\zeta^2 + 1} \right)} \tilde{R}(\tilde{\Gamma}) \quad (16.80)$$

$$B = - \frac{m_{\text{pl}}^2 n}{\left(\overline{M} + \frac{n^2 \zeta^2}{\zeta^2 + 1} \right)} \tilde{\underline{P}}. \quad (16.81)$$

Supposing $\Psi = \Psi_0 = \text{const.}$ one gets

$$A e^{2\Psi_0} + B = 0 \quad (16.82)$$

$$e^{\Psi_0} = \sqrt{-\frac{B}{A}} = \frac{4m_{\tilde{A}}}{m_{\text{pl}}} \sqrt{\frac{n|\tilde{\underline{P}}|}{(n+2)\tilde{R}(\tilde{\Gamma})}}. \quad (16.83)$$

Thus from Eq. (16.77) one gets ($z = e^v$)

$$\frac{d^2 z}{d\lambda^2} = \tilde{A} z \quad (16.84)$$

where

$$\begin{aligned} \tilde{A} &= \frac{m_{\text{pl}}^2}{8} e^{(n+2)\Psi_0} \tilde{R}(\tilde{\Gamma}) + m_{\tilde{A}}^2 e^{n\Psi_0} \tilde{\underline{P}} \\ &= m_{\tilde{A}}^2 \left(\frac{4m_{\tilde{A}}}{\alpha_s m_{\text{pl}}} \right)^n \left(\left(\frac{n}{n+2} \right) \frac{|\tilde{\underline{P}}|}{\tilde{R}(\tilde{\Gamma})} \right)^{\frac{n}{2}} \left(\frac{n-2}{n+2} \right) |\tilde{\underline{P}}| > 0. \end{aligned} \quad (16.85)$$

And eventually one gets

$$z = z_0 e^{\sqrt{\tilde{A}}\lambda} + z'_0 e^{-\sqrt{\tilde{A}}\lambda}. \quad (16.86)$$

Taking $z'_0 = 0$ we get

$$e^{2v} = z_0^2 e^{2\sqrt{\tilde{A}}\lambda}. \quad (16.87)$$

Simply rescaling a time in the metric (16.73) we finally get

$$ds_5^2 = e^{2\sqrt{\tilde{A}}\lambda} dt^2 - d\vec{r}^2 - r^2 d\lambda^2. \quad (16.88)$$

In this way we get a funny toy model. We are confined on 3-dimensional brane in a 4-dimensional euclidean space for a $\lambda = 0$. Moreover λ is changing from 0 to 2π resulting

in some interesting possibilities of communication and travel in extra dimensions. This is due to a fact that an effective speed of light depends on λ in an exponential way:

$$c_{\text{eff}} = e^{\sqrt{\bar{A}}\lambda}. \quad (16.89)$$

If we move in λ direction (even a little, remember $r = \frac{1}{m_A}$) from 0 to $\lambda_0 < 2\pi$ and after this move in space direction from $\vec{r}_0$ to $\vec{r}_1$ and again from λ_0 to 0 (we effectively travel from $\vec{r}_0$ to $\vec{r}_1$), then we can be in a point $\vec{r}_1$ even very distant from $\vec{r}_0$ ($L = |\vec{r}_1 - \vec{r}_0|$) in a much shorter time than $\frac{L}{c}$ (where c is the velocity of light taken to be equal 1). In some sense we travel in hyperspace from *Star Wars* or *Wing Commander*.

Thus we consider a metric (see Ref. [116])

$$ds_5^2 = e^{2\sqrt{\bar{A}}\lambda} dt^2 - d\vec{r}^2 - r^2 d\lambda^2. \quad (16.90)$$

We get the following geodetic equations for a signal travelling in $\vec{r}$, λ direction with initial velocity at $\lambda = 0$, $\frac{d\lambda}{dt}(0) = u$, $\frac{d\vec{r}}{dt}(0) = \vec{v}$,

$$\frac{d\vec{r}}{dt} = \vec{v} e^{\sqrt{\bar{A}}\lambda} \quad (16.91)$$

$$\left(\frac{d\lambda}{dt}\right)^2 = e^{\sqrt{\bar{A}}\lambda} - (1 - u^2)e^{2\sqrt{\bar{A}}\lambda}. \quad (16.92)$$

One integrates

$$\vec{r}(\lambda) = \vec{r}_0 - \vec{v}_0 \cdot \frac{2(1 - u^2)^{3/2}}{\sqrt{\bar{A}}} \operatorname{arctg} \left(\sqrt{1 - \frac{1}{(1 - u^2)^{1/2}} e^{-\sqrt{\bar{A}}\lambda}} \right), \quad (16.93)$$

$$\lambda = \frac{1}{\sqrt{\bar{A}}} \ln \left(\frac{4(1 - u^2)^{5/2}}{\bar{A}(t - t_0)^2 + 4(1 - u^2)^3} \right) \quad (16.94)$$

and

$$\vec{r}(t) = \vec{r}_0 - \vec{v}_0 \frac{(1 - u^2)^{3/2}}{\sqrt{\bar{A}}} \operatorname{arctg} \left(\sqrt{1 + \frac{4(1 - u^2)^2}{\bar{A}(t - t_0)^2 + 4(1 - u^2)^3}} \right), \quad (16.95)$$

$$0 < \lambda < \min \left[2\pi, \frac{1}{2\sqrt{\bar{A}}} \ln \left(\frac{1}{1 - u^2} \right) \right]. \quad (16.96)$$

Let us consider a signal travelling along an axis x from 0 to x_0 . The time of this travel is simply x_0 . But now we can consider a travel from zero to λ_0 (in λ direction) and after this from $(0, \lambda_0)$ to (x_0, λ_0) and after to $(x_0, 0)$. The time of this travel is as follows:

$$t = 2t_1 + t_2 \quad (16.97)$$

where

$$t_1 = r \int_0^{\lambda_0} \frac{d\lambda}{e\sqrt{\tilde{A}\lambda}} = \frac{r}{\sqrt{\tilde{A}}} \left(1 - e^{-\sqrt{\tilde{A}\lambda_0}} \right) \quad (16.98)$$

and

$$t_2 = x_0 e^{-\sqrt{\tilde{A}\lambda_0}}. \quad (16.99)$$

For r is of order of a scale of G.U.T. the first term is negligible and

$$t = x_0 e^{-\sqrt{\tilde{A}\lambda_0}}. \quad (16.100)$$

Taking maximal value of $\lambda_0 = 2\pi$ we get

$$t = x_0 e^{-2\pi\sqrt{\tilde{A}}}. \quad (16.101)$$

Thus we achieve really shorter time of a travelling signal through the fifth dimension.

Let us consider the more general situation when we are travelling along a time-like curve via fifth dimension from $\vec{r} = \vec{r}_0$ to $\vec{r} = \vec{r}_1$. Let the parametric equations of the curve have the following shape:

$$\begin{aligned} \vec{r} &= \vec{r}(\xi) \\ t &= t(\xi) \\ \lambda &= \lambda(\xi). \\ 0 &\leq \xi \leq 1, \\ \lambda(0) &= 0, \quad \lambda(1) = \lambda_0, \\ \vec{r}(0) &= \vec{r}_0 \\ \vec{r}(1) &= \vec{r}_1. \end{aligned} \quad (16.102)$$

Let a tangent vector to the curve be $(\dot{t}(\xi), \dot{\vec{r}}(\xi), \dot{\lambda}(\xi))$, where dot “ $\dot{}$ ” means a derivative with respect to ξ . For a total time measured by a local clock of an observer one gets

$$\Delta t = \int_0^1 d\xi \left(e^{2\sqrt{\tilde{A}\lambda}(\xi)} \left(\frac{dt}{d\xi} \right)^2 - \left(\frac{d\vec{r}}{d\xi} \right)^2 - r^2 \left(\frac{d\lambda}{d\xi} \right)^2 \right)^{\frac{1}{2}}. \quad (16.104)$$

Let us consider a three segment curve such that

- 1) a straight line from $\vec{r} = 0, \lambda = 0$ to $\vec{r} = 0, \lambda = \lambda_0$,
- 2) a straight line from $\vec{r} = 0, \lambda = \lambda_0$ to $\vec{r} = \vec{r}_0, \lambda = \lambda_0$,
- 3) a straight line from $\vec{r} = \vec{r}_0, \lambda = \lambda_0$ to $\vec{r} = \vec{r}_0, \lambda = 0$.

The additional assumptions are such that the traveller travels with a constant velocity v_λ on the first segment and on the third segment and with a velocity $\vec{v}$ on the second one.

Thus we have a parametrization:

1) On the first segment

$$\begin{aligned}\vec{r} &= 0 \\ t &= \frac{\lambda_0}{v_\lambda} \xi, \quad 0 \leq \xi \leq 1 \\ \lambda &= \xi \lambda_0.\end{aligned}\tag{16.105}$$

2) On the second segment

$$\begin{aligned}\vec{r}(\xi) &= \vec{r}_0 \xi \\ t &= \frac{|\vec{r}_0|}{|\vec{v}|} \xi, \quad 0 \leq \xi \leq 1 \\ \lambda &= \lambda_0.\end{aligned}\tag{16.106}$$

3) On the third segment

$$\begin{aligned}\vec{r} &= \vec{r}_0 \\ t &= \frac{\lambda_0}{|\vec{v}|} (1 - \xi), \quad 0 \leq \xi \leq 1 \\ \lambda &= (1 - \xi) \lambda_0.\end{aligned}\tag{16.107}$$

From Eq. (16.104) one easily gets

$$\Delta t = \Delta t_1 + \Delta t_2 \tag{16.108}$$

$$\begin{aligned}\Delta t_1 &= \frac{2r}{\sqrt{\tilde{A}}} \left(\frac{1}{\sqrt{v_\lambda \lambda_0}} \left(\sqrt{e^{2\sqrt{\tilde{A}}\lambda_0} - v_\lambda \lambda_0} - \sqrt{1 - v_\lambda \lambda_0} \right) \right. \\ &\quad \left. + \operatorname{arctg} \left(\sqrt{\frac{e^{2\sqrt{\tilde{A}}\lambda_0} - v_\lambda \lambda_0}{v_\lambda \lambda_0}} \right) - \operatorname{arctg} \left(\sqrt{\frac{1 - v_\lambda \lambda_0}{v_\lambda \lambda_0}} \right) \right)\end{aligned}\tag{16.109}$$

$$\Delta t_2 = \frac{|\vec{r}_0|}{|\vec{v}|} \sqrt{e^{2\sqrt{\tilde{A}}\lambda_0} - |\vec{v}|^2} \tag{16.110}$$

For r is of order of an inverse scale of G.U.T., the first term in Eq. (16.108) is negligible and

$$\Delta t \simeq \frac{|\vec{r}_0|}{|\vec{v}|} \sqrt{e^{2\sqrt{\tilde{A}}\lambda_0} - |\vec{v}|^2}. \tag{16.111a}$$

The traveller passing from $\vec{r} = 0$ to $\vec{r} = \vec{r}_0$ (having $\lambda = 0$ during his travel) reached $\vec{r} = \vec{r}_0$ at time

$$\Delta t' = \frac{|\vec{r}_0|}{|\vec{v}|} (1 - |\vec{v}|^2)^{1/2}. \tag{16.111b}$$

The first traveller has the following condition for $\vec{v}$:

$$|\vec{v}| \leq e^{\sqrt{\tilde{A}}\lambda_0}, \quad 0 \leq \lambda_0 \leq 2\pi. \tag{16.112a}$$

The second

$$|\vec{v}| \leq 1. \quad (16.112b)$$

Thus we see that we can travel quicker taking a bigger $\vec{v}$.

For an unmoving observer the time needed for a travel in the first case is

$$\begin{aligned} \Delta t &= \frac{2r\lambda_0}{v_\lambda} + \frac{|\vec{r}_0|}{|\vec{v}|} \simeq \frac{|\vec{r}_0|}{|\vec{v}|}, \\ |\vec{v}| &\leq e^{\sqrt{\tilde{A}}\lambda_0}, \quad 0 \leq \lambda_0 \leq 2\pi. \end{aligned} \quad (16.111a*)$$

In the second case

$$\begin{aligned} \Delta t' &= \frac{|\vec{r}_0|}{|\vec{v}|}, \\ |\vec{v}| &\leq 1 \end{aligned} \quad (16.111b*)$$

Thus in some sense we get a dilatation of time:

$$\Delta t' = \left(e^{2\sqrt{\tilde{A}}\lambda_0} - \vec{v}^2 \right)^{1/2} \Delta t \quad (16.111a**) \quad (16.111b**)$$

in the first case,

$$\Delta t' = (1 - \vec{v}^2)^{1/2} \Delta t \quad (16.111b**)$$

in the second case.

Let us come back to the Eqs (16.94–96). We see that a particle coming along a geodesic is confined in a shell surrounding a brane in the fifth dimension

$$0 \leq \lambda \leq \frac{r}{\sqrt{\tilde{A}}} \ln \left(\frac{1}{(1 - u^2)^{1/2}} \right). \quad (16.113)$$

The particle oscillates in the shell. Moreover $0 \leq \lambda \leq 2\pi$, thus if λ excesses 2π , we take for a new value $\lambda - 2\pi$ (i.e. modulo 2π). Usually for u not close to 1 this is a really thin shell of order r (a length of a unification scale).

Let us come back to our toy model for a cone with $\tilde{A} = 0$. In this case an equation of cone in 5-dimensional Minkowski space is

$$x^2 + y^2 + z^2 + r^2\lambda^2 = t^2. \quad (16.114)$$

An interesting question is what is an analog for (16.114) if $\tilde{A} \neq 0$. One easily gets that

$$x^2 + y^2 + z^2 + r^2\lambda^2 = t^2 r^2 \lambda^2 \tilde{A}^2 e^{2\sqrt{\tilde{A}}\lambda}. \quad (16.115)$$

There is no simple way to get Eq. (16.114) from Eq. (16.115). Moreover if we change a coordinate t into $rt\lambda\tilde{A}$ one gets

$$x^2 + y^2 + z^2 + r^2\lambda^2 = t^2 e^{2\sqrt{\tilde{A}}\lambda}. \quad (16.116)$$

and for $\tilde{A} = 0$ we get Eq. (16.114).

Let us notice that Eq. (16.116) is necessary to be considered for λ modulo 2π . It means even λ could be any real number in the equation we should put in a place of λ , $[\lambda]$, where $[\lambda]$ is defined as follows

$$\lambda = [\lambda] + 2\pi n, \quad [\lambda] \in \langle 0, 2\pi \rangle,$$

n is an integer and $\lambda \in (-\infty, +\infty)$. In this way in a place of Eq. (16.116) we get

$$x^2 + y^2 + z^2 + r^2[\lambda]^2 = t^2 e^{2\sqrt{\bar{A}}[\lambda]}. \quad (16.116*)$$

The last equation defines an interesting 4-dimensional hypersurface in 5-dimensional space ($\mathbb{R}^5$).

For $\lambda = 0$ we get a light cone on a brane (4-dimensional Minkowski space). For branes with $0 < \lambda = \lambda_0 < 2\pi$ we have a “cone”

$$x^2 + y^2 + z^2 = t^2 e^{2\sqrt{\bar{A}}\lambda_0} - \lambda_0^2 \quad (16.117)$$

which is a 3-dimensional hyperboloid.

In our model we get in a quite natural way a warp factor known from Ref. [117]. Moreover we have a different reason to get it. The reason of this warp factor is a many dimensional dependence of a scalar field $\Psi = \Psi(x, \lambda)$. In particular higher-dimensional excitations of Ψ forming a tower of massive scalar fields. In this model Ψ can be developed in a Fourier series

$$\Psi(x, \lambda) = \sum_{m=0}^{\infty} \psi_m^1(x) \cos(m\lambda) + \sum_{m=1}^{\infty} \psi_m^2(x) \sin(m\lambda) \quad (16.118)$$

where

$$\psi_m^1(x) = \frac{1}{2\pi} \int_0^{2\pi} \Psi(x, \lambda) \cos(m\lambda) d\lambda \quad (16.119)$$

$$\psi_m^2(x) = \frac{1}{2\pi} \int_0^{2\pi} \Psi(x, \lambda) \sin(m\lambda) d\lambda \quad (16.120)$$

$$m = 1, 2, \dots$$

$$\frac{\psi_0^1(x)}{2} = \frac{1}{2\pi} \int_0^{2\pi} \Psi(x, \lambda) d\lambda. \quad (16.121)$$

Thus an interesting point will be to find a spectrum of mass for this tower. This is simply

$$m_m^2 = m_A^2 \left(\bar{M} + \frac{n^2 \zeta^2}{\zeta^2 + 1} \right) m^2, \quad m = 0, 1, 2, \dots \quad (16.122)$$

Now we come to the calculation of a content of a warp factor (a content in terms of a tower of scalar fields). Thus we calculate

$$\psi_m^1 = \frac{1}{2\pi} \int_0^{2\pi} e^{2\sqrt{\tilde{A}}\lambda} \sin(m\lambda) d\lambda, \quad m = 1, 2, 3, \dots \quad (16.123)$$

$$\psi_m^2 = \frac{1}{2\pi} \int_0^{2\pi} e^{2\sqrt{\tilde{A}}\lambda} \cos(m\lambda) d\lambda, \quad m = 1, 2, 3, \dots \quad (16.124)$$

$$\frac{1}{2}\psi_0^1 = \frac{1}{2\pi} \int_0^{2\pi} e^{2\sqrt{\tilde{A}}\lambda} d\lambda. \quad (16.125)$$

One gets

$$\frac{1}{2}\psi_0^1 = \frac{1}{4\pi\sqrt{\tilde{A}}} \left(e^{4\pi\sqrt{\tilde{A}}} - 1 \right) \quad (16.126)$$

$$\psi_m^1 = -\frac{m}{2\pi} \frac{(e^{4\pi\sqrt{\tilde{A}}} - 1)}{(4\tilde{A} + m^2)} \quad (16.127)$$

$$\psi_m^2 = \frac{\sqrt{\tilde{A}}(e^{4\pi\sqrt{\tilde{A}}} - 1)}{\pi(4\tilde{A} + m^2)}. \quad (16.128)$$

The simple question is what is the energy to excite the warp factor in terms of a tower of scalar fields. The answer is as follows:

$$E_{\text{wf}} = \int_0^{2\pi} T_{44}^{\text{scal}}(e^{2\sqrt{\tilde{A}}\lambda}) d\lambda \quad (16.129)$$

where T_{44}^{scal} is a time component of an energy-momentum tensor for a scalar field calculated for $\Psi = e^{2\sqrt{\tilde{A}}\lambda}$.

One gets

$$T_{44}^{\text{scal}} = \frac{1}{2} \left(\frac{d\Psi}{d\lambda} \right)^2 + \frac{1}{2} \lambda_{c_0}(\Psi), \quad \lambda_{c_0}(\Psi) = \frac{1}{2} |\gamma| e^{n\Psi} - \frac{\beta}{2} e^{(n+2)\Psi} \quad (16.130)$$

$$\int_0^{2\pi} \frac{1}{2} \left(\frac{d}{d\lambda} e^{2\sqrt{\tilde{A}}\lambda} \right)^2 d\lambda = \frac{\sqrt{\tilde{A}}}{2} \left(e^{8\pi\sqrt{\tilde{A}}} - 1 \right) \quad (16.131)$$

and

$$\frac{1}{2} \lambda_{c_0}(e^{2\sqrt{\tilde{A}}\lambda}) = -\frac{1}{2} e^{n(e^{2\sqrt{\tilde{A}}\lambda})} \left(\beta e^{2(e^{2\sqrt{\tilde{A}}\lambda})} - |\gamma| \right), \quad (16.132)$$

where

$$\gamma = \frac{m_{\tilde{A}}^2}{\alpha_s^2} \tilde{P} = \frac{1}{r^2} \tilde{P},$$

$$\beta = \frac{\alpha_s^2}{l_{\text{pl}}^2} \tilde{R}(\tilde{\Gamma}) = \alpha_s^2 m_{\text{pl}}^2 \tilde{R}(\tilde{\Gamma}).$$

One easily gets

$$E_{\text{wf}} = \frac{\sqrt{\tilde{A}}}{2} (e^{8\pi\sqrt{\tilde{A}}} - 1) - \frac{1}{4\sqrt{\tilde{A}}} \left[|\gamma| \left(\text{Li} \left(\exp \left(\frac{e^{4\pi\sqrt{\tilde{A}}}}{n} \right) \right) - \text{Li}(\sqrt[n]{e}) \right) \right. \\ \left. - \beta \left(\text{Li} \left(\exp \left(\frac{e^{4\pi\sqrt{\tilde{A}}}}{(n+2)} \right) \right) - \text{Li}(\sqrt[n+2]{e}) \right) \right] \quad (16.133)$$

where

$$\text{Li}(x) = \int_0^x \frac{dz}{\ln z}, \quad x > 1, \quad (16.134)$$

is an integral logarithm. The integral is considered in the sense of a principal value.

Let us consider Eq. (16.133) using asymptotic formula for $\text{Li}(x)$ for large x :

$$\text{Li}(x) \sim x \left[\frac{1}{\ln x} + \sum_{m=1}^{\infty} \frac{m!}{\ln^{m+1} x} \right].$$

One gets

$$E_{\text{wf}} = \frac{\sqrt{\tilde{A}}}{2} (e^{8\pi\sqrt{\tilde{A}}} - 1) - \frac{e^{-4\pi\sqrt{\tilde{A}}}}{4\sqrt{\tilde{A}}} \left(n|\gamma| \exp \left(\frac{e^{4\pi\sqrt{\tilde{A}}}}{n} \right) \right. \\ \left. - (n+2)\beta \exp \left(\frac{e^{4\pi\sqrt{\tilde{A}}}}{n+2} \right) \right). \quad (16.133a)$$

However, let us think quantum-mechanically in a following direction. Let us create (maybe in an accelerator as some kind of resonances) scalar particles of mass m_n (Eq. (16.122)) with an amount of $|\psi_m^1|^2 + |\psi_m^2|^2$. In this way we create the full warp factor (for they due to interference will create it as a Fourier series). The question is what is an energy needed to create those particles. The answer is simply

$$\overline{E} = \sum_{m=1}^{\infty} (|\psi_m^1|^2 + |\psi_m^2|^2) m_m. \quad (16.135)$$

One easily gets

$$\overline{E} = \frac{m_{\tilde{A}}}{4\pi^2} \left(e^{4\pi\sqrt{\tilde{A}}} - 1 \right)^2 \left(\overline{M} + \frac{n^2\zeta^2}{\zeta^2 + 1} \right)^{\frac{1}{2}} \sum_{m=1}^{\infty} \frac{m}{4\tilde{A} + m^2}. \quad (16.136)$$

The series $\sum_{m=1}^{\infty} \frac{m}{4\tilde{A} + m^2}$ is divergent for

$$\frac{m}{4\tilde{A} + m^2} \simeq \frac{1}{m}. \quad (16.137)$$

Thus we need a regularization technique to sum it. We use ζ -function regularization technique similar as in Casimir-effect theory. Let us introduce a $\zeta_{\tilde{A}}$ -function on a complex plane

$$\zeta_{\tilde{A}} = \sum_{m=1}^{\infty} \left(\frac{m}{4\tilde{A} + m^2} \right)^s \quad (16.138)$$

for $\text{Re}(s) \geq 1 + \delta$, $\delta > 0$.

This function can be extended analytically on a whole complex plane. Obviously $\zeta_{\tilde{A}}$ has a pole for $s = 1$ (as a Riemann ζ function). Thus we should regularize it at $s = 1$. One gets

$$\overline{E} = \frac{m_{\tilde{A}}}{4\pi^2} \left(e^{4\pi\sqrt{\tilde{A}}} - 1 \right)^2 \left(\overline{M} + \frac{n^2\zeta^2}{\zeta^2 + 1} \right)^{\frac{1}{2}} \zeta_{\tilde{A}}^r(1) \quad (16.139)$$

where $\zeta_{\tilde{A}}^r$ means a regularized $\zeta_{\tilde{A}}$.

Let us introduce a ζ -Riemann function

$$\zeta_R(s) = \sum_{m=1}^{\infty} \frac{1}{m^s}, \quad \text{Re}(s) \geq 1 + \delta, \quad \delta > 0. \quad (16.140)$$

One can write

$$“\zeta_{\tilde{A}}(1)” = “\zeta_R(1)” + 4\tilde{A} \sum_{m=1}^{\infty} \frac{1}{m(4\tilde{A} + m^2)} \quad (16.141)$$

or

$$“\zeta_{\tilde{A}}(1)” = “\zeta_R(1)” + 4\tilde{A}\zeta_R(3) - 16\tilde{A}^2 \sum_{m=1}^{\infty} \frac{1}{m^3(4\tilde{A}^2 + m^2)}. \quad (16.142)$$

The series in Eqs (16.141–142) are convergent. Moreover $\zeta_R(s)$ has a pole at $s = 1$ and

$$\lim_{s \rightarrow 1} \left[\zeta_R(s) - \frac{1}{s-1} \right] = \gamma_E \quad (16.143)$$

where γ_E is an Euler constant.

Thus we can regularize $\zeta_{\tilde{A}}(s)$ in such a way that

$$\zeta_{\tilde{A}}^r(s) = \zeta_R(s) - \frac{1}{s-1} + 4\tilde{A} \sum_{m=1}^{\infty} \frac{1}{m(4\tilde{A} + m^2)}. \quad (16.144)$$

Thus

$$\zeta_{\tilde{A}}^r(1) = \gamma_E + 4\tilde{A} \sum_{m=1}^{\infty} \frac{1}{m(4\tilde{A} + m^2)}. \quad (16.145)$$

The series

$$\sum_{m=1}^{\infty} \frac{1}{m(4\tilde{A} + m^2)} \quad (16.146)$$

is convergent for every $z^2 = -4\tilde{A}$ and defines an analytic function

$$f(z) = \sum_{m=1}^{\infty} \frac{1}{m(m^2 - z^2)}. \quad (16.147)$$

Using some properties of an expansion in simple fraction for

$$\psi(x) = \frac{d}{dx} \log \Gamma(x) \quad (16.148)$$

where $\Gamma(x)$ is an Euler Γ -function, one gets

$$\psi(x) = -\gamma_E - \sum_{m=0}^{\infty} \left(\frac{1}{m+x} - \frac{1}{m+1} \right) \quad (16.149)$$

$$\zeta_{\tilde{A}}^r(1) = -\frac{1}{2} \left(\psi(2\sqrt{\tilde{A}}) + \psi(-2\sqrt{\tilde{A}}) \right) \quad (16.150)$$

or

$$\zeta_{\tilde{A}}^r(1) = \frac{1}{4\sqrt{\tilde{A}}} + \frac{\pi}{2} \operatorname{ctg} 2\sqrt{\tilde{A}}\pi \quad (16.151)$$

(where we use $\Gamma(z)\Gamma(-z) = -\frac{\pi}{z \sin \pi z}$).

And finally

$$\overline{E} = \frac{m_{\tilde{A}}}{8\pi^2} \left(\overline{M} + \frac{n^2 \zeta^2}{\zeta^2 + 1} \right) \left(e^{4\pi\sqrt{\tilde{A}}} - 1 \right)^2 \left(\frac{1}{2\sqrt{\tilde{A}}} + \pi \operatorname{ctg}(2\sqrt{\tilde{A}}\pi) \right). \quad (16.152)$$

It is easy to see that $\overline{E} = 0$ for $\tilde{A} = 0$,

$$\sqrt{\tilde{A}} = m_{\tilde{A}} \left(\frac{4\sqrt{n} m_{\tilde{A}}}{\alpha_s \sqrt{n+2} m_{\text{pl}}} \right)^{\frac{n}{2}} \left(\frac{|\tilde{P}|}{\tilde{R}(\tilde{\Gamma})} \right)^{\frac{n}{4}} \left(\frac{n-2}{n+2} |\tilde{P}| \right)^{\frac{1}{2}}. \quad (16.153)$$

For we use a dimensionless coordinate λ (not $\tilde{\lambda} = r \cdot \lambda = \frac{\lambda}{m_{\tilde{A}}}$), we can omit a factor $m_{\tilde{A}}$ in front of the right-hand side in the formula (16.153).

It is interesting to notice that for $\tilde{A} = 0$, $e^{2\sqrt{\tilde{A}}} = 1$ and we have to do with a Fourier expansion of a constant mode only ($m = 0$). This mode is a massless mode. However, a massless mode can obtain a mass from some different mechanism (as in Section 14). Thus in some sense we have to do with an energy

$$\frac{1}{4} \left| \psi_0^1(\tilde{A} = 0) \right|^2 m_\psi = m_\psi . \quad (16.154)$$

One gets the following formula

$$m_\psi = m_{\text{pl}} \sqrt{\frac{|\tilde{P}|(n+2)}{(\tilde{M} + \frac{n^2 \zeta^2}{\zeta^2 + 1})}} \left(\frac{4\sqrt{n} \sqrt{|\tilde{P}|} m_{\tilde{A}}}{\alpha_s \sqrt{n+2} \sqrt{\tilde{R}(\tilde{\Gamma})} m_{\text{pl}}} \right)^{\frac{n}{2}} . \quad (16.155)$$

Thus it seems that in order to create a warp factor $e^{2\sqrt{\tilde{A}}\lambda}$ in a front of dt^2 we should deposite an energy of a tower of scalar particles

$$\tilde{E} = E + m_\psi . \quad (16.156)$$

Even a factor equal to 1 is not free. This is of course supported by a classical field formula (Eq. (16.134)). For $\tilde{A} = 0$ we get

$$E_{\text{wf}} = -\pi \cdot e^n (\beta e^2 - |\gamma|) \quad (16.157)$$

or

$$E_{\text{wf}} = -\frac{\pi e^n}{\alpha_s^2} \left(e^2 \alpha_s^4 m_{\text{pl}}^2 \tilde{R}(\tilde{\Gamma}) - m_{\tilde{A}}^2 |\tilde{P}| \right) . \quad (16.158)$$

To be honest we should multiply E_{wf} and E by V_3 , where V_3 is a volume of a space and a result will be divergent. Moreover, if we want to excite a warp factor only locally, V_3 can be finite and the result also finite.

Let us come back to the Eq. (16.63). The variational principle based on (16.63) lagrangian is really in $(n_1 + 4)$ -dimensional space $E \times G/G_0$, where a size of a compact space $M = G/G_0$ is r . In this way the gravity lives on $(n_1 + 4)$ -dimensional manifold, where n_1 dimensions are curled into a compact space ([117]). The scalar field Ψ lives also on $(n_1 + 4)$ -dimensional manifold. The matter is 4-dimensional. Thus we can repeat some conclusions from Randall-Sundrum ([118]) and Arkani-Hamed, Savas Dimopoulos and Dvali ([119]) theory. Similarly as in their case we have gravity for two regions ($V(R)$ is a Newtonian gravitational potential).

1) $R \ll r$

$$V(R) \cong \frac{m_1 \cdot m_2}{m_{\text{pl}(n_1+4)}^{n_1+2}} \frac{1}{R^{n_1+1}} , \quad (16.159)$$

2) $R \gg r$

$$V(R) \cong \frac{m_1 \cdot m_2}{m_{\text{pl}(n_1+4)}^{n_1+2} r^{n_1}} \frac{1}{R} , \quad (16.160)$$

where $m_{\text{pl}(n_1+4)}$ is a $(n_1 + 4)$ -dimensional Planck's mass. So

$$m_{\text{pl}}^2 = m_{\text{pl}(n_1+4)}^{n_1+2} \cdot r^{n_1} = \frac{m_{\text{pl}(n_1+4)}^{n_1+2}}{m_{\tilde{A}}^{n_1}} = \left(\frac{m_{\text{pl}(n_1+4)}}{m_{\tilde{A}}} \right)^{n_1} \cdot m_{\text{pl}(n_1+4)}^2. \quad (16.161)$$

Thus

$$m_{\text{pl}(n_1+4)} = (m_{\text{pl}})^{\frac{2}{n_1+2}} \cdot (m_{\tilde{A}})^{\frac{n_1}{n_1+2}}. \quad (16.162)$$

Eqs (16.161–162) give us a constraint on parameters in our theory and establish a real strength of gravitational interactions given by $m_{\text{pl}(n_1+4)}$.

However, in our case we have to do with a scalar-tensor theory of gravity. Thus our gravitational constant is effective

$$G_{\text{eff}} = G_N e^{-(n+2)\Psi} = G_N \rho^{n+2}. \quad (16.163)$$

In this case we have to do with effective “Planck's masses” in $n_1 + 4$ and 4 dimensions

$$m_{\text{pl}(n_1+4)}^2 \longrightarrow m_{\text{pl}(n_1+4)}^2 \cdot e^{(n+2)\Psi_{(n_1+4)}} \quad (16.164)$$

$$m_{\text{pl}}^2 \longrightarrow m_{\text{pl}}^2 \cdot e^{(n+2)\Psi} \quad (16.165)$$

where $\Psi_{(n_1+4)}$ is a scalar field Ψ for $R \ll r$ and Ψ is for $R \gg r$.

In this way Eq. (16.161) reads

$$m_{\text{pl}}^2 e^{(n+2)\Psi} = \left(\frac{m_{\text{pl}(n_1+4)} e^{\frac{(n+2)}{2}\Psi_{(n_1+4)}}}{m_{\tilde{A}}} \right)^{n_1} \cdot m_{\text{pl}(n_1+4)}^2 \cdot e^{(n+2)\Psi_{(n_1+4)}}. \quad (16.166)$$

For our contemporary epoch we can take $e^{(n+2)\Psi} \simeq 1$.

If we take as in the case of Ref. [119]

$$m_{\text{pl}(n_1+4)} \simeq m_{\text{EW}}, \quad (16.167)$$

where m_{EW} is an electro-weak energy scale. We get

$$\left(\frac{m_{\text{pl}}}{m_{\text{EW}}} \right)^2 = \left(\frac{m_{\text{EW}}}{m_{\tilde{A}}} \right)^{n_1} \cdot e^{\frac{1}{2}(n+2)(n_1+2)\Psi_{(n_1+4)}}. \quad (16.168)$$

For $R \ll r$ the field $\Psi_{(n_1+4)}$ has the following behaviour

$$\Psi_{(n_1+4)} \cong \Psi_0 + \frac{\alpha}{R^{n_1+1}}. \quad (16.169)$$

Thus we can approximate $\Psi_{(n_1+4)}$ by Ψ_0 where Ψ_0 is a critical point of a selfinteracting potential for Ψ . In this way one gets

$$\left(\frac{m_{\text{pl}}}{m_{\text{EW}}} \right) \left(\frac{m_{\tilde{A}}}{m_{\text{EW}}} \right)^{n_1} = \exp \left(\frac{(n+2)(n_1+2)}{2} \Psi_0 \right) \quad (16.170)$$

where Ψ_0 is given by Eq. (14.204). Thus one gets:

$$\begin{aligned} & \left(\frac{m_{\text{pl}}}{m_{\text{EW}}} \right) \left(\frac{m_{\tilde{A}}}{m_{\text{EW}}} \right)^{n_1} \left(\frac{m_{\text{pl}}}{m_{\tilde{A}}} \right)^{\frac{(n+2)(n_1+2)}{2}} \\ &= \left(\frac{1}{\alpha_s} \right)^{\frac{(n+2)(n_1+2)}{2}} \left(\frac{n|\underline{\tilde{P}}|}{(n+2)|\tilde{R}(\tilde{\Gamma})|} \right)^{\frac{(n+2)(n_1+2)}{4}}. \end{aligned} \quad (16.171)$$

It is easy to see that we can achieve a solution for a hierarchy problem if $\left| \frac{\underline{\tilde{P}}}{\tilde{R}(\tilde{\Gamma})} \right|$ is sufficiently big, e.g. taking $\tilde{R}(\tilde{\Gamma})$ sufficiently small and/or $\underline{\tilde{P}}$ sufficiently big playing with constants ξ and ζ . In this way a real strength of gravity will be the same as electro-weak interactions and the hierarchy problem has been reduced to the problem of smallness of a cosmological constant (i.e. $\tilde{R}(\tilde{\Gamma}) \simeq 0$).

Let us consider Eq. (16.152) in order to find such a value of $\tilde{A}$ for which $\overline{E}$ is minimal. One finds that

$$\overline{E} = 0 \quad (16.172)$$

for

$$\tilde{A} = \frac{x^2}{4\pi^2} \quad (16.173)$$

where x satisfies an equation

$$x = -\text{tg } x, \quad x > 0. \quad (16.174)$$

Eq. (16.174) has an infinite number of roots. One can find some roots of Eq. (16.174) for $x > 0$. We get

$$\begin{aligned} x_1 &= 4.913\dots \\ x_2 &= 7.978\dots \\ x_3 &= 11.086\dots \\ x_4 &= 14.207\dots \end{aligned} \quad (16.175)$$

We are interested in large roots (if we want to have c_{eff} considerably big).

In this case one finds

$$x = \varepsilon + \frac{\pi}{2}(2l+1) \quad (16.176)$$

where $\varepsilon > 0$ is small and $l = 1, 2, \dots$. Moreover, to be in line in our approximation, l should be large. One gets

$$\frac{\pi}{2}(2l+1) + \varepsilon = \text{ctg } \varepsilon. \quad (16.177)$$

For ε is small, one gets

$$\text{ctg } \varepsilon \simeq \frac{1}{\varepsilon}. \quad (16.178)$$

Finally one gets

$$\varepsilon^2 + (2l+1)\frac{\pi}{2} - 1 = 0 \quad (16.179)$$

and

$$\varepsilon \simeq \frac{2}{(2l+1)\pi} \quad (16.180)$$

$$x = \frac{\pi}{2}(2l+1) + \frac{2}{(2l+1)\pi} \quad (16.180a)$$

$$\tilde{A} = \frac{1}{4\pi^2} \left(\frac{\pi}{2}(2l+1) + \frac{2}{(2l+1)\pi} \right)^2. \quad (16.181)$$

It is easy to notice that our approximation (16.180a) is very good even for small l , e.g. for $l = 4$ we get x_4 (Eq. (16.175)).

In this way an effective velocity of light can be arbitrarily large,

$$c_{\text{eff}}^{\text{max}} = c \exp \left(\frac{1}{2} \left(\frac{\pi}{2}(2l+1) + \frac{2}{(2l+1)\pi} \right) \right) \quad (16.182)$$

(i.e. for $\lambda = 2\pi$). However, in order to get such large c_{eff} we should match Eq. (16.153) with (16.181). One gets

$$\frac{\pi}{2}(2l+1) + \frac{2}{(2l+1)\pi} = 2\pi \left(\frac{4\sqrt{n}m_{\tilde{A}}}{\alpha_s\sqrt{n+2}m_{\text{pl}}} \right)^{\frac{n}{2}} \left(\frac{|\underline{\tilde{P}}|}{\tilde{R}(\tilde{\Gamma})} \right)^{\frac{n}{4}} \left(\frac{n-2}{n+2} |\underline{\tilde{P}}| \right)^{\frac{1}{2}}. \quad (16.183)$$

In this way we get some interesting relations between parameters in our theory. To have a large c_{eff} we need to make $|\underline{\tilde{P}}|$ large and $\tilde{R}(\tilde{\Gamma})$ small. In this case only the first term of the left-hand side of (16.183) is important and we should play with the ratio $\left(\frac{m_{\tilde{A}}}{\alpha_s m_{\text{pl}}} \right)$ in order to get the relation

$$(2l+1) = 4 \left(\frac{4\sqrt{n}m_{\tilde{A}}}{\alpha_s\sqrt{n+2}m_{\text{pl}}} \right)^{\frac{n}{2}} \left(\frac{|\underline{\tilde{P}}|}{\tilde{R}(\tilde{\Gamma})} \right)^{\frac{n}{4}} \left(\frac{n-2}{n+2} |\underline{\tilde{P}}| \right)^{\frac{1}{2}} \quad (16.184)$$

where l is a big natural number. In this way we can travel in hyperspace (through extra dimension) almost for free. The energy to excite the warp factor is zero (or almost zero). Thus it could be created even from a fluctuation in our Universe.

Let us notice that Eq. (16.174) is in some sense universal for a warp factor. It means that it is the same for any realization of the Nonsymmetric Jordan-Thiry (Kaluza-Klein) Theory.

However, if we take Eq. (16.133) and if we demand

$$E_{\text{wf}} = 0, \quad (16.185)$$

we get some roots (they exist) depending on n , β and $|\gamma|$. Thus the roots, some values of $\tilde{A} > 0$ depend on details of the theory, they are not universal as in the case

of Eqs (16.172–174). In this case the Eq. (16.153) is in some sense a selfconsistency condition for a theory and more restrictive. The fluctuations of a tower of scalar fields in the case of $\sqrt{\tilde{A}} = \frac{1}{4}(2l+1)$ are as follows

$$\frac{1}{2}\psi_0^1 = \frac{(e^{\pi(2l+1)} - 1)}{2l+1} \quad (16.186)$$

$$\psi_m^1 = -\frac{m}{\pi} \cdot \frac{(e^{\pi(2l+1)} - 1)}{((2l+1)^2 + 4m^2)} \quad (16.187)$$

$$\psi_m^2 = \frac{(2l+1)(e^{\pi(2l+1)} - 1)}{\pi((2l+1)^2 + 4m^2)}. \quad (16.188)$$

In the case of very large l (which is really considered) one gets

$$\frac{1}{2}\psi_0^1 = \frac{e^{\pi(2l+1)}}{2l} \quad (16.186a)$$

$$\psi_m^1 = -\frac{m}{2\pi} \cdot \frac{e^{\pi(2l+1)}}{4(l^2 + m^2)} \quad (16.187a)$$

$$\psi_m^2 = \frac{l \cdot e^{\pi(2l+1)}}{2\pi(l^2 + m^2)} \quad (16.188a)$$

$$m = 1, 2, \dots$$

According to our calculations the energy of the fluctuations (excitations) (16.186a–188a) is zero.

Thus the only difficult problem to get a warp factor is to excite a tower of scalar fields coherently in a shape of (16.186–188a) or to wait for such a fluctuation. In the first case it is possible to use some techniques from quantum optics applied to the scalar fields ψ_m . In our case we get

$$\psi_0 = \frac{e^{\pi(2l+1)}}{2l} \quad (16.189)$$

$$\psi_m = \frac{e^{\pi(2l+1)}\sqrt{16l^2 + m^2}}{8\pi(l^2 + m^2)} \sin(m\lambda + \delta_m)e^{im_m t} \quad (16.190)$$

where

$$\text{tg } \delta_m = -\frac{m}{4l} \quad (16.191)$$

and m_m is given by the formula (16.122).

For ψ_0 is constant in time and does not contribute to the total energy of a warp factor (only to the energy of a warp factor equal to 1), we consider only

$$\psi(\lambda, t) = \frac{e^{\pi(2l+1)}}{8\pi} \left[\sum_{m=1}^{\infty} \frac{\sqrt{16l^2 + m^2}}{l^2 + m^2} \sin(m\lambda + \delta_m)e^{im_m t} + \frac{4\pi}{l} \right]. \quad (16.192)$$

For $t = 0$ we get

$$\psi(\lambda, 0) = e^{\pi(2l+1)\lambda} \quad (16.193)$$

and the wave packet (16.192) will disperse. Thus we should keep the wave packet to not decohere in such a way that

$$\psi(\lambda, t) = \frac{e^{\pi(2l+1)\lambda}}{8\pi} \left[\sum_{m=1}^{\infty} \frac{\sqrt{16l^2 + m^2}}{l^2 + m^2} \sin(m\lambda + \delta_m) + \frac{4\pi}{l} \right]. \quad (16.194)$$

It means $\psi(\lambda, t)$ should be a pure zero mode for all the time $t \geq 0$ (see Ref. [120]).

Let us come back to the Eq. (16.184) (earlier to Eq. (16.153)) taking under consideration that in a real model $m_{\tilde{A}} = \frac{\alpha_s}{r}$. One can rewrite Eq. (16.184) in the following way:

$$\left(\frac{2l+1}{4\alpha_s} \right)^{4/n} \left(\frac{n+2}{(n-2)|\tilde{P}|} \right)^{2/n} \left(\frac{\alpha_s \sqrt{n+2} m_{\text{pl}}}{4\sqrt{n} m_{\tilde{A}}} \right)^2 = \frac{|\tilde{P}|}{\tilde{R}(\tilde{\Gamma})} \quad (16.195)$$

or

$$\tilde{R}(\tilde{\Gamma}) = \left(\frac{4\alpha_s}{2l+1} \right)^{4/n} \left(\frac{(n-2)|\tilde{P}|}{n+2} \right)^{2/n} \left(\frac{4\sqrt{n} m_{\tilde{A}}}{\alpha_s \sqrt{n+2} m_{\text{pl}}} \right)^2 |\tilde{P}| = 2F(\zeta). \quad (16.196)$$

We want to find some conditions for μ and ζ (or ξ and ζ) in order to satisfy Eq. (16.153) for large l in a special model with $H = G2$, $\dim G2 = n = 14$ for $M = S^2$ and $\tilde{R}(\tilde{\Gamma})$ calculated for $G = SO(3)$. In this case one finds

$$F(\zeta) = 3^{2/7} \cdot 7 \cdot \left(\frac{\alpha_s}{l} \right)^{4/7} \left(\frac{m_{\tilde{A}}}{\alpha_s m_{\text{pl}}} \right)^2 |\tilde{P}|^{15/7} \quad (16.197)$$

where $\tilde{P}$ is given by the formula (8.64).

We calculate $\tilde{R}(\tilde{\Gamma})$ for a group $SO(3)$ in the fourth point of Ref. [18] (Eq. (7.21)) and we get

$$\tilde{R}(\tilde{\Gamma}) = \frac{2(2\mu^3 + 7\mu^2 + 5\mu + 20)}{(\mu^2 + 4)^2}. \quad (16.198)$$

The polynomial

$$W(\mu) = 2\mu^3 + 7\mu^2 + 5\mu + 20 \quad (16.199)$$

possesses only one real root

$$\mu_0 = -\frac{\sqrt[3]{1108 + 3\sqrt{135645}}}{6} - \frac{7}{6} - \frac{19}{6 \cdot \sqrt[3]{1108 + 3\sqrt{135645}}} = -3.581552661 \dots \quad (16.200)$$

It is interesting to notice that $W(-3.581552661) = 2.5 \cdot 10^{-9}$ and for 70-digit approximation of μ_0 , $\tilde{\mu}$ equal to

$$-3.581552661076733712599740215045436907383569800816123632201827285932446,$$

we have

$$W(\tilde{\mu}) = 0.1 \cdot 10^{-67}.$$

The interesting point in $\tilde{\mu}$ is that any truncation of $\tilde{\mu}$ (it means, removing some digits from the end of this number) results in growing W , i.e.,

$$0 < W(\tilde{\mu}_n) < W(\tilde{\mu}_{n-1}),$$

where $\tilde{\mu}_n$ means a truncation of $\tilde{\mu}$ with only n digits, $n \leq 70$, $\tilde{\mu}_{70} = \tilde{\mu}$. This can help us in some approximation for $\tilde{R}(\tilde{\Gamma}) > 0$, close to zero.

Substituting (16.198) into Eq. (16.196) we arrive to the following equation:

$$-F(\zeta)\mu^4 + 2\mu^3 + \mu^2(7 - 4F(\zeta)) + 5\mu + 4(-2F(\zeta) + 5) = 0. \quad (16.201)$$

Considering $F(\zeta)$ small we can write a root of this polynomial as

$$\mu = \mu_0 + \varepsilon \quad (16.202)$$

where μ_0 is the root of the polynomial (16.199) and ε is a small correction. In this way one gets the solution

$$\mu = -3.581552661 \dots + 47 \left(\frac{\alpha_s}{l} \right)^{4/7} \left(\frac{m_{\tilde{A}}}{m_{\text{pl}}} \right)^2 q(\zeta) \quad (16.203)$$

where $q(\zeta)$ is given by the formula

$$\begin{aligned} q(\zeta) = & \left| \frac{16|\zeta|^3(\zeta^2 + 1)}{3(2\zeta^2 + 1)(1 + \zeta^2)^{5/2}} \left(\zeta^2 E \left(\frac{|\zeta|}{\sqrt{\zeta^2 + 1}} \right) - (2\zeta^2 + 1) K \left(\frac{|\zeta|}{\sqrt{\zeta^2 + 1}} \right) \right) \right. \\ & \left. + 8 \ln \left(|\zeta| \sqrt{\zeta^2 + 1} \right) + \frac{4(1 + 9\zeta^2 - 8\zeta^4)|\zeta|^3}{3(1 + \zeta^2)^{3/2}} \right|^{\frac{15}{7}} \\ & \times \left[\ln \left(|\zeta| + \sqrt{\zeta^2 + 1} \right) + 2\zeta^2 + 1 \right]^{-\frac{15}{7}}, \end{aligned} \quad (16.204)$$

$|\zeta| > |\zeta_0| = 1.36$ (see Fig. 6), and K and E are given by the formulae (8.65a) and (8.65b).

In this way taking sufficiently large l we can choose μ such that $\tilde{R}(\tilde{\Gamma}) > 0$ and quite arbitrary ζ ($\tilde{P} < 0$) to satisfy a zero energy condition for an excitation of a warp factor.

Taking $m_{\tilde{A}} = m_{\text{EW}} \simeq 80 \text{ GeV}$, $m_{\text{pl}} \simeq 2.4 \cdot 10^{18} \text{ GeV}$, $\alpha_s = \sqrt{\alpha_{\text{em}}} = \frac{1}{\sqrt{137}}$, one gets from Eq. (16.203)

$$\mu = -3.581552661 \dots + 4.1 \cdot 10^{-35} \frac{q(\zeta)}{l^{4/7}} \quad (16.205)$$

which justifies our approximation for ε and simultaneously gives an account for a smallness of a cosmological constant $\tilde{R}(\tilde{\Gamma}(\mu)) \simeq 0$. Thus it seems that in this simple toy model we can achieve a large c_{eff} without considering large cosmological constant.

Let us give some numerical estimation for a time needed to travel a distance of 200 Mps

$$t_{\text{travel}} = \frac{L}{c_{\text{eff}}^{\text{max}}} = \frac{L}{c} \cdot e^{-(2l+1)\pi/4}.$$

One gets for $L = 200 \text{ Mps}$, $t_{\text{travel}} = 2 \cdot 10^{13} \text{ s} \cdot e^{-l\pi/2}$ and for $l = 100$, $t_{\text{travel}} \cong 12 \text{ ns}$.

Let us notice that for $l = 100$, μ from Eq. (16.205) is equal to

$$\mu = -3.581552661 \dots + 5.94 \cdot 10^{-44} q(\zeta). \quad (16.206)$$

$q(\zeta)$ cannot be too large (it is a part of cosmological term $q(\zeta) = |\tilde{P}|^{15/7}$). If $\tilde{P}$ is of order 0.1, q is of order 10^{-16} ; if $\tilde{P}$ is of order 1.1, q is of order 0.6. Thus μ is very close to μ_0 for some reasonable values of $|\tilde{P}|$. It is interesting to ask how to develop this model to more dimensional case. It means, to the manifold M which is not a circle. It is easy to see that in this case we should consider a warp factor which depends on more coordinates taking under consideration some ansatzes. For example we can consider a warp factor in a shape

$$\exp\left(\sum_{i=1}^m \sqrt{\tilde{A}_i} f_i(y)\right) \quad (16.207)$$

where $f_i(y) = x_i$, $i = 1, 2, \dots, m = \frac{1}{2}n_1(n_1 + 1)$ are parametric equations of the manifold M in m -dimensional euclidean space and

$$ds_M^2 = \left(\sum_{k=1}^m \left(\frac{\partial f^k}{\partial y^i}\right) \left(\frac{\partial f^k}{\partial y^j}\right)\right) dy^i \otimes dy^j \quad (16.207a)$$

is a line element of the manifold M . In the case of the manifold S^2 it could be

$$\begin{aligned} f_1 &= \cos \theta \sin \lambda \\ f_2 &= \cos \theta \cos \lambda \\ f_3 &= \sin \theta \\ m &= 3 \end{aligned} \quad (16.208)$$

and a warp factor takes the form

$$\begin{aligned} &\exp\left(\sqrt{\tilde{A}_1} \cos \theta \sin \lambda + \sqrt{\tilde{A}_2} \cos \theta \cos \lambda + \sqrt{\tilde{A}_3} \sin \theta\right), \\ &\theta \in \langle 0, \pi \rangle, \lambda \in \langle 0, 2\pi \rangle. \end{aligned} \quad (16.209)$$

Afterwards we should develop a warp factor in a series of spherical harmonics on the manifold M and calculate an energy of a tower of scalar particles connected with this development. We should of course regularize the series (divergent) using ζ -functions techniques. In this case we should use ζ -functions connected with the Beltrami-Laplace operator on the manifold M . Afterwards we should find a zero energy condition for special types of constants $\tilde{A}_i$, $i = 1, 2, \dots, m$. We can of course calculate the energy of an excitation of the warp factor using some formulae from classical field theory. In

the case of $M = S^2$ we have to do with an ordinary Laplace operator on S^2 and with ordinary spherical harmonics with spectrum given by Eq. (16.14). The Fourier analysis of (16.209) can be proceeded on S^2 .

Moreover a toy model with S^1 can be extended to the cosmological background in such a way that

$$ds_5^2 = e^{\sqrt{\tilde{A}}\lambda} dt^2 - R^2(t)(dx^2 + dy^2 + dz^2) - r^2 d\lambda^2. \quad (16.210)$$

It seems that from practical (calculational) point of view it is easier to consider a metric of the form

$$ds_5^2 = dt^2 - e^{-\sqrt{\tilde{A}}\lambda} R^2(t)(dx^2 + dy^2 + dz^2) - r^2 d\lambda^2 \quad (16.211)$$

where $R(t)$ is a scale factor of the Universe. For further investigations we should also consider more general metrics, i.e.

$$ds_6^2 = dt^2 - \exp\left(-\sqrt{\tilde{A}}(\cos\theta(\sin\lambda + \cos\lambda) + \sin\theta)\right) \times R^2(t)(dx^2 + dy^2 + dz^2) - r^2(d\theta^2 + \sin^2\theta d\lambda^2) \quad (16.212)$$

or even

$$ds_{4+n_1}^2 = dt^2 - \exp\left(-\sum_{i=1}^m \sqrt{\tilde{A}_1} f_i(y)\right) \cdot R^2(t) \frac{dx^2 + dy^2 + dz^2}{1 - k(x^2 + y^2 + z^2)} - r^2 \left(\left(\sum_{k=1}^m \left(\frac{\partial f^k}{\partial y^i} \right) \left(\frac{\partial f^k}{\partial y^j} \right) \right) \cdot dy^i \otimes dy^j \right) \quad (16.213)$$

where

$$m = \frac{1}{2} n_1(n_1 + 1). \quad (16.214)$$

Let us notice that our toy model with a travel in a hyperspace (this is really a travel along dimensions of a vacuum state manifold) can go to some “acausal properties”. This is evident if we consider two signals: one sent in an ordinary way with a speed of light and a second via the fifth dimension to the same point. The second signal will appear earlier than the first one at the point. Thus it will affect this point earlier. Effectively it means a superluminal propagation of signals in a Minkowski space (or in Friedmann-Robertson-Walker Universe). In this way we can construct a time-machine using this solution (even if it does not introduce tachyons in a 4-dimensional space-time), i.e. a space-time where there are closed time-like (or null) curves.

Let us notice that given a superluminal signal we can always find a reference frame in which it is travelling backwards in time. Suppose we send a signal from A to B (at time $t = 0$) at an effective speed $u_{\text{eff}} > 1$ in frame S_1 (it means we send a signal through a path described here earlier and an effective speed $u_{\text{eff}} > 1$) with coordinates (t, x) (we consider only one space dimension). In a frame S_2 moving with respect to S_1 with

velocity $v > \frac{1}{u_{\text{eff}}}$, the signal travels backwards in t' time. This follows from the Lorentz transformation, i.e.

$$t'_B = \frac{t_B(1 - vu_{\text{eff}})}{\sqrt{1 - v^2}} \quad (16.215)$$

$$x'_B = \frac{t_B(1 - \frac{v}{u_{\text{eff}}})}{\sqrt{1 - v^2}}. \quad (16.216)$$

We require both $x'_B > 0$, $t'_B < 0$, that is $\frac{1}{u_{\text{eff}}} < v < u_{\text{eff}}$ which is possible only if $u_{\text{eff}} > 1$.

However this by itself does not mean a violation of a causality. For that we require that the signal can be returned from B to a point in the past of light cone of A . Moreover, if we return the signal from B to C with the same effective speed (it means we send a signal through the fifth dimension with the same effective speed using the warp factor) in frame S_1 , then it arrives at C in the future cone of B . The situation is physically equivalent in the Lorentz boosted frame S_2 —the return signal travels forward in time t' and arrives at C in the future cone of A . This is a frame-independent statement. If a backwards-in-time signal AC is possible in frame S_1 , then a return signal sent with the same speed u_{eff} will arrive at a point D in the past light cone of A creating a closed time-like curve $ACDA$. In a Minkowski space (a space for $\lambda = 0$, or in general for $\lambda = \lambda_0 = \text{const.}$), local Lorentz invariance results in that if a superluminal signal such as AB is possible, then so is one of AC for it is just given by an appropriate Lorentz boost. The existence of a global inertial frames in a Minkowski space guarantees the existence of the return signal CD .

Thus we get unacceptable closed time loops. Moreover the theory for the five-dimensional world is still causal, as in five-dimensional General Relativity. Let us notice that in a cosmological background described by a metric (16.211) the situation will be very similar. We have here also global inertial frame for $\lambda = 0$, and similar possibility to send a superluminal signal through fifth dimension. However, in this case we should change a shape of Lorentz transformation to include a scale factor $R(t)$ which slightly complicates considerations. Moreover in order to create a warp factor we need a zero energy condition. If this condition is not satisfied we need an infinite amount of an energy to create it. The zero energy condition imposes some restriction on parameters which are involved in our unified theory. Maybe it is impossible to satisfy these conditions in our world. In this way it would be an example of Hawking's chronology protection conjecture: "*the laws of physics do not allow the appearance of closed timelike curves*". However, if a zero energy condition is satisfied in a realistic Nonsymmetric Kaluza–Klein (Jordan–Thiry) Theory (it means in such a theory which is consistent with a phenomenology of a contemporary elementary particle physics) then we meet possibility of effective closed time-like loops (even the higher dimensions are involved). In this case we should look for some weaker conjectures (especially in an expanding Universe, not just in a Minkowski space). Such weaker condition protects us from some different causal paradoxes as the following paradox: a highly energetic

signal can destroy a laboratory before it was created. In some sense it means some kind of selfconsistency conditions.

Thus what is a prescription to construct a time-machine in our theory? It is enough to have a 5-dimensional (or $(n_1 + 4)$ -dimensional) space-time with a warp factor described here earlier. Afterwards we need two frames S_1 and S_2 in relative motion with relative velocity $v > \frac{1}{u_{\text{eff}}}$ where u_{eff} is an effective superluminal velocity $u_{\text{eff}} > 1$. This could be arranged e.g. as two spaceships with relative motion (A and C). Sending a signal with effective velocity u_{eff} from A to C we get the signal in a past of C . Afterwards we send a signal (through the fifth dimension) from C to A . In this way we arrive in a past of A .

If we want to travel into a future we should reverse a procedure with

$$-1 < v < 0.$$

In both cases we use an effective time-like loop, constructed due to a warp factor in a five-dimensional space-time. Using this loop several times we can go to the future or to the past as far as we want. (As in a TVP spectacle *Through the Fifth Dimension*.) An interesting point which can be raised consists in an effectiveness of such a travel. It means how much time we need to get e.g. 1 s to past or 1 s to future. It depends on a relative velocity v and u_{eff} . The effectiveness can be easily calculated.

$$\eta = \frac{u_{\text{eff}} v}{\sqrt{1 - v^2}}. \quad (16.217)$$

If we use our estimation for c_{eff} (see Eq. (16.182)), we get

$$\eta = \frac{v}{\sqrt{1 - v^2}} e^{(2l+1)\pi/4} \quad (16.218)$$

for l large. It is easy to see that $\eta \rightarrow \infty$ if $v \rightarrow 1$ and if $l \rightarrow \infty$.

Let us take $v = \frac{1}{10}$ and $l = 100$. In this case one gets

$$\eta \approx 10^{67}. \quad (16.219)$$

What does this number mean? It simply means that in order to go 1 s into the past or into the future we loose 10^{-67} s of our life.

However, to be honest, we should use as u_{eff} a velocity different from c_{eff} . In this case a time needed to travel through the fifth dimension really matters. Let a speed along the fifth dimension be equal to the velocity of light (or close to it). In this case one gets

$$u_{\text{eff}} = \frac{c_{\text{eff}}}{1 + 2(\frac{r_0}{L})c_{\text{eff}}} \quad (16.220)$$

where $r_0 \simeq 10^{-16}$ cm, L is the distance between A and B , e.g. 1000 km. In this case u_{eff} is smaller,

$$u_{\text{eff}} = 10^{26} \quad (16.221)$$

and

$$\eta \simeq 10^{25}. \quad (16.222)$$

In this case we can estimate how much time we loose to go 2000 years back or forward. It is $6.3 \cdot 10^{-15}$ s. However, if we want to see dinos we should go back 100 mln yr and we loose 10^{-10} s. Of course in the formula for u_{eff} we can use smaller L and a velocity in the place c_{eff} can be smaller too. We can travel more comfortably changing smoothly a velocity. Any time a time to loose to travel back or forward in time for our time machine is very small. In some sense it is very effective (if it is possible to construct).

Let us notice that the warp factor (or $\tilde{A}$) is fixed by the constants of the theory (especially by a cosmological constant λ_{c0}). In this way only in some special cases we get a zero energy condition. However, in the case of the warp factor existence we have to do with a tower of scalar fields (not only with a quintessence). The quintessence field is a zero mode of this tower. Thus in the case of an existing of a tower of scalar fields we will have to do with a corrected cosmological constant, which is really an averaged cosmological term over S^1 (in general over M)

$$\lambda_{c0}^{\text{corrected}} = \frac{1}{2\pi} \int_0^{2\pi} (|\gamma| - \beta e^{2\Psi(\lambda)}) e^{n\Psi(\lambda)} d\lambda \quad (16.223)$$

where $\Psi(\lambda)$ is given by Eq. (16.118). In this way according to our considerations concerning an energy of excitation

$$\lambda_{c0}^{\text{corrected}} = E_{\text{wf}}. \quad (16.224)$$

However, in the formula for E_{wf} we dropped an energy for a zero mode for a tower of scalar fields (i.e., for a quintessence). Thus

$$\lambda_{c0}^{\text{tot}} = \lambda_{c0}(\Psi_0) + \lambda_{c0}^{\text{corrected}}. \quad (16.225)$$

If the zero energy condition for a warp factor is satisfied, we get

$$\lambda_{c0}^{\text{tot}} = \lambda_{c0} \quad (16.226)$$

and we do not need any tuning of parameters in the theory. However, now we should write down full 5-dimensional equations in our theory ($n_1 + 4$ in general) to look for more general solutions mentioned here earlier.

In Fig. 8 we draw a time-like loop in our theory together with a realistic path in five-dimensional space-time with a warp factor corresponding to the loop.

Let us consider a $(n_1 + 4)$ formalism of the lagrangian in our theory given by Eq. (7.5). In this case we do not suppose that a scalar field does not depend on a point of a manifold M . In general $\Psi = \Psi(x, y)$, $x \in E$, $y \in M$. Simultaneously we should suppose that $g_{\mu\nu} = g_{\mu\nu}(x, y)$ for we use the transformation (7.4). From the variation principle

$$\delta \int_U d^4x d^{n_1}y L(g, \overline{W}, \tilde{A}, \Psi, \Phi) \sqrt{-g} \sqrt{\tilde{g}} = 0, \quad U \subset E \times M \quad (16.227)$$




Fig. 8. A closed time-like curve (a time-like loop) $ACDA$ in a Minkowski space with a warp factor. $AA'C'C''D'D''A''A$ means a realistic path in a 5-dimensional space-time with a warp factor.

we get equations (9.3÷9.18a) with some differences connected with the fact that now we do not average over a manifold M . It means that in definitions for $V(\Phi)$ there are no integrals and the same is true for $\tilde{P}$. Now in the place of $\tilde{P}$ we have simply $\hat{\tilde{R}}(\hat{\tilde{T}})$ (i.e. a scalar curvature for a metric $g_{\tilde{a}\tilde{b}}$). In the place of $\tilde{g}^{[\tilde{a}\tilde{b}]}$ we have simply $g^{[\tilde{a}\tilde{b}]}$ and so on. Moreover the only one “material source” which lives in $(n_1 + 4)$ -dimensional space-time is a scalar field Ψ . For this reason in a lagrangian of this field we have the formula (16.4) (or (16.3)). In this way a full set of field equations should be supplemented with

$$R_{(\mu\tilde{a})} = \frac{8\pi K}{c^4} T_{(\mu\tilde{a})}^{\text{scal}}(\Psi) \quad (16.228)$$

$$R_{(\tilde{a}\tilde{b})} = \frac{8\pi K}{c^4} T_{(\tilde{a}\tilde{b})}^{\text{scal}}(\Psi) \quad (16.229)$$

and

$$g_{\mu\nu,\tilde{s}} - \Gamma_{\mu\tilde{s}}^{\xi} g_{\xi\nu} - \Gamma_{\tilde{s}\nu}^{\xi} g_{\mu\xi} = 0 \quad (16.230)$$

where $T_{(\mu\tilde{a})}^{\text{scal}}$, $T_{(\tilde{a}\tilde{b})}^{\text{scal}}$ are remaining parts of an energy-momentum tensor for a scalar field Ψ and $\Gamma_{\mu\tilde{s}}^{\xi}$, $\Gamma_{\tilde{s}\nu}^{\xi}$ are remaining coefficients of a connection. $R_{\mu\nu}$ are constructed from $\overline{W}^{\alpha}_{\beta}$ connection and $R_{(\mu\tilde{a})}$, $R_{(\tilde{a}\tilde{b})}$ are constructed from the connection induced

by relations (9.5), (6.25), (9.4) and the condition

$$\Gamma_{\xi\tilde{s}}^\xi = \Gamma_{\tilde{s}\xi}^\xi. \quad (16.231)$$

A metric tensor has a general form

$$ds_{n_1+4}^2 = g_{(\mu\nu)}(x, y) dx^\mu \otimes dx^\nu - r^2 g_{(\tilde{a}\tilde{b})} \theta^{\tilde{a}} \otimes \theta^{\tilde{b}}. \quad (16.232)$$

We have the formula

$$\begin{aligned} T_{AB}^{\text{scal}}(\Psi) = & -\frac{e^{(n+2)\Psi}}{16\pi} \left\{ (g_{KA}g_{ZB} + g_{ZA}g_{KB}) \right. \\ & \times \tilde{g}^{(CK)} \tilde{g}^{(NZ)} \left[\frac{n^2}{2} (g^{SM}g_{NS} - \delta^M_N) + \overline{M}\Psi_{,N} \right] \Psi_{,C} \\ & \left. - g_{AB} \left[\overline{M}\tilde{g}^{(NM)} + n^2 g^{[MN]} g_{[DM]} \tilde{g}^{(CD)} \Psi_{,N} \Psi_{,C} \right] \right\} \end{aligned} \quad (16.233)$$

g_{AB} tensor is here in the place of γ_{AB} in order to stress that g_{AB} depends on x and y . We suppose however that g_{AB} has the following structure

$$g_{AB} = \left(\begin{array}{c|c} g_{\mu\nu} & 0 \\ \hline 0 & r^2 g_{\tilde{a}\tilde{b}} \end{array} \right). \quad (16.234)$$

This tensor is of course nonsymmetric, $g = \det(g_{AB})$.

These field equations are fundamental equations to find our solutions. For we are looking for some cosmological solutions (with warp factor), we suppose additionally that

$$g_{\mu\nu} = g_{(\mu\nu)}(x, y) \quad (16.235)$$

$$g_{\tilde{a}\tilde{b}} = g_{\tilde{a}\tilde{b}}(y) \quad (16.235a)$$

The metric on a manifold M is still nonsymmetric.

We will also suppose that the total energy momentum tensor (except for a scalar field) is given by a sum of several hydrodynamics energy-momentum tensors including a dust and a radiation. We get five-dimensional field equations for $M = S^1$ and 6-dimensional for $M = S^2$ (a 6-dimensional Weinberg–Salam model). Moreover, there is a different approach to derive $(n_1 + 4)$ -dimensional equations in the Nonsymmetric Kaluza–Klein (Jordan–Thiry) Theory.

Let us consider an $(n_1 + 4)$ generalization of 4-dimensional field equations in the Nonsymmetric Kaluza–Klein (Jordan–Thiry) Theory. According to the second point

of Ref. [18] we get (see Eqs (4.8.3)÷(4.8.14)):

$$\bar{R}_{MN}(\bar{W}) - \frac{1}{2} g_{MN} \bar{R}(\bar{W}) = 8\pi K \left(T_{MN}^{\text{gauge}} + T_{MN}^{\text{scal}} + g_{MN} F \right) \quad (16.236)$$

$$\tilde{g}^{[MN]}_{,N} = 0 \quad (16.237)$$

$$g_{MN,S} - g_{ZN} \bar{\Gamma}^Z_{MS} - g_{MZ} \bar{\Gamma}^Z_{SN} = 0 \quad (16.238)$$

$$\begin{aligned} \nabla_M^{\text{gauge}} \left(l_{ab} \tilde{L}^{aAM} \right) &= 2 \tilde{g}^{[AB]} \nabla_B^{\text{gauge}} \left(h_{ab} \tilde{g}^{[MN]} H_{MN}^a \right) \\ &+ (n+2) \partial_B \Psi \left[l_{ab} L^{aBA} - 2g^{[BA]} \left(h_{ab} \tilde{g}^{[MN]} H_{MN}^a \right) \right] \end{aligned} \quad (16.239)$$

$$\begin{aligned} \left[(n^2 + 2\bar{M}) \tilde{g}^{(AM)} - n^2 g^{NM} \tilde{g}^{(AS)} \right] \frac{\partial^2 \Psi}{\partial x^A \partial x^M} + \frac{1}{\sqrt{-g}} \partial_M \left\{ \sqrt{-g} \left(n^2 \tilde{g}^{(MA)} \right. \right. \\ \left. \left. - \frac{n^2}{2} g_{DN} \left(g^{NA} \tilde{g}^{(MD)} + g^{NM} \tilde{g}^{(AD)} \right) - 2\bar{M} \tilde{g}^{(MA)} \right) \right\} \frac{\partial \Psi}{\partial x^A} \\ - 8\pi(n+2)\pi e^{-(n+2)\Psi} (\mathcal{L}_{YM} - 2F) = 0 \end{aligned} \quad (16.240)$$

where

$$\begin{aligned} T_{AB}^{\text{gauge}} &= -\frac{l_{ab}}{4\pi} \left\{ g_{CB} g^{TZ} g^{EC} L_{ZA}^a L_{TE}^b - 2g^{[MN]} H_{MN}^{(a} H_{AB}^{b)} \right. \\ &\left. - \frac{1}{4} g_{AB} \left[L^{aMN} H_{MN}^b - 2 \left(g^{[MN]} H_{MN}^a \right) \left(g^{[CS]} H_{[CS]}^b \right) \right] \right\} \end{aligned} \quad (16.241)$$

is the energy momentum tensor for the $(n_1 + 4)$ -dimensional gauge (Yang-Mills) field, $T_{AB}^{\text{scal}}(\Psi)$ is given by the Eq. (16.233)

$$K = G_{(n_1+4)} e^{-(n+2)\Psi} \quad (16.242)$$

$$F = e^{2(n+2)\Psi} \frac{\tilde{R}(\tilde{\Gamma})}{16\pi} \quad (16.243)$$

$$\tilde{L}^{aMN} = \sqrt{-g} g^{BM} g^{CA} L_{BC}^a \quad (16.244)$$

$$\tilde{g}^{[MN]} = \sqrt{-g} g^{[MN]} \quad (16.245)$$

where

$$G_{(n_1+4)} = \frac{1}{M_{\text{pl}(n_1+4)}}$$

is an $(n_1 + 4)$ -dimensional gravitational constant, ∇_N^{gauge} means a gauge derivative for a connection defined on P (over an $(n_1 + 4)$ -dimensional base),

$$l_{dc} g_{MB} g^{CM} L_{CA}^d + l_{cd} g_{AM} g^{MC} L_{BC}^d = 2l_{cd} g_{AM} g^{MC} H_{BC}^d \quad (16.246)$$

$$\mathcal{L}_{YM} = \frac{1}{8\pi} l_{cd} (H^c H^d - L^{cMN} H_{MN}^d) \quad (16.247)$$

$$H^c = g^{[AB]} H_{AB}^c \quad (16.248)$$

H_{AB}^c is a curvature of a connection ω on a fibre bundle P (over an $(n_1 + 4)$ -dimensional base).

Now let us suppose that a base manifold is a Cartesian product $E \times M$, $M = G/G_0$ (as in sec. 4) and let us suppose a symmetry conditions on a connection ω over $E \times M$ ($\dim M = n_1$), i.e. an invariancy of ω with respect to the action of the group G on $E \times G/G_0$ (see sec. 4 for details). Simultaneously we suppose that g_{AB} has a form given by Eq. (16.234), where $g_{\tilde{a}\tilde{b}}$ is defined and examined in sec. 4. In this way we can reduce the field equation to simpler form described above.

The connection $\overline{W}^A_B$ is given by the equation

$$\overline{W}^A_B = \overline{\omega}^A_B - \frac{2}{(n_1 + 3)} \overline{W} \delta^A_B \quad (16.249)$$

where

$$\overline{\omega}^A_B = \overline{\Gamma}^A_{BC} \theta^C \quad (16.250)$$

$$\overline{W} = \overline{W}_C \theta^C = \frac{1}{2} \left(\overline{W}^S_{CS} - \overline{W}^S_{SC} \right) \theta^C \quad (16.251)$$

The dimension reduction procedure (described in sec. 4) reduces $(n_1 + 4)$ -dimensional Yang-Mills' field to a 4-dimensional Higgs' field. R_{AB} is a Ricci (Moffat–Ricci) tensor for the connection $\overline{W}^A_B$. In our approximation we can consider those fields as 4-dimensional matter fields (i.e. hydrodynamics energy-momentum tensors). Remembering that only $g_{\mu\nu} = g_{\nu\mu}(x, y)$ and $\Psi = \Psi(x, y)$ are functions depending on $y \in M$ we can try to solve our equations for metrics defined earlier. Eventually let us remark that both approaches described here are not strictly equivalent. Moreover for the class of problems considered here they are equivalent.

17. The Approximation Procedure for the Lagrangian of the Higgs Field in the Nonsymmetric Jordan–Thiry Theory

In this section we develop the approximation procedure for the lagrangian involving Higgs' field up to the second order of expansion in ζ and $h_{\mu\nu} = g_{\mu\nu} - \eta_{\mu\nu}$. We also deal with geodetic equations on the Jordan-Thiry manifold in the first order of approximation with respect to ζ and $h_{\mu\nu}$ for terms coupled to Higgs' field. We find also the field equations in linear approximation and an influence of $\overline{W}_\mu$ and electromagnetic fields in this approximation. In this section we use p_{ab} to denote the Killing–Cartan tensor.

Let us consider the lagrangian for the Higgs' field i.e.:

$$\mathcal{L}(\Phi, \Psi) = \frac{2}{r^2} e^{-2\Psi} \mathcal{L}_{\text{kin}}(\overset{\text{gauge}}{\nabla} \Phi) - \frac{1}{r^4} e^{(n-2)\Psi} V(\Phi) - \frac{4}{r^4} e^{(n-2)\Psi} \mathcal{L}_{\text{int}}(\Phi, \tilde{A}) \quad (17.1)$$

where

$$\mathcal{L}_{\text{kin}}(\overset{\text{gauge}}{\nabla} \Phi) = \ell_{ab} (g^{\tilde{b}\tilde{n}} g^{\alpha\mu} L^a_{\tilde{\alpha}} \overset{\text{gauge}}{\nabla}_\mu \Phi^b_{\tilde{n}})_{av} \quad (17.2a)$$

$$V(\Phi) = 2\ell_{ab} (\underline{g}^{([\tilde{m}\tilde{n}][\tilde{a}\tilde{b}]}) H^a_{\tilde{m}\tilde{n}} H^b_{\tilde{a}\tilde{b}} - \underline{g}^{\tilde{a}\tilde{m}\tilde{b}\tilde{n}} L^a_{\tilde{a}\tilde{b}} H^b_{\tilde{m}\tilde{n}})_{av} \quad (17.2b)$$

$$\mathcal{L}_{\text{int}}(\Phi, \tilde{A}) = p_{ab} \mu^a_i \tilde{H}^i \underline{g}^{[\tilde{a}\tilde{b}]} H^b_{\tilde{a}\tilde{b}} \quad (17.2c)$$

$\underline{g}^{([\tilde{m}\tilde{n}][\tilde{a}\tilde{b}]})$, $\underline{g}^{\tilde{a}\tilde{m}\tilde{b}\tilde{n}}$ and $\underline{g}^{[\tilde{a}\tilde{b}]}$ are given by the formula (6.24) and $H^b_{\tilde{a}\tilde{b}}$ by the formula (4.55).

Moreover for $L^a_{\tilde{\alpha}\tilde{b}}$ and $L^a_{\tilde{a}\tilde{b}}$ we have

$$\ell_{dc} g_{\beta\mu} g^{\gamma\beta} L^d_{\gamma\tilde{a}} + \ell_{cd} g_{\tilde{a}\tilde{m}} g^{\tilde{m}\tilde{c}} L^d_{\mu\tilde{c}} = 2\ell_{cd} g_{\tilde{a}\tilde{m}} g^{\tilde{m}\tilde{c}} \overset{\text{gauge}}{\nabla}_\mu \Phi^d_{\tilde{c}} \quad (17.3a)$$

$$\ell_{dc} g_{\tilde{m}\tilde{b}} g^{\tilde{c}\tilde{m}} L^d_{\tilde{c}\tilde{a}} + \ell_{cd} g_{\tilde{a}\tilde{m}} g^{\tilde{m}\tilde{c}} L^d_{\tilde{b}\tilde{c}} = 2\ell_{cd} g_{\tilde{a}\tilde{m}} g^{\tilde{m}\tilde{c}} H^d_{\tilde{b}\tilde{c}} \quad (17.3b)$$

Let us consider an expansion for $g_{\tilde{a}\tilde{b}}$ and $g^{\tilde{a}\tilde{b}}$

$$\begin{aligned} g_{\tilde{a}\tilde{b}} &= h^0_{\tilde{a}\tilde{b}} + \zeta k^0_{\tilde{a}\tilde{b}} \\ g^{\tilde{m}\tilde{a}} g_{\tilde{n}\tilde{a}} &= (h^{0\tilde{m}\tilde{a}} + h^{1\tilde{m}\tilde{a}} + h^{2\tilde{m}\tilde{a}} + \dots) \cdot (h^0_{\tilde{n}\tilde{a}} + \zeta k^0_{\tilde{n}\tilde{a}}) = \delta^{\tilde{m}}_{\tilde{n}} \end{aligned} \quad (17.4)$$

From (17.4) we get:

$$h^{1\tilde{m}\tilde{n}} = -\zeta h^{0\tilde{m}\tilde{a}} h^{0\tilde{n}\tilde{b}} k^0_{\tilde{b}\tilde{a}} \quad (17.5)$$

$$h^{2\tilde{m}\tilde{n}} = -\zeta h^{0\tilde{n}\tilde{b}} h^{1\tilde{m}\tilde{a}} k^0_{\tilde{b}\tilde{a}} = \zeta^2 h^{0\tilde{n}\tilde{b}} h^{0\tilde{m}\tilde{e}} h^{0\tilde{a}\tilde{s}} k^0_{\tilde{s}\tilde{e}} k^0_{\tilde{b}\tilde{a}} \quad (17.6)$$

and

$$g^{\tilde{m}\tilde{n}} = h^{0\tilde{m}\tilde{n}} - \zeta h^{0\tilde{m}\tilde{a}} h^{0\tilde{n}\tilde{b}} k_{\tilde{b}\tilde{a}}^0 + \zeta^2 h^{0\tilde{n}\tilde{b}} h^{0\tilde{m}\tilde{e}} h^{0\tilde{a}\tilde{s}} k_{\tilde{s}\tilde{e}}^0 k_{\tilde{b}\tilde{a}}^0 + \dots \quad (17.7)$$

Let us write the Eq. (17.3b) in a more convenient form:

$$\begin{aligned} \ell_{dc}(\delta_{\tilde{b}}^{\tilde{c}} - 2\zeta k_{\tilde{b}\tilde{d}}^0 \tilde{g}^{\tilde{c}\tilde{d}}) L_{\tilde{c}\tilde{a}}^d + \ell_{cd}(\delta_{\tilde{a}}^{\tilde{c}} - 2\zeta k_{\tilde{d}\tilde{a}}^0 \tilde{g}^{\tilde{d}\tilde{c}}) L_{\tilde{b}\tilde{c}}^d = \\ = 2(\delta_{\tilde{a}}^{\tilde{c}} - 2\zeta k_{\tilde{d}\tilde{a}}^0 \tilde{g}^{\tilde{d}\tilde{c}}) H_{\tilde{b}\tilde{c}}^d \ell_{cd}. \end{aligned} \quad (17.8)$$

Let us expand $L_{\tilde{b}\tilde{c}}^d$ into a power series with respect to ζ :

$$L_{\tilde{b}\tilde{c}}^d = L_{\tilde{b}\tilde{c}}^{(0)d} + L_{\tilde{b}\tilde{c}}^{(1)d} + L_{\tilde{b}\tilde{c}}^{(2)d} + \dots \quad (17.9)$$

Using (17.5) and (17.6) one gets up to the second order:

$$\begin{aligned} \ell_{dc} L_{\tilde{b}\tilde{a}}^d + \ell_{cd} L_{\tilde{b}\tilde{a}}^d = 2\ell_{cd} H_{\tilde{b}\tilde{a}}^d - 4\zeta k_{\tilde{d}\tilde{a}}^0 \ell_{cd} H_{\tilde{b}\tilde{c}}^d \cdot \\ \cdot (h^{0\tilde{d}\tilde{c}} - \zeta h^{0\tilde{d}\tilde{t}} h^{0\tilde{t}\tilde{s}} k_{\tilde{s}\tilde{t}}^0 + \zeta^2 h^{0\tilde{d}\tilde{e}} h^{0\tilde{c}\tilde{r}} h^{0\tilde{s}\tilde{t}} k_{\tilde{s}\tilde{e}}^0 k_{\tilde{r}\tilde{t}}^0) + \\ + 2\zeta (h^{0\tilde{c}\tilde{d}} - \zeta h^{0\tilde{c}\tilde{s}} h^{0\tilde{t}\tilde{d}} k_{\tilde{t}\tilde{s}}^0 + \zeta^2 h^{0\tilde{c}\tilde{r}} h^{0\tilde{d}\tilde{e}} h^{0\tilde{s}\tilde{t}} k_{\tilde{s}\tilde{r}}^0 k_{\tilde{e}\tilde{t}}^0) \cdot \\ \cdot (k_{\tilde{b}\tilde{d}}^0 \ell_{cd} L_{\tilde{c}\tilde{a}}^d + k_{\tilde{c}\tilde{a}}^0 \ell_{cd} L_{\tilde{b}\tilde{d}}^d) \end{aligned} \quad (17.10)$$

One easily finds:

$$L_{\tilde{b}\tilde{a}}^{(0)a} = p^{ab} \ell_{bc} H_{\tilde{b}\tilde{a}}^c = H_{\tilde{b}\tilde{a}}^a + \xi p^{ab} k_{bc} H_{\tilde{b}\tilde{a}}^c \quad (17.11)$$

$$L_{\tilde{b}\tilde{a}}^{(1)a} = \zeta h^{0\tilde{d}\tilde{c}} p^{ac} p^{bd} \ell_{be} (k_{\tilde{d}\tilde{a}}^0 \ell_{cd} H_{\tilde{b}\tilde{a}}^e + k_{\tilde{b}\tilde{d}}^0 \ell_{dc} H_{\tilde{c}\tilde{a}}^e) - 2\zeta k_{\tilde{d}\tilde{a}}^0 p^{ac} \ell_{cd} H_{\tilde{b}\tilde{c}}^d h^{0\tilde{d}\tilde{c}} \quad (17.12)$$

or

$$L_{\tilde{b}\tilde{a}}^{(1)a} = \zeta (\delta_e^a - \xi^2 p^{ab} k_{bc} p^{cd} k_{de}) \cdot h^{0\tilde{c}\tilde{d}} (k_{\tilde{b}\tilde{d}}^0 H_{\tilde{c}\tilde{a}}^e - k_{\tilde{a}\tilde{d}}^0 H_{\tilde{c}\tilde{b}}^e) \quad (17.13)$$

$$\begin{aligned} L_{\tilde{b}\tilde{a}}^{(2)a} = \zeta^2 \xi h^{0\tilde{c}\tilde{d}} h^{0\tilde{t}\tilde{s}} (k_{+b}^a k_{-c}^b) p^{cr} k_{rd} \times \\ \times (k_{\tilde{b}\tilde{d}}^0 H_{\tilde{s}\tilde{b}}^d k_{\tilde{a}\tilde{b}}^0 - k_{\tilde{c}\tilde{t}}^0 k_{\tilde{d}\tilde{b}}^0 H_{\tilde{s}\tilde{a}}^d + 2k_{\tilde{b}\tilde{t}}^0 k_{\tilde{d}\tilde{a}}^0 H_{\tilde{s}\tilde{c}}^d) \end{aligned} \quad (17.14)$$

where

$$k_{\pm b}^a = \delta_b^a \pm \xi p^{ar} k_{rb} \quad (17.15)$$

Let us consider the potential for the Higgs' field $V(\Phi)$ and expand it into a power series with respect to ζ .

$$V(\Phi) = V^{(0)}(\Phi) + V^{(1)}(\Phi) + V^{(2)}(\Phi) + \dots \quad (17.16)$$

Using (17.12–14) and (17.2b) one gets after some calculations:

$$V^{(0)}(\Phi) = \int_M \frac{\sqrt{h^0}}{V_2} dm(y) (p_{cb} + \xi^2 k_{cr} p^{rs} k_{sb}) H^c_{\tilde{b}\tilde{c}} H^b_{\tilde{m}\tilde{a}} h^{0\tilde{b}\tilde{m}} h^{0\tilde{c}\tilde{a}} \quad (17.17)$$

$$V^{(1)}(\Phi) = \int_M \frac{\sqrt{h^0}}{V_2} 2\zeta \xi dm(y) (p_{ab} + \xi k_{ab}) h^{0\tilde{b}\tilde{s}} h^{0\tilde{m}\tilde{t}} h^{0\tilde{c}\tilde{a}} p^{ar} \times (k_{rs} + \xi k_{rp} p^{pq} k_{qs}) k^0_{\tilde{t}\tilde{s}} H^s_{\tilde{b}\tilde{c}} H^b_{\tilde{m}\tilde{a}} \quad (17.18)$$

$$V^{(2)}(\Phi) = \frac{\zeta^2}{V_2} \int_M \sqrt{h^0} dm(y) (p_{ab} + \xi k_{ab}) H^s_{\tilde{b}\tilde{c}} H^b_{\tilde{m}\tilde{a}} k^0_{\tilde{t}\tilde{s}} k^0_{\tilde{d}\tilde{e}} \times \left[(2h^{0\tilde{b}\tilde{e}} h^{0\tilde{m}\tilde{t}} h^{0\tilde{s}\tilde{d}} h^{0\tilde{a}\tilde{c}} + h^{0\tilde{b}\tilde{s}} h^{0\tilde{m}\tilde{t}} h^{0\tilde{c}\tilde{e}} h^{0\tilde{a}\tilde{d}}) k^a_{+s} + \right. \\ + (h^{0\tilde{e}\tilde{s}} h^{0\tilde{m}\tilde{t}} h^{0\tilde{c}\tilde{a}} h^{0\tilde{b}\tilde{d}} - h^{0\tilde{d}\tilde{s}} h^{0\tilde{m}\tilde{t}} h^{0\tilde{c}\tilde{a}} h^{0\tilde{b}\tilde{e}} + \\ + h^{0\tilde{c}\tilde{s}} h^{0\tilde{m}\tilde{t}} h^{0\tilde{e}\tilde{a}} h^{0\tilde{b}\tilde{d}} - h^{0\tilde{c}\tilde{s}} h^{0\tilde{m}\tilde{t}} h^{0\tilde{d}\tilde{a}} h^{0\tilde{b}\tilde{e}}) k^a_{+r} k^r_{-s} + \\ + \frac{\xi}{2} (h^{0\tilde{c}\tilde{m}} h^{0\tilde{t}\tilde{a}} h^{0\tilde{d}\tilde{b}} h^{0\tilde{e}\tilde{s}} + h^{0\tilde{c}\tilde{m}} h^{0\tilde{t}\tilde{a}} h^{0\tilde{e}\tilde{b}} h^{0\tilde{d}\tilde{s}} + \\ - h^{0\tilde{c}\tilde{m}} h^{0\tilde{s}\tilde{a}} h^{0\tilde{d}\tilde{b}} h^{0\tilde{e}\tilde{t}} - h^{0\tilde{c}\tilde{m}} h^{0\tilde{s}\tilde{a}} h^{0\tilde{e}\tilde{b}} h^{0\tilde{d}\tilde{t}}) k^a_{+r} k^r_{-s} + \\ + \frac{\xi}{2} (h^{0\tilde{c}\tilde{m}} h^{0\tilde{t}\tilde{a}} h^{0\tilde{d}\tilde{b}} h^{0\tilde{e}\tilde{s}} + h^{0\tilde{c}\tilde{m}} h^{0\tilde{t}\tilde{a}} h^{0\tilde{e}\tilde{b}} h^{0\tilde{d}\tilde{s}} + \\ - h^{0\tilde{c}\tilde{m}} h^{0\tilde{s}\tilde{a}} h^{0\tilde{d}\tilde{b}} h^{0\tilde{e}\tilde{t}} - h^{0\tilde{c}\tilde{m}} h^{0\tilde{s}\tilde{a}} h^{0\tilde{e}\tilde{b}} h^{0\tilde{d}\tilde{t}}) (k^a_{+r} k^r_{-s}) p^{pq} k_{qs} + \\ + \frac{\xi}{2} (h^{0\tilde{c}\tilde{d}} h^{0\tilde{t}\tilde{b}} h^{0\tilde{s}\tilde{m}} h^{0\tilde{e}\tilde{a}} - h^{0\tilde{c}\tilde{d}} h^{0\tilde{s}\tilde{b}} h^{0\tilde{t}\tilde{m}} h^{0\tilde{e}\tilde{a}} + \\ - h^{0\tilde{c}\tilde{e}} h^{0\tilde{t}\tilde{b}} h^{0\tilde{s}\tilde{m}} h^{0\tilde{d}\tilde{a}} + h^{0\tilde{c}\tilde{e}} h^{0\tilde{s}\tilde{b}} h^{0\tilde{t}\tilde{m}} h^{0\tilde{d}\tilde{a}}) (k^a_{+r} k^r_{-p}) p^{pq} k_{qs} + \\ - \frac{\xi}{2} (h^{0\tilde{s}\tilde{m}} h^{0\tilde{a}\tilde{t}} h^{0\tilde{b}\tilde{e}} h^{0\tilde{c}\tilde{d}} - h^{0\tilde{m}\tilde{t}} h^{0\tilde{a}\tilde{s}} h^{0\tilde{b}\tilde{e}} h^{0\tilde{c}\tilde{d}} + \\ \left. - h^{0\tilde{s}\tilde{m}} h^{0\tilde{a}\tilde{t}} h^{0\tilde{b}\tilde{e}} h^{0\tilde{c}\tilde{e}} + h^{0\tilde{m}\tilde{t}} h^{0\tilde{a}\tilde{s}} h^{0\tilde{b}\tilde{d}} h^{0\tilde{c}\tilde{e}}) \delta^a_s \right] \quad (17.19)$$

Up to the first order of approximation in ζ one gets:

$$V(\Phi) = \int_M [(p_{cb} + \xi^2 k_{cr} p^{rs} k_{sb}) H^c_{\tilde{b}\tilde{c}} H^b_{\tilde{m}\tilde{a}} h^{0\tilde{b}\tilde{m}} h^{0\tilde{c}\tilde{a}} + \\ + 2\xi \zeta p^{ar} k_{rs} (p_{ab} + \xi k_{ab}) (\delta^s_p + \xi p^{sq} k_{qp}) \times \\ \times h^{0\tilde{b}\tilde{s}} h^{0\tilde{m}\tilde{t}} h^{0\tilde{c}\tilde{a}} k^o_{\tilde{t}\tilde{s}} H^p_{\tilde{m}\tilde{c}} H^b_{\tilde{m}\tilde{c}}] \frac{\sqrt{|h^0|}}{V_2} dm(y). \quad (17.20)$$

It is easy to see that $V(\Phi)$ has only the first type of the critical point corresponding to the trivial gauge configuration for the Higgs' field up to the first order of expansion

with respect to ζ . $V^{(2)}(\Phi)$ gives a contribution to the Higgs' potential resulting in the second type of critical point for the potential corresponding to the nontrivial gauge configuration of the Higgs' field.

Let us consider Eq. (17.3a) and rewrite it in a more convenient form.

$$\begin{aligned} \ell_{dc}(\delta_\beta^\gamma - 2g_{[\beta\delta]}g^{\gamma\delta})L^d_{\gamma\tilde{a}} + \ell_{cd}(\delta_{\tilde{a}}^{\tilde{c}} - 2\zeta k^o_{\tilde{d}\tilde{a}}\tilde{g}^{\tilde{d}\tilde{c}})L^d_{\beta\tilde{c}} = \\ = 2\ell_{cd}(\delta_{\tilde{a}}^{\tilde{c}} - 2\zeta k^o_{\tilde{d}\tilde{a}}\tilde{g}^{\tilde{d}\tilde{c}})\overset{\text{gauge}}{\nabla}_\beta\Phi_{\tilde{c}}^d. \end{aligned} \quad (17.21)$$

Let us expand $L^d_{\mu\tilde{c}}$ into a power series with respect to ζ and with respect to $h_{\mu\nu} = g_{\mu\nu} - \eta_{\mu\nu}$

$$\begin{aligned} L^d_{\mu\tilde{c}} &= L^{(0)}_{\mu\tilde{c}} + L^{(1)}_{\mu\tilde{c}} + L^{(2)}_{\mu\tilde{c}} + \dots, \\ g^{\mu\alpha}g_{\nu\alpha} &= (h^{\mu\alpha} + \overset{1}{h}^{\mu\alpha} + \overset{2}{h}^{\mu\alpha} + \dots). \end{aligned} \quad (17.22)$$

Using (17.21), (17.5), (17.6) one gets up to the second order in ζ and $h_{\mu\nu}$.

$$\begin{aligned} \ell_{dc}L^d_{\beta\tilde{a}} + \ell_{cd}L^d_{\beta\tilde{a}} - 2\zeta k^o_{\tilde{d}\tilde{a}}(h^{o\tilde{d}\tilde{c}} - \zeta h^{o\tilde{d}\tilde{p}}h^{o\tilde{c}\tilde{b}}k^o_{\tilde{b}\tilde{p}} + \zeta^2 h^{o\tilde{c}\tilde{b}}h^{o\tilde{d}\tilde{e}}h^{o\tilde{p}\tilde{s}}k^o_{\tilde{s}\tilde{e}}k^o_{\tilde{b}\tilde{p}})\ell_{cd}L^d_{\beta\tilde{c}} + \\ - 2h_{[\beta\delta]}(\eta^{\gamma\delta} - \eta^{\gamma\alpha}\eta^{\delta\mu}h_{\mu\alpha} + \eta^{\gamma\mu}\eta^{\delta\sigma}\eta^{\alpha\nu}h_{\nu\mu}h_{\delta\alpha})\ell_{dc}L^d_{\gamma\tilde{a}} = \\ = 2\ell_{cd}\overset{\text{gauge}}{\nabla}_\beta\Phi_{\tilde{a}}^d - 4\zeta\ell_{cd}k^o_{\tilde{d}\tilde{a}}(h^{o\tilde{d}\tilde{c}} - \zeta h^{o\tilde{d}\tilde{p}}h^{o\tilde{c}\tilde{b}}k^o_{\tilde{b}\tilde{p}} + \\ + \zeta^2 h^{o\tilde{c}\tilde{b}}h^{o\tilde{d}\tilde{e}}h^{o\tilde{p}\tilde{s}}k^o_{\tilde{s}\tilde{e}}k^o_{\tilde{b}\tilde{p}})\overset{\text{gauge}}{\nabla}_\beta\Phi_{\tilde{c}}^b. \end{aligned} \quad (17.23)$$

From (17.23) one obtains:

$$L^a_{\mu\tilde{c}} = p^{ab}\ell_{bc}\overset{\text{gauge}}{\nabla}_\mu\Phi_{\tilde{c}}^c = \overset{\text{gauge}}{\nabla}_\mu\Phi_{\tilde{c}}^a + \xi p^{ab}k_{bc}\overset{\text{gauge}}{\nabla}_\mu\Phi_{\tilde{c}}^c, \quad (17.24)$$

$$\begin{aligned} L^1_{\beta\tilde{a}} &= \zeta k^o_{\tilde{d}\tilde{a}}h^{o\tilde{d}\tilde{c}}p^{fc}(\ell_{cd}p^{db}\ell_{bp} - 2\ell_{cp})\overset{\text{gauge}}{\nabla}_\beta\Phi_{\tilde{c}}^p + \\ &+ \eta^{\gamma\delta}h_{[\beta\delta]}p^{fc}p^{db}\ell_{bp}\ell_{cd}\overset{\text{gauge}}{\nabla}_\gamma\Phi_{\tilde{a}}^p, \end{aligned} \quad (17.25)$$

$$\begin{aligned} L^m_{\beta\tilde{a}} &= \zeta^2 p^{cm}k^o_{\tilde{d}\tilde{a}}\ell_{cd}h^{o\tilde{d}\tilde{p}}h^{o\tilde{c}\tilde{q}}[-k^o_{\tilde{q}\tilde{p}}p^{db}\ell_{bf}\overset{\text{gauge}}{\nabla}_\beta\Phi_{\tilde{c}}^f + \\ &+ k^o_{\tilde{c}\tilde{p}}(p^{dn}\ell_{nq}p^{qp}\ell_{bp}\overset{\text{gauge}}{\nabla}_\beta\Phi_{\tilde{q}}^p + p^{dq}\ell_{qp}\overset{\text{gauge}}{\nabla}_\beta\Phi_{\tilde{q}}^p + 2\overset{\text{gauge}}{\nabla}_\beta\Phi_{\tilde{c}}^d)] + \\ &+ \zeta p^{cm}\eta^{\gamma\delta}h_{[\beta\delta]}k^o_{\tilde{d}\tilde{a}}h^{o\tilde{d}\tilde{c}}(\ell_{cd}p^{nd}p^{qb}\ell_{bp}\ell_{nq} + \\ &+ \ell_{dc}p^{dq}\ell_{qa}p^{ab}\ell_{bp} + p^{dq}\ell_{dc}\ell_{qp})\overset{\text{gauge}}{\nabla}_\gamma\Phi_{\tilde{q}}^p + \\ &+ h_{[\beta\delta]}\eta^{\delta\mu}\eta^{\gamma\alpha}p^{cm}\ell_{dc}\ell_{bp}(h_{\mu\alpha}p^{db} + h_{[\mu\alpha]}p^{dq}p^{ab}\ell_{qa})\overset{\text{gauge}}{\nabla}_\gamma\Phi_{\tilde{a}}^p. \end{aligned} \quad (17.26)$$

Let us expand $\mathcal{L}_{\text{kin}}(\overset{\text{gauge}}{\nabla}\Phi)$ into a power series with respect to ζ and $h_{\mu\nu}$.

$$\mathcal{L}_{\text{kin}}(\overset{\text{gauge}}{\nabla}\Phi) = \mathcal{L}_{\text{kin}}^{(0)}(\overset{\text{gauge}}{\nabla}\Phi) + \mathcal{L}_{\text{kin}}^{(1)}(\overset{\text{gauge}}{\nabla}\Phi) + \mathcal{L}_{\text{kin}}^{(2)}(\overset{\text{gauge}}{\nabla}\Phi) + \dots \quad (17.27)$$

Using (17.2a) and (17.24–26) one gets:

$$\begin{aligned} \mathcal{L}_{\text{kin}}^{(0)}(\overset{\text{gauge}}{\nabla} \Phi) &= \frac{1}{V_2} \int_M \sqrt{|h^0|} dm(y) \eta^{\alpha\mu} h^{0\tilde{b}\tilde{n}} (\ell_{ab} p^{af} \ell_{fc}) \overset{\text{gauge}}{\nabla}_\alpha \Phi_b^c \overset{\text{gauge}}{\nabla}_\mu \Phi_n^b = \\ &= \int_M [\eta^{\alpha\mu} h^{0\tilde{b}\tilde{n}} (p_{cb} + \xi^2 p^{af} k_{ab} k_{fc}) \overset{\text{gauge}}{\nabla}_\alpha \Phi_b^c \overset{\text{gauge}}{\nabla}_\mu \Phi_n^b \sqrt{|h^0|} dm(y)], \quad (17.28) \end{aligned}$$

$$\begin{aligned} \mathcal{L}_{\text{kin}}^{(1)}(\overset{\text{gauge}}{\nabla} \Phi) &= \frac{1}{V_2} \int_M \{ -\zeta \eta^{\alpha\mu} (p_{cb} + \xi^2 p^{af} k_{ab} k_{fc}) k^o_{\tilde{d}\tilde{b}} h^{o\tilde{d}\tilde{n}} h^{o\tilde{b}\tilde{m}} \times \\ &\times (\overset{\text{gauge}}{\nabla}_\alpha \Phi_{\tilde{m}}^c \overset{\text{gauge}}{\nabla}_\mu \Phi_n^b + p^{ce} \ell_{ed} p^{db} \ell_{bp} \overset{\text{gauge}}{\nabla}_\alpha \Phi_{\tilde{m}}^p \overset{\text{gauge}}{\nabla}_\mu \Phi_n^b + \\ &- 2(p_{pb} + \xi^2 p^{qg} k_{qb} k_{gd}) p^{pi} \ell_{id} \overset{\text{gauge}}{\nabla}_\mu \Phi_n^b \overset{\text{gauge}}{\nabla}_\alpha \Phi_{\tilde{m}}^d] + \\ &- (p_{cb} + \xi^2 p^{af} k_{ab} k_{fc}) \eta^{\alpha\mu} \eta^{\gamma\delta} h^{o\tilde{b}\tilde{n}} \times \\ &\times [h_{\delta\alpha} \overset{\text{gauge}}{\nabla}_\mu \Phi_b^c \overset{\text{gauge}}{\nabla}_\gamma \Phi_n^b + h_{[\delta\alpha]} p^{ce} (p_{ep} + \xi^2 p^{df} k_{dp} k_{fe}) \times \\ &\times \overset{\text{gauge}}{\nabla}_\mu \Phi_n^b \overset{\text{gauge}}{\nabla}_\gamma \Phi_{\tilde{b}}^p] \} \sqrt{|h^0|} dm(y), \quad (17.29) \end{aligned}$$

$$\begin{aligned} \mathcal{L}_{\text{kin}}^{(2)}(\overset{\text{gauge}}{\nabla} \Phi) &= \int_M \frac{\sqrt{|h^0|}}{V_2} dm(y) \times \\ &\times [\zeta^2 (\eta^{\alpha\mu} \ell_{ab} p^{ad} \ell_{dc} h^{o\tilde{b}\tilde{c}} h^{o\tilde{n}\tilde{s}} h^{o\tilde{a}\tilde{p}} k^o_{\tilde{p}\tilde{c}} k^o_{\tilde{s}\tilde{a}} \overset{\text{gauge}}{\nabla}_\mu \Phi_n^b \overset{\text{gauge}}{\nabla}_\alpha \Phi_b^c + \\ &+ \eta^{\alpha\mu} h^{o\tilde{b}\tilde{r}} h^{o\tilde{n}\tilde{s}} h^{o\tilde{d}\tilde{c}} k^o_{\tilde{d}\tilde{b}} k^o_{\tilde{r}\tilde{s}} \ell_{ab} p^{ac} \ell_{cd} p^{dq} \ell_{qp} \overset{\text{gauge}}{\nabla}_\alpha \Phi_{\tilde{c}}^p \overset{\text{gauge}}{\nabla}_\mu \Phi_n^b + \\ &+ 2\eta^{\alpha\mu} h^{o\tilde{b}\tilde{r}} h^{o\tilde{m}\tilde{s}} h^{o\tilde{d}\tilde{c}} k^o_{\tilde{s}\tilde{r}} k^o_{\tilde{d}\tilde{b}} \ell_{ab} p^{ac} \ell_{cd} \overset{\text{gauge}}{\nabla}_\alpha \Phi_{\tilde{c}}^d \overset{\text{gauge}}{\nabla}_\mu \Phi_n^b) + \\ &+ \zeta (2\eta^{\alpha\rho} \eta^{\mu\beta} h_{\beta\rho} h^{o\tilde{b}\tilde{n}} h^{o\tilde{d}\tilde{c}} k^o_{\tilde{d}\tilde{b}} \ell_{ab} p^{ac} \ell_{cd} \overset{\text{gauge}}{\nabla}_\mu \Phi_n^b \overset{\text{gauge}}{\nabla}_\alpha \Phi_{\tilde{c}}^d + \\ &+ \eta^{\alpha\rho} \eta^{\mu\beta} h_{\beta\rho} h^{o\tilde{b}\tilde{a}} h^{o\tilde{n}\tilde{r}} k^o_{\tilde{r}\tilde{a}} \ell_{ab} p^{ac} \ell_{cd} \overset{\text{gauge}}{\nabla}_\mu \Phi_n^b \overset{\text{gauge}}{\nabla}_\alpha \Phi_{\tilde{c}}^c + \\ &- \eta^{\alpha\rho} \eta^{\mu\beta} h_{\beta\rho} h^{o\tilde{b}\tilde{n}} h^{o\tilde{d}\tilde{c}} k^o_{\tilde{d}\tilde{b}} \ell_{ab} p^{ac} \ell_{cd} \ell_{dq} \overset{\text{gauge}}{\nabla}_\mu \Phi_n^b \overset{\text{gauge}}{\nabla}_\alpha \Phi_{\tilde{c}}^p + \\ &- \eta^{\alpha\mu} \eta^{\gamma\delta} h_{[\alpha\delta]} h^{o\tilde{b}\tilde{r}} h^{o\tilde{m}\tilde{s}} k^o_{\tilde{s}\tilde{r}} \ell_{ab} p^{ac} p^{dq} \ell_{qp} \ell_{cd} \overset{\text{gauge}}{\nabla}_\gamma \Phi_{\tilde{b}}^p \overset{\text{gauge}}{\nabla}_\mu \Phi_{\tilde{m}}^b + \\ &+ \eta^{\alpha\mu} \eta^{\gamma\delta} h_{[\alpha\delta]} h^{o\tilde{b}\tilde{n}} h^{o\tilde{d}\tilde{c}} k^o_{\tilde{d}\tilde{b}} \ell_{ab} p^{ca} \ell_{cd} p^{nd} p^{qm} \ell_{mp} \ell_{nq} \times \\ &\times \overset{\text{gauge}}{\nabla}_\mu \Phi_n^b \overset{\text{gauge}}{\nabla}_\gamma \Phi_{\tilde{c}}^p + \\ &+ \eta^{\alpha\mu} \eta^{\gamma\delta} h_{[\alpha\delta]} h^{o\tilde{b}\tilde{n}} h^{o\tilde{d}\tilde{c}} k^o_{\tilde{d}\tilde{b}} \ell_{ab} p^{ca} \ell_{dc} p^{dq} \ell_{qn} p^{nm} \ell_{mp} \times \\ &\times \overset{\text{gauge}}{\nabla}_\mu \Phi_n^b \overset{\text{gauge}}{\nabla}_\gamma \Phi_{\tilde{c}}^p + \\ &+ \eta^{\alpha\mu} \eta^{\gamma\delta} h_{[\alpha\delta]} h^{o\tilde{b}\tilde{n}} h^{o\tilde{d}\tilde{c}} k^o_{\tilde{d}\tilde{b}} \ell_{ab} p^{ca} \ell_{dc} p^{dq} \ell_{qp} \overset{\text{gauge}}{\nabla}_\mu \Phi_n^b \overset{\text{gauge}}{\nabla}_\gamma \Phi_{\tilde{c}}^p) + \\ &+ h^{o\tilde{n}\tilde{b}} (\eta^{\alpha\mu} \eta^{\delta\nu} \eta^{\gamma\rho} h_{[\alpha\delta]} h_{(\nu\rho)} \ell_{ab} p^{ca} \ell_{dc} \ell_{np} p^{dq} p^{nm} \ell_{qm} \times \end{aligned}$$

$$\begin{aligned}
& \times \nabla_\mu^{\text{gauge}} \Phi_{\tilde{n}}^b \nabla_\gamma^{\text{gauge}} \Phi_{\tilde{b}}^p + \\
& + 2\eta^{\alpha\mu} \eta^{\delta\nu} \eta^{\gamma\rho} h_{[\alpha\delta]} h_{[\nu\rho]} \ell_{ab} p^{ca} \ell_{dc} \ell_{np} p^{dq} p^{mn} \ell_{qm} \times \\
& \times \nabla_\mu^{\text{gauge}} \Phi_{\tilde{n}}^b \nabla_\gamma^{\text{gauge}} \Phi_{\tilde{b}}^p + \\
& + \eta^{\alpha\gamma} \eta^{\mu\sigma} \eta^{\rho\beta} h_{\beta\gamma} h_{\sigma\rho} \ell_{ab} p^{\text{ad}} \ell_{dc} \nabla_\mu^{\text{gauge}} \Phi_{\tilde{n}}^b \nabla_\alpha^{\text{gauge}} \Phi_{\tilde{c}}^c + \\
& - \eta^{\alpha\rho} \eta^{\mu\beta} \eta^{\gamma\delta} h_{\beta\rho} h_{[\gamma\delta]} \ell_{ab} p^{ac} p^{dq} \ell_{qp} \ell_{cd} \nabla_\mu^{\text{gauge}} \Phi_{\tilde{n}}^b \nabla_\alpha^{\text{gauge}} \Phi_{\tilde{b}}^p]. \quad (17.30)
\end{aligned}$$

It is easy to see that the skewon field $h_{[\mu\nu]}$ enters the lagrangian $\mathcal{L}_{\text{kin}}(\nabla^{\text{gauge}} \Phi)$ in the first order of approximation similarly as for the Yang–Mills' field. Moreover it couples here the gauge derivative of the Higgs' field.

Let us consider $\mathcal{L}_{\text{int}}(\Phi, \tilde{A})$. One gets:

$$\mathcal{L}_{\text{int}}^{(0)}(\Phi, \tilde{A}) = \mathcal{L}_{\text{int}}^{(1)}(\Phi, \tilde{A}) = 0 \quad (17.31)$$

and

$$\begin{aligned}
\mathcal{L}_{\text{int}}^{(2)}(\Phi, \tilde{A}) = & \frac{\zeta}{V_2} \int_M \sqrt{|h^0|} dm(y) \times \\
& \times (p_{ab} \mu_i^a \eta^{\mu\alpha} \eta^{\nu\beta} h_{\beta\alpha} h^{o\tilde{m}\tilde{a}} h^{o\tilde{n}\tilde{b}} k_{\tilde{b}\tilde{a}}^o \tilde{H}_{\mu\nu}^i H_{\tilde{m}\tilde{n}}^b). \quad (17.32)
\end{aligned}$$

The nonminimal interaction term enters from the second order of approximation. (17.32) gives us an interpretation of a constant ζ . It plays the role of a coupling constant between the skewon field $h_{[\mu\nu]}$ and the Higgs' field. The coupling constant ζ is simultaneously connected to a part of the cosmological constant in our theory. Thus we get an interesting relation between cosmological constant and coupling constants ζ and ξ between skewon field $h_{[\tilde{m}\nu]}$ and Higgs' and Yang–Mills' fields.

Thus we get $V(\Phi)$, $\mathcal{L}_{\text{kin}}(\nabla^{\text{gauge}} \Phi)$, $\mathcal{L}_{\text{int}}(\Phi, \tilde{A})$ up to the second order in an approximation with respect to ζ and $h_{\mu\nu} = g_{\mu\nu} - \eta_{\tilde{m}\nu}$. The higher order terms can be obtained in a similar way.

Let us give a remark concerning a convergence of series appearing here. They are double power series with respect to ζ and $h_{\mu\nu}$ and they converge for sufficiently small ζ and $h_{\mu\nu}$. However all the functions of ζ and $h_{\mu\nu}$ considered here (i.e. $L_{\tilde{b}\tilde{c}}^a$, $L_{\alpha\tilde{a}}^a$, $g^{\tilde{a}\tilde{b}}$, $g^{\mu\nu}$, $\mathcal{L}_{\text{kin}}(\nabla^{\text{gauge}} \Phi)$, $V(\Psi)$, $\mathcal{L}_{\text{int}}(\Phi, \tilde{A})$) are well defined for any ζ and $h_{\mu\nu}$. They are rational functions of these variables. Moreover the exact form of all of these functions is hard to get.

Let us consider the terms involving Higgs' field in the geodetic equation in the first order of approximation with respect to ζ and $h_{\mu\nu}$ i.e. the fourth term in Eq. (15.2) and

Eq. (15.3). One gets:

$$\begin{aligned}
& \left(\frac{q^c}{m_o} \right) u^{\tilde{b}} [\eta^{\alpha\delta} \ell_{cd} \overset{\text{gauge}}{\nabla}_\delta \Phi_{\tilde{b}}^d - \xi \eta^{\alpha\delta} k_{cd} p^{db} \ell_{bf} \overset{\text{gauge}}{\nabla}_\delta \Phi_{\tilde{b}}^f + \\
& - \zeta \xi \eta^{\alpha\delta} k_{cd} k^o_{\tilde{a}\tilde{b}} h^{o\tilde{d}\tilde{c}} p^{df} (\ell_{fe} p^{eq} \ell_{qp} - 2\ell_{fp}) \overset{\text{gauge}}{\nabla}_\delta \Phi_{\tilde{c}}^p + \\
& - \eta^{\alpha\rho} \eta^{\delta\beta} h_{\beta\rho} \ell_{cd} \overset{\text{gauge}}{\nabla}_\delta \Phi_{\tilde{b}}^d - \frac{1}{2} (\ell_{cd} \eta^{\alpha\delta} \eta^{\rho\beta} - \ell_{dc} \eta^{\delta\rho} \eta^{\alpha\beta}) \times \\
& \times h_{\beta\rho} p^{db} \ell_{bf} \overset{\text{gauge}}{\nabla}_\delta \Phi_{\tilde{b}}^f - \xi \eta^{\alpha\delta} \eta^{\gamma\rho} h_{[\delta\rho]} k_{cd} p^{df} p^{eb} \ell_{fe} \ell_{bp} \overset{\text{gauge}}{\nabla}_\gamma \Phi_{\tilde{b}}^p]. \quad (17.33)
\end{aligned}$$

This is the fourth term in Eq. (15.2) i.e. the term involving the Higgs' field in the first order of approximation. Let us write the equation for the Higgs' charge in the first order of approximation. One easily finds:

$$\begin{aligned}
& \frac{\tilde{D}u^{\tilde{a}}}{d\tau} + \frac{1}{r^2} \left(\frac{q^c}{m_o} \right) u^\beta \{ \ell_{cd} h^{o\tilde{a}\tilde{d}} \overset{\text{gauge}}{\nabla}_\beta \Phi_{\tilde{d}}^d - \xi k_{cd} h^{o\tilde{a}\tilde{d}} p^{db} \ell_{bp} \overset{\text{gauge}}{\nabla}_\beta \Phi_{\tilde{d}}^p + \\
& + \frac{1}{2} \zeta [(\ell_{cd} h^{o\tilde{a}\tilde{n}} h^{o\tilde{d}\tilde{m}} - \ell_{dc} h^{o\tilde{d}\tilde{n}} h^{o\tilde{a}\tilde{m}}) k^o_{\tilde{m}\tilde{n}} p^{db} \ell_{bp} \overset{\text{gauge}}{\nabla}_\beta \Phi_{\tilde{d}}^p + \\
& - 2h^{o\tilde{a}\tilde{n}} h^{o\tilde{d}\tilde{m}} k^o_{\tilde{m}\tilde{n}} \overset{\text{gauge}}{\nabla}_\beta \Phi_{\tilde{d}}^d] + \\
& - \zeta \xi k_{cd} h^{o\tilde{a}\tilde{d}} k^o_{\tilde{f}\tilde{d}} h^{o\tilde{f}\tilde{c}} p^{fe} (\ell_{ep} p^{pr} \ell_{rq} - 2\ell_{eq}) \overset{\text{gauge}}{\nabla}_\beta \Phi_{\tilde{c}}^q + \\
& - \xi \eta^{\gamma\delta} h_{[\beta\delta]} k_{cd} p^{dr} p^{sp} \ell_{pf} \ell_{rs} \overset{\text{gauge}}{\nabla}_\beta \Phi_{\tilde{d}}^f \} + \\
& + \frac{1}{r^2} \left(\frac{q^c}{m_o} \right) u^{\tilde{b}} \{ \ell_{cd} h^{o\tilde{a}\tilde{d}} \overset{\text{gauge}}{\nabla}_\beta H_{\tilde{d}\tilde{b}}^d - \xi k_{cd} h^{o\tilde{a}\tilde{d}} p^{df} \ell_{fe} H_{\tilde{b}\tilde{d}}^e + \\
& + \frac{1}{2} \zeta [(\ell_{cd} h^{o\tilde{a}\tilde{n}} h^{o\tilde{d}\tilde{m}} - \ell_{dc} h^{o\tilde{d}\tilde{n}} h^{o\tilde{a}\tilde{m}}) k^o_{\tilde{m}\tilde{n}} p^{df} \ell_{fe} H_{\tilde{b}\tilde{d}}^e + \\
& - 2h^{o\tilde{a}\tilde{n}} h^{o\tilde{d}\tilde{m}} k^o_{\tilde{m}\tilde{n}} H_{\tilde{d}\tilde{b}}^d] + \\
& - \zeta \xi k_{cd} h^{o\tilde{a}\tilde{d}} (h^{o\tilde{f}\tilde{c}} p^{dp} p^{bq} \ell_{be} (k^o_{\tilde{f}\tilde{d}} \ell_{pq} H_{\tilde{b}\tilde{c}}^e + k^o_{\tilde{b}\tilde{f}} \ell_{qp} H_{\tilde{c}\tilde{d}}^e) + \\
& - 2k^o_{\tilde{e}\tilde{d}} h^{o\tilde{e}\tilde{c}} p^{dp} \ell_{pq} H_{\tilde{b}\tilde{c}}^q) \} \}. \quad (17.34)
\end{aligned}$$

Eventually one easily writes down the explicit form of the mass matrix for massive vector bosons (i.e. $M^2_{ij}(\Phi_{\text{crt}}^k)$ from (8.21–22) in zeroth order of expansion with respect ζ .

$$\begin{aligned}
M^2_{ij}(\Phi_{\text{crt}}^k) &= - \left(\frac{\alpha_s}{r} \right)^2 \left(\frac{\hbar}{c} \right)^2 h^{o\tilde{b}\tilde{n}} (\ell_{ab} p^{af} \ell_{fc}) \cdot \\
&\cdot ((\Phi_{\text{crt}}^k)^p_{\tilde{b}} C^c_{pq} \alpha_j^q + (\Phi_{\text{crt}}^k)^c_{\tilde{p}} f_{\tilde{b}j}^{\tilde{p}}) \cdot ((\Phi_{\text{crt}}^k)^p_{\tilde{b}} C^b_{ms} \alpha_i^s + (\Phi_{\text{crt}}^k)^b_{\tilde{m}} f_{\tilde{n}i}^{\tilde{m}}). \quad (17.35)
\end{aligned}$$

The next terms i.e. the first and the second can be obtained similarly from Eq. (17.28–30). Thus we get an expansion with respect to ζ

$$M^2_{ij}(\Phi^k_{\text{crt}}) = \overset{(0)}{M^2}_{ij}(\Phi^k_{\text{crt}}) + \overset{(1)}{M^2}_{ij}(\Phi^k_{\text{crt}}) + \overset{(2)}{M^2}_{ij}(\Phi^k_{\text{crt}}) + \dots, \quad (17.36)$$

where

$$\begin{aligned} \overset{(1)}{M^2}_{ij}(\Phi^k_{\text{crt}}) = & \zeta \int_M \left(\frac{\alpha_s}{r} \right)^2 \left(\frac{\hbar}{c} \right)^2 [(p_{cb} + \xi^2 p^{af} k_{ab} k_{fc}) \cdot k^o_{\bar{d}\bar{b}} h^{o\bar{d}\bar{n}} h^{o\bar{b}\bar{m}} \cdot \\ & \cdot ((\Phi^k_{\text{crt}})^p_{\bar{m}} C^c_{pq} \alpha^q_j + (\Phi^k_{\text{crt}})^c_{\bar{p}} f^{\bar{p}}_{\bar{m}j}) \cdot ((\Phi^k_{\text{crt}})^s_{\bar{n}} C^c_{sr} \alpha^r_i + (\Phi^k_{\text{crt}})^b_{\bar{q}} f^{\bar{q}}_{\bar{n}i}) + \\ & + p^{ce} \ell_{ed} p^{db} \ell_{bp} \cdot \\ & \cdot ((\Phi^k_{\text{crt}})^u_{\bar{m}} C^p_{uw} \alpha^w_j + (\Phi^k_{\text{crt}})^p_{\bar{u}} f^{\bar{u}}_{\bar{m}j}) \cdot ((\Phi^k_{\text{crt}})^v_{\bar{n}} C^b_{vz} \alpha^z_i + (\Phi^k_{\text{crt}})^b_{\bar{v}} f^{\bar{v}}_{\bar{n}i}) + \\ & - 2(p_{pb} + \zeta^2 p^{qg} k_{qb} h_{gp}) p^{pl} \ell_{ld} \cdot \\ & \cdot ((\Phi^k_{\text{crt}})^w_{\bar{n}} C^b_{wq} \alpha^q_j + (\Phi^k_{\text{crt}})^b_{\bar{v}} f^{\bar{v}}_{\bar{n}j}) \cdot ((\Phi^k_{\text{crt}})^z_{\bar{n}} C^d_{zv} \alpha^v_i + (\Phi^k_{\text{crt}})^d_{\bar{p}} f^{\bar{p}}_{\bar{n}i})] \cdot \\ & \cdot \frac{\sqrt{|h^0|}}{V_2} dm(y), \end{aligned} \quad (17.37)$$

$$\begin{aligned} \overset{(2)}{M^2}_{ij}(\Phi^k_{\text{crt}}) = & \int_M \zeta^2 \left(\frac{\alpha_s}{r} \right)^2 \left(\frac{\hbar}{c} \right)^2 [(\ell_{ab} + p^{ad} \ell_{dc} h^{o\bar{b}\bar{c}} h^{o\bar{n}\bar{s}} h^{o\bar{a}\bar{p}} k^o_{\bar{p}\bar{c}} k^o_{\bar{s}\bar{a}} \cdot \\ & \cdot ((\Phi^k_{\text{crt}})^u_{\bar{n}} C^b_{uq} \alpha^q_j + (\Phi^k_{\text{crt}})^b_{\bar{m}} f^{\bar{m}}_{\bar{n}j}) \cdot ((\Phi^k_{\text{crt}})^v_{\bar{b}} C^c_{vw} \alpha^w_i + (\Phi^k_{\text{crt}})^c_{\bar{q}} f^{\bar{q}}_{\bar{b}i}) + \\ & + h^{o\bar{b}\bar{r}} h^{o\bar{n}\bar{s}} h^{o\bar{d}\bar{c}} k^o_{\bar{d}\bar{b}} k^o_{\bar{r}\bar{s}} \ell_{ab} p^{ce} \ell_{cd} h^{dq} \ell_{qp} \cdot \\ & \cdot ((\Phi^k_{\text{crt}})^r_{\bar{c}} C^p_{rf} \alpha^f_j + (\Phi^k_{\text{crt}})^p_{\bar{q}} f^{\bar{q}}_{\bar{c}j}) \cdot ((\Phi^k_{\text{crt}})^s_{\bar{n}} C^b_{sg} \alpha^g_i + (\Phi^k_{\text{crt}})^b_{\bar{m}} f^{\bar{m}}_{\bar{n}i}) + \\ & - 2h^{o\bar{b}\bar{r}} h^{o\bar{n}\bar{s}} h^{o\bar{d}\bar{c}} k^o_{\bar{s}\bar{r}} k^o_{\bar{d}\bar{b}} \ell_{ab} p^{ac} \ell_{cd} \cdot \\ & \cdot ((\Phi^k_{\text{crt}})^q_{\bar{c}} C^d_{qp} \alpha^p_j + (\Phi^k_{\text{crt}})^d_{\bar{m}} f^{\bar{m}}_{\bar{c}j}) \cdot ((\Phi^k_{\text{crt}})^e_{\bar{n}} C^b_{ef} \alpha^f_i + (\Phi^k_{\text{crt}})^b_{\bar{p}} f^{\bar{p}}_{\bar{n}i})] \cdot \\ & \cdot \frac{\sqrt{|h^0|}}{V_2} dm(y). \end{aligned} \quad (17.38)$$

After a diagonalization procedure for $M^2_{ij}(\Phi^k_{\text{crt}})$ we get a mass spectrum for vector bosons. For $k = 0$ in the “true” vacuum case and for $k = 1$ for the “false” vacuum case.

Let us consider the mass matrix for Higgs’ particles (see sec. 8)

$$m^2(\Phi^k_{\text{crt}})^{\bar{a}}_{\bar{a}} \tilde{r}_r = \frac{1}{4} \left(\frac{\delta^2 V}{\delta \Phi^{\bar{a}}_{\bar{a}} \delta \Phi^{\bar{r}}_{\bar{r}}} \Big|_{\Phi=\Phi^k_{\text{crt}}} \right) \quad (17.39)$$

and expand it in a power series with respect to ζ . One gets

$$m^2(\Phi^k_{\text{crt}})^{\bar{a}}_{\bar{a}} \tilde{r}_r = \overset{(0)}{m^2}(\Phi^k_{\text{crt}})^{\bar{a}}_{\bar{a}} \tilde{r}_r + \overset{(1)}{m^2}(\Phi^k_{\text{crt}})^{\bar{a}}_{\bar{a}} \tilde{r}_r + \overset{(2)}{m^2}(\Phi^k_{\text{crt}})^{\bar{a}}_{\bar{a}} \tilde{r}_r + \dots \quad (17.40)$$

One finds

$$\begin{aligned}
\bar{m}^{(0)2}(\Phi_{\text{crt}}^k)_{\tilde{a} \tilde{r}}^{\tilde{a} \tilde{r}} &= \frac{1}{r^4} \left\{ (p_{cd} + \xi^2 p^{rs} k_{cr} k_{sd}) \cdot h^{o\tilde{n}\tilde{p}} h^{\tilde{m}\tilde{q}} \cdot \right. \\
&\cdot \left(\frac{\alpha_s}{\sqrt{\hbar c}} C_{av}^c (\delta_{\tilde{n}}^{\tilde{a}} [\Phi_{\text{crt}}^k]_{\tilde{m}}^v - [\Phi_{\text{crt}}^k]_{\tilde{n}}^v \delta_{\tilde{m}}^{\tilde{a}}) - f_{\tilde{n}\tilde{m}}^{\tilde{a}} \delta_a^c \right) \cdot \\
&\cdot \left(\frac{\alpha_s}{\sqrt{\hbar c}} C_{ru}^d (\delta_{\tilde{p}}^{\tilde{r}} [\Phi_{\text{crt}}^k]_{\tilde{q}}^u - [\Phi_{\text{crt}}^k]_{\tilde{p}}^u \delta_{\tilde{q}}^{\tilde{r}}) - f_{\tilde{p}\tilde{q}}^{\tilde{r}} \delta_d^u \right) + \\
&+ \frac{2\alpha_s}{\sqrt{\hbar c}} C_{ar}^c (\delta_{\tilde{n}}^{\tilde{a}} \delta_{\tilde{m}}^{\tilde{r}} - \delta_{\tilde{m}}^{\tilde{a}} \delta_{\tilde{n}}^{\tilde{r}}) (p_{cd} + \xi^2 p^{rs} k_{cr} k_{sd}) h^{o\tilde{p}\tilde{n}} h^{o\tilde{q}\tilde{m}} \cdot \\
&\cdot \left(\frac{\alpha_s}{\sqrt{\hbar c}} C_{zx}^b [\Phi_{\text{crt}}^k]_{\tilde{p}}^z [\Phi_{\text{crt}}^k]_{\tilde{q}}^x - [\Phi_{\text{crt}}^k]_{\tilde{z}}^b f_{\tilde{p}\tilde{q}}^{\tilde{z}} - \frac{\sqrt{\hbar c}}{\alpha_s} \mu_i^b f_{\tilde{p}\tilde{q}}^{\tilde{z}} \right) \Big\}, \quad (17.41)
\end{aligned}$$

$$\begin{aligned}
\bar{m}^{(1)2}(\Phi_{\text{crt}}^k)_{\tilde{a} \tilde{r}}^{\tilde{a} \tilde{r}} &= \frac{\zeta \xi}{r^4} \int_M \left\{ \left(\frac{\alpha_s}{\sqrt{\hbar c}} C_{av}^c (\delta_{\tilde{n}}^{\tilde{a}} [\Phi_{\text{crt}}^k]_{\tilde{m}}^v - [\Phi_{\text{crt}}^k]_{\tilde{n}}^v \delta_{\tilde{m}}^{\tilde{a}}) - f_{\tilde{n}\tilde{m}}^{\tilde{a}} \delta_a^c \right) \cdot \right. \\
&\cdot \left(\frac{\alpha_s}{\sqrt{\hbar c}} C_{ru}^d (\delta_{\tilde{p}}^{\tilde{r}} [\Phi_{\text{crt}}^k]_{\tilde{q}}^u - [\Phi_{\text{crt}}^k]_{\tilde{p}}^u \delta_{\tilde{q}}^{\tilde{r}}) - f_{\tilde{p}\tilde{q}}^{\tilde{r}} \delta_d^u \right) \cdot \\
&\cdot h^{o\tilde{n}\tilde{s}} h^{o\tilde{p}\tilde{t}} h^{o\tilde{n}\tilde{q}} p_{cd} p^{ar} (k_{rs} + \xi k_{rp} p^{pq} k_{qs}) k_{\tilde{t}\tilde{s}}^o + \\
&+ \frac{\alpha_s}{\sqrt{\hbar c}} C_{ar}^c (\delta_{\tilde{n}}^{\tilde{a}} \delta_{\tilde{m}}^{\tilde{r}} - \delta_{\tilde{m}}^{\tilde{a}} \delta_{\tilde{n}}^{\tilde{r}}) p^{ar} k_{\tilde{t}\tilde{s}}^o \cdot \\
&\cdot [\ell_{bc} h^{o\tilde{a}\tilde{m}} h^{o\tilde{b}\tilde{s}} h^{o\tilde{n}\tilde{t}} (k_{rs} + \xi k_{rp} p^{pq} k_{qs}) + \\
&+ \ell_{ab} h^{o\tilde{m}\tilde{s}} h^{o\tilde{b}\tilde{t}} h^{o\tilde{n}\tilde{a}} (k_{rc} + \xi k_{rp} p^{pq} k_{qc})] \cdot \\
&\cdot \left(\frac{\alpha_s}{\sqrt{\hbar c}} C_{zx}^b [\Phi_{\text{crt}}^k]_{\tilde{p}}^z [\Phi_{\text{crt}}^k]_{\tilde{q}}^x - [\Phi_{\text{crt}}^k]_{\tilde{z}}^b f_{\tilde{p}\tilde{q}}^{\tilde{z}} - \frac{\sqrt{\hbar c}}{\alpha_s} \mu_i^b f_{\tilde{p}\tilde{q}}^{\tilde{z}} \right) \Big\} \cdot \\
&\cdot \frac{\sqrt{|h^0|}}{V_2} dm(y), \quad (17.42)
\end{aligned}$$

$$\begin{aligned}
\bar{m}^{(2)2}(\Phi_{\text{crt}}^k)_{\tilde{a} \tilde{r}}^{\tilde{a} \tilde{r}} &= - \int_M \frac{\zeta^2}{r^4} \left\{ \frac{\alpha_s}{\sqrt{\hbar c}} C_{av}^c (\delta_{\tilde{n}}^{\tilde{a}} [\Phi_{\text{crt}}^k]_{\tilde{m}}^v - [\Phi_{\text{crt}}^k]_{\tilde{n}}^v \delta_{\tilde{m}}^{\tilde{a}}) - f_{\tilde{n}\tilde{m}}^{\tilde{a}} \delta_a^c \right\} \cdot \\
&\cdot \left(\alpha_s \sqrt{\frac{\hbar}{c}} C_{ru}^d (\delta_{\tilde{p}}^{\tilde{r}} [\Phi_{\text{crt}}^k]_{\tilde{q}}^u - [\Phi_{\text{crt}}^k]_{\tilde{p}}^u \delta_{\tilde{q}}^{\tilde{r}}) - f_{\tilde{p}\tilde{q}}^{\tilde{r}} \delta_r^d \right) \cdot \\
&\cdot \ell_{ac} k_{\tilde{t}\tilde{s}}^o k_{\tilde{d}\tilde{e}}^o \cdot \\
&\cdot \left[(2h^{o\tilde{p}\tilde{q}} h^{o\tilde{m}\tilde{t}} h^{o\tilde{s}\tilde{d}} h^{o\tilde{n}\tilde{c}} + h^{o\tilde{p}\tilde{s}} h^{o\tilde{m}\tilde{t}} h^{o\tilde{q}\tilde{e}} h^{o\tilde{n}\tilde{d}}) k_{+d}^a + \right. \\
&+ (h^{o\tilde{c}\tilde{s}} h^{o\tilde{m}\tilde{t}} h^{o\tilde{q}\tilde{n}} h^{o\tilde{p}\tilde{d}} - h^{o\tilde{d}\tilde{s}} h^{o\tilde{m}\tilde{t}} h^{o\tilde{q}\tilde{n}} h^{o\tilde{p}\tilde{c}} + \\
&+ h^{o\tilde{q}\tilde{s}} h^{o\tilde{m}\tilde{t}} h^{o\tilde{d}\tilde{n}} h^{o\tilde{p}\tilde{c}}) k_{+d}^a k_{-d}^s + \\
&+ \frac{\xi}{2} (h^{o\tilde{q}\tilde{m}} h^{o\tilde{t}\tilde{n}} h^{o\tilde{d}\tilde{p}} h^{o\tilde{e}\tilde{s}} + h^{o\tilde{q}\tilde{m}} h^{o\tilde{t}\tilde{n}} h^{o\tilde{e}\tilde{p}} h^{o\tilde{d}\tilde{s}} +
\end{aligned}$$

$$\begin{aligned}
& -h^{o\tilde{q}\tilde{m}}h^{o\tilde{s}\tilde{n}}h^{o\tilde{d}\tilde{p}}h^{o\tilde{e}\tilde{t}} - h^{o\tilde{q}\tilde{m}}h^{o\tilde{s}\tilde{n}}h^{o\tilde{e}\tilde{p}}h^{o\tilde{d}\tilde{t}})k^a_{+s}k^s_{-d}+ \\
& + \frac{\xi}{2}(h^{o\tilde{q}\tilde{m}}h^{o\tilde{t}\tilde{n}}h^{o\tilde{d}\tilde{p}}h^{o\tilde{e}\tilde{s}} + h^{o\tilde{q}\tilde{m}}h^{o\tilde{t}\tilde{n}}h^{o\tilde{e}\tilde{p}}h^{o\tilde{d}\tilde{s}}+ \\
& + h^{o\tilde{q}\tilde{m}}h^{o\tilde{s}\tilde{n}}h^{o\tilde{d}\tilde{p}}h^{o\tilde{e}\tilde{t}} - h^{o\tilde{q}\tilde{m}}h^{o\tilde{s}\tilde{n}}h^{o\tilde{e}\tilde{p}}h^{o\tilde{d}\tilde{t}})(k^a_{+s}k^s_{-d})p^{pq}k_{qd}+ \\
& - \frac{\xi}{2}(h^{o\tilde{m}\tilde{s}}h^{o\tilde{n}\tilde{t}}h^{o\tilde{p}\tilde{p}}h^{o\tilde{q}\tilde{d}} - h^{o\tilde{m}\tilde{t}}h^{o\tilde{n}\tilde{s}}h^{o\tilde{p}\tilde{e}}h^{o\tilde{q}\tilde{d}}+ \\
& - h^{o\tilde{m}\tilde{s}}h^{o\tilde{n}\tilde{t}}h^{o\tilde{p}\tilde{e}}h^{o\tilde{q}\tilde{e}} + h^{o\tilde{m}\tilde{t}}h^{o\tilde{n}\tilde{s}}h^{o\tilde{p}\tilde{d}}h^{o\tilde{q}\tilde{e}})\delta_d^a \Big] + \\
& + \frac{\alpha_s}{\sqrt{hc}}C_{ar}^c(\delta_{\tilde{n}}^{\tilde{a}}\delta_{\tilde{m}}^{\tilde{r}} - \delta_{\tilde{m}}^{\tilde{a}}\delta_{\tilde{n}}^{\tilde{r}})k^o_{\tilde{t}\tilde{s}}k^o_{\tilde{d}\tilde{e}} \cdot \\
& \cdot \left\{ \left[(2h^{o\tilde{b}\tilde{c}}h^{o\tilde{m}\tilde{t}}h^{o\tilde{s}\tilde{d}}h^{o\tilde{n}\tilde{c}} + h^{o\tilde{b}\tilde{s}}h^{o\tilde{m}\tilde{t}}h^{o\tilde{c}\tilde{e}}h^{o\tilde{n}\tilde{d}})k^a_{+b}+ \right. \right. \\
& + (h^{o\tilde{e}\tilde{s}}h^{o\tilde{m}\tilde{t}}h^{o\tilde{c}\tilde{n}}h^{o\tilde{b}\tilde{d}} - h^{o\tilde{d}\tilde{s}}h^{o\tilde{m}\tilde{t}}h^{o\tilde{c}\tilde{n}}h^{o\tilde{b}\tilde{e}}+ \\
& + h^{o\tilde{c}\tilde{s}}h^{o\tilde{m}\tilde{t}}h^{o\tilde{e}\tilde{n}}h^{o\tilde{b}\tilde{d}} - h^{o\tilde{c}\tilde{s}}h^{o\tilde{m}\tilde{t}}h^{o\tilde{d}\tilde{n}}h^{o\tilde{b}\tilde{e}})k^a_{+r}k^r_{-b}+ \\
& + \frac{\xi}{2}(h^{o\tilde{c}\tilde{m}}h^{o\tilde{t}\tilde{n}}h^{o\tilde{d}\tilde{b}}h^{o\tilde{e}\tilde{s}} + h^{o\tilde{c}\tilde{m}}h^{o\tilde{t}\tilde{n}}h^{o\tilde{e}\tilde{b}}h^{o\tilde{d}\tilde{s}}+ \\
& - h^{o\tilde{c}\tilde{m}}h^{o\tilde{s}\tilde{n}}h^{o\tilde{d}\tilde{b}}h^{o\tilde{e}\tilde{t}} - h^{o\tilde{c}\tilde{m}}h^{o\tilde{s}\tilde{n}}h^{o\tilde{e}\tilde{b}}h^{o\tilde{d}\tilde{t}})k^a_{+r}k^r_{-b}+ \\
& + \frac{\xi}{2}(h^{o\tilde{c}\tilde{m}}h^{o\tilde{t}\tilde{n}}h^{o\tilde{d}\tilde{b}}h^{o\tilde{e}\tilde{s}} + h^{o\tilde{c}\tilde{m}}h^{o\tilde{t}\tilde{n}}h^{o\tilde{e}\tilde{b}}h^{o\tilde{d}\tilde{s}}+ \\
& - h^{o\tilde{c}\tilde{m}}h^{o\tilde{s}\tilde{n}}h^{o\tilde{d}\tilde{b}}h^{o\tilde{e}\tilde{t}} - h^{o\tilde{c}\tilde{m}}h^{o\tilde{s}\tilde{n}}h^{o\tilde{e}\tilde{b}}h^{o\tilde{d}\tilde{t}})(k^a_{+s}k^r_{-p})p^{pq}k_{qd}+ \\
& + \frac{\xi}{2}(h^{o\tilde{c}\tilde{d}}h^{o\tilde{t}\tilde{b}}h^{o\tilde{s}\tilde{m}}h^{o\tilde{e}\tilde{n}} - h^{o\tilde{c}\tilde{d}}h^{o\tilde{s}\tilde{b}}h^{o\tilde{t}\tilde{m}}h^{o\tilde{e}\tilde{n}}+ \\
& + h^{o\tilde{c}\tilde{e}}h^{o\tilde{t}\tilde{b}}h^{o\tilde{s}\tilde{m}}h^{o\tilde{d}\tilde{n}} + h^{o\tilde{c}\tilde{e}}h^{o\tilde{s}\tilde{b}}h^{o\tilde{t}\tilde{m}}h^{o\tilde{d}\tilde{n}})(k^a_{+r}k^s_{-d})p^{pq}k_{qb}+ \\
& - \frac{\xi}{2}(h^{o\tilde{m}\tilde{s}}h^{o\tilde{n}\tilde{t}}h^{o\tilde{b}\tilde{s}}h^{o\tilde{c}\tilde{d}} - h^{o\tilde{m}\tilde{t}}h^{o\tilde{n}\tilde{s}}h^{o\tilde{b}\tilde{e}}h^{o\tilde{c}\tilde{d}}+ \\
& - h^{o\tilde{m}\tilde{s}}h^{o\tilde{n}\tilde{t}}h^{o\tilde{b}\tilde{d}}h^{o\tilde{c}\tilde{e}} - h^{o\tilde{m}\tilde{t}}h^{o\tilde{n}\tilde{s}}h^{o\tilde{b}\tilde{d}}h^{o\tilde{c}\tilde{e}})\delta_d^a \Big] \ell_{ac}+ \\
& - \ell_{ab}((2h^{o\tilde{h}\tilde{e}}h^{o\tilde{c}\tilde{t}}h^{o\tilde{s}\tilde{d}}h^{o\tilde{b}\tilde{m}} + h^{o\tilde{n}\tilde{s}}h^{o\tilde{c}\tilde{t}}h^{o\tilde{m}\tilde{e}}h^{o\tilde{b}\tilde{d}})k^a_{+c}+ \\
& + (h^{o\tilde{e}\tilde{s}}h^{o\tilde{c}\tilde{t}}h^{o\tilde{m}\tilde{b}}h^{o\tilde{n}\tilde{d}} - h^{o\tilde{d}\tilde{s}}h^{o\tilde{c}\tilde{t}}h^{o\tilde{m}\tilde{b}}h^{o\tilde{n}\tilde{e}}+ \\
& + (h^{o\tilde{e}\tilde{s}}h^{o\tilde{c}\tilde{t}}h^{o\tilde{m}\tilde{b}}h^{o\tilde{n}\tilde{d}} - h^{o\tilde{d}\tilde{s}}h^{o\tilde{c}\tilde{t}}h^{o\tilde{m}\tilde{b}}h^{o\tilde{n}\tilde{e}}+ \\
& + h^{o\tilde{m}\tilde{s}}h^{o\tilde{c}\tilde{t}}h^{o\tilde{e}\tilde{b}}h^{o\tilde{n}\tilde{d}} - h^{o\tilde{m}\tilde{s}}h^{o\tilde{c}\tilde{t}}h^{o\tilde{d}\tilde{b}}h^{o\tilde{n}\tilde{e}})k^a_{+r}k^r_{-c}+ \\
& + \frac{\xi}{2}(h^{o\tilde{c}\tilde{m}}h^{o\tilde{t}\tilde{b}}h^{o\tilde{d}\tilde{n}}h^{o\tilde{e}\tilde{b}} + h^{o\tilde{c}\tilde{m}}h^{o\tilde{t}\tilde{b}}h^{o\tilde{d}\tilde{n}}h^{o\tilde{e}\tilde{s}}+ \\
& - h^{o\tilde{c}\tilde{m}}h^{o\tilde{s}\tilde{b}}h^{o\tilde{d}\tilde{n}}h^{o\tilde{e}\tilde{t}} - h^{o\tilde{c}\tilde{m}}h^{o\tilde{s}\tilde{b}}h^{o\tilde{e}\tilde{n}}h^{o\tilde{d}\tilde{t}})k^a_{+r}k^r_{-c}+ \\
& + \frac{\xi}{2}(-h^{o\tilde{m}\tilde{d}}h^{o\tilde{t}\tilde{n}}h^{o\tilde{s}\tilde{c}}h^{o\tilde{e}\tilde{b}} + h^{o\tilde{m}\tilde{d}}h^{o\tilde{s}\tilde{n}}h^{o\tilde{t}\tilde{c}}h^{o\tilde{e}\tilde{b}}+
\end{aligned}$$

$$\begin{aligned}
& -h^{o\tilde{m}\tilde{e}}h^{o\tilde{t}\tilde{n}}h^{o\tilde{s}\tilde{c}}h^{o\tilde{d}\tilde{b}} + h^{o\tilde{m}\tilde{e}}h^{o\tilde{s}\tilde{n}}h^{o\tilde{t}\tilde{c}}h^{o\tilde{d}\tilde{b}})(k_{+s}^ak_{-p}^r)p^{pq}k_{qc}+ \\
& + \frac{\xi}{2}(-h^{o\tilde{c}\tilde{m}}h^{o\tilde{t}\tilde{b}}h^{o\tilde{d}\tilde{n}}h^{o\tilde{e}\tilde{s}} + h^{o\tilde{c}\tilde{m}}h^{o\tilde{t}\tilde{b}}h^{o\tilde{e}\tilde{n}}h^{o\tilde{d}\tilde{s}}+ \\
& + h^{o\tilde{c}\tilde{m}}h^{o\tilde{s}\tilde{b}}h^{o\tilde{d}\tilde{n}}h^{o\tilde{s}\tilde{t}} - h^{o\tilde{c}\tilde{m}}h^{o\tilde{s}\tilde{b}}h^{o\tilde{e}\tilde{n}}h^{o\tilde{d}\tilde{t}})(k_{+r}^ak_{-p}^r)p^{pq}k_{qc}+ \\
& - \frac{\xi}{2}(h^{o\tilde{c}\tilde{s}}h^{o\tilde{b}\tilde{t}}h^{o\tilde{n}\tilde{s}}h^{o\tilde{m}\tilde{d}} - h^{o\tilde{c}\tilde{t}}h^{o\tilde{b}\tilde{s}}h^{o\tilde{n}\tilde{e}}h^{o\tilde{m}\tilde{d}}+ \\
& - h^{o\tilde{c}\tilde{s}}h^{o\tilde{b}\tilde{t}}h^{o\tilde{n}\tilde{d}}h^{o\tilde{m}\tilde{e}} + h^{o\tilde{c}\tilde{t}}h^{o\tilde{b}\tilde{s}}h^{o\tilde{n}\tilde{d}}h^{o\tilde{m}\tilde{e}})\delta_c^d) \\
& \cdot \left(\frac{\alpha_s}{\sqrt{\hbar c}}C_{uv}^b[\Phi_{\text{crt}}^k]_{\tilde{c}}^z[\Phi_{\text{crt}}^k]_{\tilde{b}}^v - [\Phi_{\text{crt}}^{\tilde{k}}]_{\tilde{z}}^bf_{\tilde{c}\tilde{b}}^{\tilde{z}} - \frac{\sqrt{\hbar c}}{\alpha_s}\mu_i^bf_{\tilde{p}\tilde{q}}^{\tilde{i}} \right) \Bigg\} \Bigg\}. \\
& \cdot \frac{\sqrt{|h^0|}}{V_2}dm(y). \tag{17.43}
\end{aligned}$$

After a diagonalization of the matrix $m^2(\Phi_{\text{crt}}^k)_{\tilde{a}\tilde{b}}$ we get a mass spectrum for Higgs' particles. $k = 0$ corresponds to the “true” vacuum case and $k = 1$ to the “false” vacuum case. In both cases i.e. for vector bosons and Higgs' particles this is the mass spectrum up to the second order of expansion with respect to ξ . Next orders of expansion can be obtained in a similar way. The lagrangian involving Higgs' field depends on the scalar field Ψ .

Let us consider the full lagrangian of the theory in the second order of approximation with respect to $h_{[\mu\nu]}$ and $\overline{W}_\mu$. Thus let us take this part of Yang–Mill's and Higgs' lagrangian which depends on $\overline{W}_\mu$ only. We remind to the reader that we suppose in section 12 an identification $\tilde{A}_\mu^F = -\frac{2}{3}\overline{W}_\mu$ and $\tilde{H}_{\mu\nu}^F = -\frac{4}{3}\overline{W}_{[\mu,\nu]}$.

Let us take the Yang–Mill's lagrangian in the zeroth order

$$\begin{aligned}
\mathcal{L}_{YM}^{(0)} &= \frac{2}{9\pi}[\tilde{p}_{FF} + \xi^2\tilde{k}_{Fr}\tilde{p}^{rs}\tilde{k}_{sF}]\overline{W}_{[\beta,\gamma]}\overline{W}_{[\mu,\alpha]}\eta^{\beta\mu}\eta^{\gamma\alpha}+ \\
& + \frac{1}{12\pi}[\tilde{p}_{Fb} + \xi^2\tilde{k}_{Fr}\tilde{p}^{rs}\tilde{k}_{sb}]\overline{W}_{[\beta,\gamma]}H_{[\mu,\alpha]}^b\eta^{\beta\mu}\eta^{\gamma\alpha} \tag{17.44}
\end{aligned}$$

and in the first order:

$$\begin{aligned}
\mathcal{L}_{YM}^{(1)} &= -\frac{4}{9\pi}(\tilde{p}_{FF} + \xi^2\tilde{k}_{Fr}\tilde{p}^{rs}\tilde{k}_{sF})\eta^{\beta\sigma}\eta^{\mu\tau}\eta^{\gamma\alpha}h_{(\tau\sigma)}\overline{W}_{[\beta,\gamma]}\overline{W}_{[\mu,\alpha]}+ \\
& + \frac{2}{3\pi}(\tilde{p}_{Fb} + \xi^2\tilde{k}_{Fr}\tilde{p}^{rs}\tilde{k}_{sb})\eta^{\beta\sigma}\eta^{\mu\tau}\eta^{\gamma\alpha}h_{(\tau\sigma)}\overline{W}_{[\beta,\gamma]}H_{[\mu,\alpha]}^b+ \\
& - \frac{4\xi}{3\pi}(\tilde{p}_{aF} + \xi\tilde{k}_{aF})\tilde{p}^{ar}\tilde{k}_{rs}(\delta^s_F + \xi\tilde{p}^{sq}\tilde{k}_{qF})\eta^{\beta\sigma}\eta^{\mu\tau}\eta^{\gamma\alpha}h_{[\tau\sigma]}\overline{W}_{[\beta,\gamma]}\overline{W}_{[\mu,\alpha]}+ \\
& - \frac{\xi}{3\pi}(\tilde{p}_{aF} + \xi\tilde{k}_{aF})\tilde{p}^{ar}\tilde{k}_{rs}(\delta^s_p + \xi\tilde{p}^{sq}\tilde{k}_{qp})\eta^{\beta\sigma}\eta^{\mu\tau}\eta^{\gamma\alpha}h_{[\tau\sigma]}\overline{W}_{[\beta,\gamma]}\overline{W}_{[\mu,\alpha]}+
\end{aligned}$$

$$-\frac{4\tilde{\xi}}{3\pi}(\tilde{p}_{ab} + \tilde{\xi}\tilde{k}_{ab})\tilde{p}^{ar}\tilde{k}_{rs}(\delta^s_F + \tilde{\xi}\tilde{p}^{sq}\tilde{k}_{sF})\eta^{\beta\sigma}\eta^{\mu\tau}\eta^{\gamma\alpha}h_{[\tau\sigma]}\overline{W}_{[\beta,\gamma]}\overline{W}_{[\mu,\alpha]}^b, \quad (17.45)$$

where $\tilde{\ell}_{ab} = \tilde{p}_{ab} + \tilde{\xi}\tilde{k}_{\tilde{a}\tilde{b}}$ is the nonsymmetric right-invariant tensor on the gauge group G (being a subgroup of H). The field $\overline{W}_\mu$ appears also in $\mathcal{L}_{\text{int}}(\Phi, \tilde{A})$. One gets

$$\begin{aligned} \mathcal{L}^{(2)}(\Phi, \tilde{A}) = & -\frac{4\tilde{\xi}}{3V_2} \int_M \sqrt{|h^0|} dm(y) \cdot \\ & \cdot \{\tilde{p}_{ab}\mu^a{}_F \eta^{\mu\alpha}\eta^{\alpha\beta}h_{\beta\alpha}h^{o\tilde{m}\tilde{a}}h^{o\tilde{n}\tilde{b}}k^o_{\tilde{b}\tilde{a}}\overline{W}_{[\mu,\nu]}H^b_{\tilde{m}\tilde{n}}\}. \end{aligned} \quad (17.46)$$

Moreover, the field $\tilde{A}^F_\mu = -\frac{2}{3}\overline{W}_\mu$ couples the Higgs field via the gauge derivative. Thus we should consider $\mathcal{L}_{\text{kin}}(\overset{\text{gauge}}{\nabla}\Phi)$ up to the second order of $\overline{W}_\mu$. Moreover, we do not consider in this case the full lagrangian $\mathcal{L}_{\text{kin}}(\overset{\text{gauge}}{\nabla}\Phi)$ because according to our assumption the $\tilde{A}^F_\mu$ field becomes massive after a symmetry breaking from G to G_0 and it is enough to consider a mass term for this field in the lagrangian

$$\frac{1}{2}\eta^{\mu\nu}m_F^2\tilde{A}^F_\mu\tilde{A}^F_\nu\tilde{p}_{FF} = \frac{2}{9}m_F^2\tilde{p}_{FF}\eta^{\mu\nu}\overline{W}_\mu\overline{W}_\nu, \quad (17.47)$$

where m_F is the mass of boson corresponding to the broken $\tilde{A}^F_\mu$ field (the $\mathbf{U}(1)_F$ subgroup G , $\mathbf{U}(1)_F \not\subset G_0$). The remaining part of the full lagrangian of the theory containing $\overline{W}_\mu$ is the term

$$\frac{2}{3}g^{[\mu\nu]}\overline{W}_{[\mu,\nu]}. \quad (17.48)$$

Taking this term up to the second order with respect to $h_{(\mu\nu)}$, $h_{[\mu\nu]}$ and $\overline{W}_\mu$ one gets:

$$-\frac{2}{3}\eta^{[\mu|\alpha|}\eta^{\nu]}h_{\beta\alpha}\overline{W}_{[\mu,\nu]}. \quad (17.49)$$

We should add of course a quadratic approximation of $R_4(\tilde{F})$ in N.G.T. to have full lagrangian containing $\overline{W}_\mu$. This is known in N.G.T. The Nonsymmetric–Nonabelian Jordan–Theory (Kaluza–Klein) Theory with Higgs’ mechanism gives us several additional couplings between $h_{[\mu\nu]}$, $h_{(\mu\nu)}$ and $\overline{W}_\mu$, which are absent in pure N.G.T. It seems that they can help in solving some serious problems in N.G.T. raised recently by Damour (see Ref. [121]).

In order to have a more clear situation we take only this terms which does not depend on the gauge field (broken or unbroken) and which are at least quadratic in $\overline{W}_\mu$, $h_{[\mu\nu]}$, $h_{(\mu\nu)}$. One gets:

$$\begin{aligned} \mathcal{L}_{\text{quadratic}}(\overline{W}) = & \frac{2}{9\pi}(\tilde{p}_{FF} + \tilde{\xi}^2\tilde{k}_{F\tau}\tilde{p}^{rs}\tilde{k}_{sF})\eta^{\beta\mu}\eta^{\gamma\alpha}\overline{W}_{[\beta,\gamma]}\overline{W}_{[\mu,\alpha]} + \\ & -\frac{2}{9}\tilde{p}_{FF}m_F^2\eta^{\mu\nu}\overline{W}_\mu\overline{W}_\nu - \frac{2}{3}\eta^{\nu\beta}\eta^{\mu\alpha}h_{[\beta,\alpha]}\overline{W}_{[\mu,\nu]}. \end{aligned} \quad (17.50)$$

Thus we get the lagrangian which is a Proca-field lagrangian interacting with $h_{[\beta\alpha]}$.

Let us consider linearization of Eqs. (12.44 ÷ 49) with respect to $h_{(\mu\nu)}$, $h_{[\mu\nu]}$ and $\overline{W}_\mu$ and let us neglect all Yang–Mill’s field and Higgs’ field except $\overline{W}_\mu$ field ($\tilde{A}^F_\mu$). One easily gets

$$\begin{aligned} \tilde{R}_{\mu\rho} + \frac{1}{4}F^\lambda_{\mu\rho,\lambda} + \left(\alpha + \frac{5}{6}\right)h_{[\mu\lambda],\rho}{}^\lambda + \\ - \left(\alpha + \frac{1}{2}\right)h_{[\rho\xi],\mu}{}^\lambda + \frac{1}{2}h_{[\rho\mu],\lambda}{}^\lambda + \frac{2}{3}(1 + 4\alpha)\overline{W}_{[\mu\rho]} = ah_{\mu\rho}. \end{aligned} \quad (17.51)$$

Notice that we omit the energy—momentum tensor for $\overline{W}_\mu$, because it is of the second order with respect to $\overline{W}_\mu$, $ah_{\mu\rho}$ is a cosmological term with a being a constant, $\tilde{R}_{\mu\rho}$ is the usual Ricci tensor for $h_{(\mu\nu)}$ in the linear approximation

$$F_{\lambda\mu\rho} = 3h_{[[\lambda\mu],\rho]}. \quad (17.52)$$

In the formula (17.51) we take for a full Ricci tensor the following sum.

$$R_{\mu\nu}(\overline{W}) = R^\lambda_{\mu\nu\lambda} + \alpha R^\lambda_{\chi\mu\nu}. \quad (17.53)$$

In the work we are using $\alpha = \frac{1}{2}$.

The most interesting case is the interaction of the skewon field $h_{[\mu\nu]}$ with $\overline{W}_\mu$.

The lagrangian for the $\overline{W}_\mu$ field looks as follows

$$L(\overline{W}) = -\frac{A^2}{8\pi}\overline{W}_{[\mu,\nu]}\overline{W}^{[\mu,\nu]} + B^2\overline{W}_\mu\overline{W}^\mu + \frac{2}{3}(1 + 4\alpha)h^{[\mu\nu]}\overline{W}_{[\mu,\nu]}, \quad (17.54)$$

where A and B are constants and

$$h^{[\mu\nu]} = \eta^{\mu\alpha}\eta^{\nu\beta}h_{[\alpha\beta]}, \quad (17.55)$$

$\eta_{\mu\nu}$ is a Minkowski’s tensor.

The constants A and B depends on the details of the theory presented before (i.e. groups, constants ξ , μ , ζ).

Now we redefine the field $\overline{W}_\mu$ introducing a new field V_μ

$$V_\mu = \frac{1}{2}A\overline{W}_\mu. \quad (17.56)$$

One gets:

$$L(V) = -\frac{1}{8\pi}V_{\mu\nu}V^{\mu\nu} + \frac{m^2}{4\pi}V_\mu V^\mu - \frac{2}{3}g_V h^{[\mu\nu]}V_{\mu\nu}, \quad (17.57)$$

where

$$V_{\mu\nu} = \partial_\mu V_\nu - \partial_\nu V_\mu \quad (17.58)$$

and m_V is a mass of the Proca field V_μ , and g_V is a coupling constant, $g_V > 0$. The equation for the field V_μ is as follows:

$$(-\eta^{\mu\lambda}\square + \partial^\mu\partial^\lambda - \eta^{\mu\lambda}m_V^2)V_\lambda = \frac{8\pi}{3}g_V\partial_\nu h^{[\nu\mu]}. \quad (17.59)$$

The equation can be rewritten in the following form:

$$\partial_\nu V^{\nu\mu} + m_V^2 V^\mu = -\frac{8\pi}{3}g_V\partial_\nu h^{[\nu\mu]}. \quad (17.60)$$

Now we come back to Eq. (17.51) and write them for the symmetric and antisymmetric part eliminating $\bar{W}_\mu$ field. One gets:

$$\begin{aligned} \tilde{R}_{\mu\rho} + 2\left(\alpha + \frac{5}{6}\right)(h_{[\mu\lambda],\rho}{}^\lambda + h_{[\rho\lambda],\mu}{}^\lambda) + \\ -\frac{1}{2}\left(\alpha + \frac{1}{2}\right)(h_{[\rho\lambda],\mu}{}^\lambda + h_{[\mu\lambda],\rho}{}^\lambda) = ah_{(\mu\rho)}, \end{aligned} \quad (17.61)$$

$$\begin{aligned} F_{[\mu\rho,|\lambda|,\gamma]}^\lambda + 2\left(\alpha + \frac{5}{6}\right)(h_{[[\mu|\lambda|],\rho}{}^\lambda, \gamma] + h_{[[\rho|\lambda|],\mu}{}^\lambda, \gamma]) + \\ -2\left(\alpha + \frac{1}{2}\right)(h_{[[\rho|\lambda|],\mu}{}^{|\lambda|}, \gamma] - h_{[[\mu|\lambda|],\rho}{}^{|\lambda|}, \gamma]) + \frac{1}{2}h_{[[\rho\mu],|\lambda|,|\gamma]} = \frac{1}{3}aF_{\mu\rho\gamma}. \end{aligned} \quad (17.62)$$

The Eq. (17.61) describes linear G.R.T. with skewon sources and Eq. (17.62) is the equation for skewon field which couples to the V_μ field in Eq. (17.60). For $\tilde{R}_{\mu\rho}$ we have a simple formula

$$\tilde{R}_{\mu\rho} = -\square\bar{h}_{\mu\rho}, \quad (17.63)$$

where $\bar{h}_{\mu\rho} = h_{(\mu\rho)} - \frac{1}{2}\eta^{\alpha\beta}h_{(\alpha\beta)}\eta_{\mu\rho}$, and $\bar{h}^{\mu\nu}{}_{,\nu} = 0$.

Thus the Eq. (17.61) takes the form

$$\begin{aligned} -\square\bar{h}_{\mu\rho} + \frac{1}{2}\left(\alpha + \frac{5}{6}\right)(h_{[\mu\lambda],\rho}{}^\lambda + h_{[\rho\lambda],\mu}{}^\lambda) + \\ -\frac{1}{2}\left(\alpha + \frac{1}{2}\right)(h_{[\rho\lambda],\mu}{}^\lambda + h_{[\mu\lambda],\rho}{}^\lambda) = a\bar{h}_{\mu\rho} - \frac{a}{2}\eta^{\alpha\beta}\bar{h}_{(\alpha\beta)}\eta_{\mu\rho}, \end{aligned} \quad (17.64)$$

where $\bar{h}^{\mu\rho}{}_{,\rho} = 0$, $\bar{h}_{\mu\rho} = \bar{h}_{\rho\mu}$.

Moreover, it seems that we should consider also some nonlinear terms connected to V_μ as a source of gravitational field in (17.64). It seems also that we should include an electromagnetic field, which remains massless after symmetry breaking and the scalar field Ψ . In this way we consider lagrangians and energy—momentum tensors for V_μ , A_μ and Ψ .

In the lowest order the lagrangians for electromagnetic and scalar fields take forms

$$\mathcal{L}_{\text{em}} = -\frac{1}{8\pi} F_{\mu\nu} F^{\mu\nu}, \quad (17.65)$$

$$\mathcal{L}_{\text{scal}} = \frac{1}{8\pi} \left(\overline{M} \eta^{\mu\nu} \Psi_{,\mu} \Psi_{,\nu} + \frac{m^2}{2} \Psi^2 \right), \quad (17.66)$$

where $\overline{M} = (\ell^{[dc]} \ell_{[dc]} - 3n(n-1))$.

According to our considerations the scalar field Ψ obtains a non-zero mass. Moreover, we should redefine the field Ψ in such a way that

$$\mathcal{L}_{\text{scal}} = \frac{1}{8\pi} \eta^{\mu\nu} \varphi_{,\mu} \varphi_{,\nu} + \frac{m_\varphi^2}{16\pi} \varphi^2, \quad (17.67)$$

where m_φ is a mass of the field φ .

It is easy to see that

$$\varphi = \overline{M} \Psi. \quad (17.68)$$

According to the Eq. (12.44–50) we really need the quantity

$$\overline{T}_{\mu\rho} = T_{\mu\rho} - \frac{1}{2} \eta_{\mu\rho} (\eta^{\alpha\beta} T_{\alpha\beta}). \quad (17.69)$$

One easily finds for V_μ and electromagnetic fields

$$\overline{T}_{\alpha\rho} = \frac{1}{4\pi} \left[\eta^{\tau\mu} V_{\mu\alpha} V_{\tau\rho} - \frac{1}{4} \eta_{\mu\rho} (V_{\mu\beta} V^{\mu\beta}) + m_V^2 V_\alpha V_\rho \right], \quad (17.70)$$

$$\overline{T}_{\alpha\rho}^{\text{em}} = \frac{1}{4\pi} \left(\eta^{\tau\mu} F_{\mu\alpha} F_{\tau\rho} - \frac{1}{4} \eta_{\alpha\rho} (F_{\mu\nu} F^{\mu\nu}) \right). \quad (17.71)$$

For scalar field φ we have

$$\overline{T}_{\alpha\rho}^{\text{scal}} = \frac{1}{4\pi} \left(\varphi_{,\alpha} \varphi_{,\rho} - \frac{1}{4} \eta_{\alpha\rho} (\varphi_{,\mu} \varphi^{,\mu}) + \frac{m_\varphi^2}{4} \eta_{\alpha\rho} \varphi^2 \right). \quad (17.72)$$

We should add of course to the set of equations of fields, equations for electromagnetic and scalar fields. Thus one gets:

$$\begin{aligned} & -\square \overline{h}_{\mu\rho} + \frac{1}{2} \left(\alpha + \frac{5}{6} \right) (h_{[\mu\lambda],\rho}{}^{,\lambda} + h_{[\rho\lambda],\mu}{}^{,\lambda}) - \frac{1}{2} \left(\alpha + \frac{1}{2} \right) (h_{[\rho\lambda],}{}^{,\lambda}{}_{,\mu} + h_{[\mu\lambda],}{}^{,\lambda}{}_{,\rho}) = \\ & = a \overline{h}_{\mu\rho} - \frac{a}{2} \eta^{\alpha\beta} \overline{h}_{(\alpha\beta)} \eta_{\mu\rho} + \frac{\varkappa}{4\pi} \left(\eta^{\tau\beta} (F_{\beta\mu} F_{\tau\rho} + V_{\beta\mu} V_{\tau\rho}) + \varphi_{,\mu} \varphi_{,\rho} - \frac{1}{4} \eta_{\mu\rho} (V_{\alpha\beta} V^{\alpha\beta} + \right. \end{aligned}$$

$$+F_{\alpha\beta}F^{\alpha\beta} + \varphi_{,\alpha}\varphi^{,\alpha} - m_\varphi^2\varphi^2) + m_V^2 V_\mu V_\rho \Big), \quad (17.73)$$

$$\partial_\mu F^{\mu\nu} = 0, \quad F_{[\rho\lambda,\nu]} = 0, \quad (17.74)$$

$$(\square + m_\varphi^2)\varphi = 0, \quad (17.75)$$

$$\partial_\nu V^{\nu\mu} + m_V^2 V^\mu = -\frac{8\pi}{3}g_V\partial_\nu h^{[\nu\mu]}, \quad V_{[\mu\nu,\lambda]} = 0, \quad (17.76)$$

$$\begin{aligned} F_{[\mu\rho,|\lambda|,\gamma]}^\lambda + 2\left(\alpha + \frac{5}{6}\right)(h_{[[\mu|\lambda|],\rho,|\lambda|],\gamma} + h_{[[\rho|\lambda|],\mu,|\lambda|],\gamma}) + \\ -2\left(\alpha + \frac{1}{2}\right)(h_{[[\rho|\lambda|],|\lambda|,\mu,\gamma]} - h_{[[\mu|\lambda|],|\lambda|,\rho,\gamma]}) + \frac{1}{2}h_{[[\rho\mu],|\lambda|,|\lambda|,\gamma]} = \frac{1}{3}aF_{\mu\rho\gamma}, \end{aligned} \quad (17.77)$$

where

$$F_{\mu\rho\gamma} = 3h_{[[\mu\rho],\gamma]}, \quad (17.78)$$

$$\bar{h}_{\mu\rho} = h_{(\mu\rho)} - \frac{1}{2}\eta^{\alpha\beta}h_{(\alpha\beta)}\eta_{\mu\rho}, \quad (17.79)$$

$$\bar{h} = \eta^{\mu\rho}\bar{h}_{\mu\rho} \quad (17.80)$$

and

$$\bar{h}^{\mu\nu}{}_{,\nu} = 0 \quad (17.81)$$

and $\varkappa = \frac{8\pi G_N}{c^4}$.

The Eqs. (17.73 ÷ 17.81) are full set of equations for coupled fields in our approximation.

Let us notice that in Eq. (17.77) we have on the right-hand side a term $\frac{1}{3}aF_{\mu\rho\gamma}$. This term induces a mass term for the field $h_{[\mu\nu]}$. In this way the skewon field $h_{[\mu\nu]}$ is massive (due to cosmological term) and some problems raised in Ref. [121] are not applicable.

Let us recall that the fields V_μ , φ are also massive. $h_{[\mu\nu]}$, V_μ , φ are additional fields which are connected to gravitational interaction (except $h_{(\mu\nu)}$ -field). They are massive and in linear approximation they are of short range (having a Yukawa-type behaviour). Due to this we have no problems with gravitational radiation from closed binary systems. Only the famous quadrupole radiation formula enters the equation for a secular changing of an orbital period. This is exactly the same as in G.R.T. ([122]).

Recently J. W. Moffat has proposed a reformulation of N.G.T. ([123]). It seems that many of additional phenomenological terms in his gravitational lagrangian can be obtained from our theory. In particular in the approximation presented here they are coming from an effective interactions with massive multiplet of Higgs fields. In our theory Higgs' field is a part of multidimensional gauge field in nonsymmetric Kaluza–Klein (Jordan–Thiry) theory. In some sense this is a little similar to Connes Noncommutative Geometry ([124]). Moreover in order to be closer to this approach we should

extend our formalism to include supersymmetry and supergravity in a nonsymmetric version of our theory.

The interesting point in our approach is to find a quantum version of this classical field theory. It simply means how to quantize the theory? It seems that the best way to achieve it is to use Ashtekar approach of canonical quantization ([125]). In his approach applied to gravity there is a possibility to extend the formalism to include gauge fields and Higgs' fields. In such a way we can work with a group $SU(2) \otimes G(SL(2, \mathbb{C}) \otimes G)$ in the place of $SU(2)(SL(2, \mathbb{C}))$.

Let us notice that there is a different possibility to quantize our theory (even in a perturbative regime). This possibility is to consider the theory as an effective one to some energy scale. In this way our Lagrangian is an "effective Lagrangian". It describes low energy interactions and we are using only those degrees of freedom and interactions appropriate for those energies. Whether or not is this theory fundamental, at low energy it can be described by an effective theory. The lowest energy Lagrangian can be used to determine the propagators and low energy vertices, and the rest can be treated as perturbations. When we quantize the theory (introducing propagators for graviton, skewon, photon, gluons, higgses, scalarons, intermediate bosons and vertices for them) and calculate loops, the usual ultraviolet divergences can be absorbed into definitions of renormalized couplings in the most general Lagrangian.

Those effects can be left-over in the amplitudes for long distance propagation (low energy). Roughly speaking we can consider a full Lagrangian of the theory as a sum of local Lagrangians (the most general local effective Lagrangian) absorbing infinities into all couplings' constants. Afterwards we use an energy expansion and order the most general Lagrangian in powers of the low energy scale divided by the high energy scale. If we want to quantize a full theory with a tower of scalar fields, we can use those massive fields as regulator fields in some kind of Pauli-Villars' renormalization scheme. In this way we can get some interesting results similar to J. Donoghue's approach in quantization of General Relativity (see Ref. [126]). In some sense J. Donoghue's approach is an attempt to renormalize nonrenormalizable theory. In both approaches to quantization of this unified theory we can use generalized functions of J. F. Colombeau (see Ref. [127]).

The interesting problem to consider in our theory is to find a reason to existence of a cosmological constant ($\lambda_{c0} = \overline{\Lambda}$, see Eq. (14.296)). A natural assumption is to put both cosmological terms to zero. This is possible by taking special values of μ and ζ such that

$$\tilde{R}(\tilde{\Gamma}(\mu_0)) = \tilde{P}(\zeta_0) = 0. \quad (17.82)$$

In this way we control a value of a cosmological constant. In order to get a specific value of λ_{c0} we should take some values of μ and ζ different from μ_0 and ζ_0 . For μ and ζ enter some observables connected to elementary interactions and other cosmological predictions (see sections 18 and 19), we should be very careful. In this way we have to do with some kind of fine tuning of μ and ζ . It seems moreover that the non-zero

cosmological terms are connected to a breaking of some kind of invariance in the theory. Probably to a conformal invariance.

In order to examine this conjecture let us linearize full field equations of our theory (Eqs (9.3)÷(9.26)). It is easy to see that those (linearized) equations are conformally invariant if both cosmological terms are zero and $r \rightarrow \infty$. We linearize $g_{\mu\nu}$ around $\eta_{\mu\nu}$ and Ψ around Ψ_0 . Equations for Yang-Mills field are not linearized. In a general case in order to get a scale invariance we should have a condition

$$T^{\text{tot}}_{\alpha\beta} g^{\alpha\beta} = 0 \quad (17.83)$$

where $T^{\text{tot}}_{\alpha\beta}$ is a total energy momentum tensor (including cosmological terms). For this we should develop a theory to include fermions in order to get more conclusive results. Thus we should extend this approach to supermanifolds with anticommuting coordinates and to superfibrebundles with super Lie groups (algebras).

18. On some Cosmological Consequences of the Nonsymmetric Kaluza–Klein (Jordan–Thiry) Theory

We consider some consequences of the Nonsymmetric Kaluza–Klein (Jordan–Thiry) Theory. We calculate primordial fluctuation spectrum functions, spectral indices, and first derivatives of spectral indices.

Let us consider Eq. (14.341).

$$\Phi(t) = \frac{(5+3a)\sqrt{5+4a}}{8\alpha_s(A_1 \operatorname{ns}^2(u) + A_2)} + \frac{(1+\sqrt{5+4a})}{2\alpha_s} \quad (18.1)$$

where

$$u = \sqrt[4]{\frac{Ae^{n\Psi_1} (5Ae^{n\Psi_1} - 3\alpha_s^2 r)}{r^2}} \cdot \sqrt{\sin \frac{\varphi + \pi}{3}} (t - t_0) \quad (18.2)$$

$$A_1 = \sqrt{\frac{5}{3} - a^2} \sin \frac{\varphi + \pi}{3} \quad (18.3)$$

$$A_2 = \sqrt{\frac{5}{3} - a^2} \cos \frac{\varphi + 4\pi}{3} + \frac{1}{12}(5+6a) \quad (18.4)$$

and

$$\cos \varphi = -\frac{\sqrt{A}e^{\frac{n}{2}\Psi_1} (9\alpha_s^2 Br^2 + 40Ae^{n\Psi_1})}{12(5Ae^{n\Psi_1} - 3\alpha_s^2 Br^2)^{3/2}} \quad (18.5)$$

$$0 < a = \sqrt{\frac{\alpha_s^2 Br^2}{Ae^{n\Psi_1}}} < 0.3115, \quad (18.6)$$

ns is the Jacobi elliptic function for the modulus

$$m^2 = \frac{\sin\left(\frac{\varphi}{3}\right)}{\sin\left(\frac{\varphi+2\pi}{3}\right)}, \quad (18.7)$$

B is an integration constant, r is a radius of manifold of vacuum states (a scale of energy), α_s a coupling constant. A is given by the equation (14.321) and is fixed by the details of the theory. Ψ_1 is a critical value of a scalar field Ψ . n is a dimension of the group H . Eq. (18.1) gives a solution to the equation of an evolution of the Higgs field under some special assumptions (in the cosmological background). Due to this solution we are able to calculate $P_R(K)$ —a spectrum of primordial fluctuations in the Universe.

According to the Eq. (14.186)

$$P_R(K) = \left(\frac{H_0}{2\pi} \right)^2 \sum_{\vec{m}, d} \left(\frac{H_0}{\frac{d}{dt}(\Phi_{\vec{m}}^d)} \right)^2 \Big|_{t=t^*}. \quad (18.8)$$

Using Eq. (18.1) and the assumption

$$\Phi_{\vec{m}}^d(t) = \delta_{\vec{m}}^d \Phi(t) \quad (18.9)$$

one gets for $t = t^* = \frac{1}{H_0} \ln \frac{K}{R_0 H_0}$

$$P_R(K) = P_0 \frac{(A_1 + A_2 \operatorname{sn}^2(v))^2 \operatorname{sc}^2(v)}{\operatorname{dn}^2(v)} \quad (18.10)$$

where H_0 is a Hubble constant

$$P_0 = \frac{12H_0^4 \alpha_s^2 n_1 A^{3/2} e^{(3/2)n\Psi_1} \cdot \sin^{-3} \left(\frac{\varphi + \pi}{3} \right) \cdot \pi^{-2}}{\left(5\sqrt{Ae^{n\Psi_1}} + 3\alpha_s \sqrt{Br^2} \right) \left(5A\sqrt{Ae^{n\Psi_1}} + 4\alpha_s \sqrt{Br^2} \right) (5Ae^{n\Psi_1} - 3\alpha_s^2 Br^2)} \quad (18.11)$$

$$v = (\ln K - t_0 H_0 - \ln(H_0 R_0)) \frac{1}{H_0} \sqrt[4]{\frac{Ae^{n\Psi_1} (5Ae^{n\Psi_1} - 3\alpha_s^2 Br^2)}{r^4}} \sqrt{\sin \frac{\varphi + \pi}{3}}, \quad (18.12)$$

n_1 means a dimension of the manifold $M = G/G_0$ (a vacuum states manifold), $K^2 = \vec{K}^2$ means as usual the square of the length of a fluctuations wave vector. We calculate a spectral index of a power spectrum

$$n_s(K) - 1 = \frac{d \ln P_R(K)}{d \ln K}, \quad (18.13)$$

$$n_s(K) - 1 = 2C \left(\frac{\operatorname{dn}(v)}{\operatorname{cn}(v) \operatorname{sn}(v)} - m^2 \frac{\operatorname{sn}(v) \operatorname{cn}(v)}{\operatorname{dn}(v)} + \frac{2A_2 \operatorname{sn}(v) \operatorname{cn}(v) \operatorname{dn}(v)}{A_1 + A_2 \operatorname{sn}^2(v)} \right) \quad (18.14)$$

and an important parameter

$$\begin{aligned} \frac{dn_s(K)}{d \ln K} = 2C^2 & \left[(\operatorname{sn}^2(v) \operatorname{dn}(v) - \operatorname{cn}^2(v) \operatorname{dn}(v) - m^2 \operatorname{sn}(v) \operatorname{cn}(v)) \right. \\ & \times \left(\frac{1}{\operatorname{cn}(v) \operatorname{sn}(v)} + \frac{m^2}{\operatorname{dn}(v)} \right) \\ & + 2A_2 \frac{\operatorname{cn}^2(v) \operatorname{dn}^2(v) - \operatorname{sn}^2(v) \operatorname{dn}^2(v) - m^2 \operatorname{sn}^2(v) \operatorname{cn}^2(v)}{A_1 + A_2 \operatorname{sn}^2(v)} \\ & \left. + 4A_2^2 \frac{\operatorname{sn}^2(v) \operatorname{cn}^2(v) \operatorname{dn}^2(v)}{(A_1 + A_2 \operatorname{sn}^2(v))^2} \right] \end{aligned} \quad (18.15)$$

where v is given by the formula (18.12) and all the Jacobi elliptic functions in the formulae (18.10), (18.14–15) have the same modulus

$$m^2 = \frac{\sin \frac{\varphi}{3}}{\sin \frac{\varphi+2\pi}{3}},$$

$$C = \frac{1}{H_0} \sqrt[4]{\frac{Ae^{n\Psi_1} (5Ae^{n\Psi_1} - 3\alpha_s^2 Br^2)}{r^4}} \sqrt{\sin \frac{\varphi + \pi}{3}}. \quad (18.16)$$

In this theory we have some arbitrary parameters which can be translated to parameters A_1 , A_2 , C and m . Thus we can consider Eqs (18.10), (18.14–15) as parametrized by P_0 , A_1 , A_2 , C and m . Simultaneously v can be rewritten as

$$v = C (\ln K - K'_0) \quad (18.17)$$

where K'_0 is an arbitrary parameter. Let us remind that in our theory we have arbitrary parameters: r , ζ , ξ , α_s (they could be fixed by some data from elementary particle physics, however, they are free as some parameters of the theory) and integration constants B and t_0 (R_0 also in some sense). The interesting idea for future research consists in applying CMBFAST or CMBEASY software [128] to calculate all the cosmological CMB characteristics with mentioned parameters considered as free. Afterwards we should fit them to existing cosmological data [104] as good as possible in order to find some constraints for parameters from our theory.

Let us consider the function $P_R(K)$ from Eq. (18.10). It is easy to see that the function

$$f(v) = \frac{(A_1 + A_2 \operatorname{sn}^2(v))^2 \operatorname{sc}^2(v)}{\operatorname{dn}^2(v)} \quad (18.18)$$

is a periodic function

$$f(v) = f(v + 2K(m^2)l), \quad l = 0, \pm 1, \pm 2, \dots, \quad (18.19)$$

$$K(m^2) = \int_0^{\pi/2} \frac{d\theta}{\sqrt{1 - m^2 \sin^2 \theta}}. \quad (18.20)$$

Thus we get

$$P_R(K) = P_R(Ke^{l\eta}), \quad (18.21)$$

$$\eta = \frac{2K(m^2)H_0 \sqrt[4]{\frac{r^4}{Ae^{n\Psi_1}(5Ae^{n\Psi_1} - 3\alpha_s^2 Br^2)}}}{\sqrt{\sin \frac{\varphi + \pi}{3}}} \quad (18.22)$$

or

$$\eta = \frac{2K(m^2)}{C}. \quad (18.23)$$

Thus it is enough to consider $P_R(K)$ in the interval

$$e^{-\eta/2} \leq K \leq e^{+\eta/2} \quad (18.24)$$

(a little similar to the first Brillouin zone). In order to have an interesting cosmological region we should have

$$\eta \simeq 20 \quad (18.25)$$

which gives us a constraint

$$\frac{2K(m^2)}{C} = \frac{2K(m^2)H_0}{\sqrt{\sin \frac{\varphi+\pi}{3}}} \sqrt[4]{\frac{r^4}{Ae^{n\Psi_1} (5Ae^{n\Psi_1} - 3\alpha_s^2 Br^2)}} \approx 10. \quad (18.26)$$

Moreover we should consider a region with $n_s(K) \cong 1$. This can be satisfied if

$$\begin{aligned} \operatorname{dn}^2(v) (A_1 + A_2 \operatorname{sn}^2(v)) - m^2 \operatorname{sn}^2(v) \operatorname{cn}^2(v) (A_1 + A_2 \operatorname{sn}^2(v)) \\ + 2A_2 \operatorname{sn}^2(v) \operatorname{cn}^2(v) \operatorname{dn}^2(v) = 0 \end{aligned} \quad (18.27)$$

with the condition

$$\operatorname{sn}(v) \operatorname{cn}(v) \operatorname{dn}(v) (A_1 + A_2 \operatorname{sn}^2(v)) \neq 0. \quad (18.28)$$

Using some simple relations among Jacobi elliptic functions and introducing a new variable

$$x = \operatorname{sn}^2(v) \quad (18.29)$$

one gets a cubic equation

$$x^3 + \frac{A_1 m^2 - 4A_2 m^2 - 2A_2}{3A_2 m^2} x^2 + \frac{3A_2 - 2A_1 m^2}{3A_2 m^2} x + \frac{A_1}{3A_2 m^2} = 0 \quad (18.30)$$

or

$$x^3 + \frac{\rho m^2 - 4m^2 - 2}{3m^2} x^2 + \frac{3 - 2\rho m^2}{3m^2} x + \frac{\rho}{3m^2} = 0, \quad (18.31)$$

where

$$\rho = \frac{A_1}{A_2} = \frac{\sqrt{\frac{5}{3} - a^2} \sin \frac{\varphi+\pi}{3}}{\sqrt{\frac{5}{3} - a^2} \cos \frac{\varphi+4\pi}{3} + \frac{5+6a}{12}}. \quad (18.32)$$

This equation can be reduced to

$$y^3 + py + q = 0 \quad (18.33)$$

where

$$p = \frac{-m^4(\rho^2 + 10\rho + 16) + m^2(4\rho + 11) - 4}{27m^4} \quad (18.34)$$

$$q = \frac{2m^6(\rho^3 + 15\rho^2 + 92\rho - 64) + 6m^4(49 + 25\rho - 2\rho^2) + 6m^2(43\rho - 64) - 16}{27^2m^6} \quad (18.35)$$

$$x = y - \left(\frac{\rho m^2 - 4m^2 - 2}{9m^2} \right). \quad (18.36)$$

The discriminant of Eq. (18.33) can be written as

$$D = \frac{q^2}{4} + \frac{p^3}{27} \quad (18.37)$$

and one gets after some tedious algebra

$$\begin{aligned} D(\rho, m) = \frac{1}{27^4 m^{10}} \bigg[& -m^{10} (387\rho^4 + 7376\rho^3 + 29456\rho^2 + 114944\rho + 258048) \\ & + m^8 (243\rho^4 + 6660\rho^3 + 19062\rho^2 + 15000\rho + 116352) \\ & + m^6 (246\rho^4 + 2292\rho^3 - 4716\rho^2 + 3299\rho + 2473) \\ & + m^4 (144\rho^3 + 1512\rho^2 + 10476\rho + 19251) \\ & + m^2 (-1548\rho^3 + 3584\rho^2 + 35190\rho - 26460) \\ & + 48(75 - 39\rho) \bigg]. \end{aligned} \quad (18.38)$$

We use the Cardano formulae to find a root x_0 . Afterwards we get

$$v_{\pm} = \pm \int_0^{\arcsin \sqrt{x_0}} \frac{d\theta}{\sqrt{1 - m^2 \sin^2 \theta}}. \quad (18.39)$$

Thus we need $0 < x_0 < 1$.

It is easy to see that every

$$v_l = v_{\pm} + 2K(m^2)l, \quad l = 0, \pm 1, \pm 2, \dots, \quad (18.40)$$

satisfies Eq. (18.27).

If $D(m, \rho) > 0$, we have only one real root

$$x_0 = \sqrt[3]{-\frac{q}{2} + \sqrt{D}} + \sqrt[3]{-\frac{q}{2} - \sqrt{D}} - \left(\frac{\rho m^2 - 4m^2 - 2}{9m^2} \right) \quad (18.41)$$

where q is given by Eq. (18.35) and D by Eq. (18.38). Moreover we need $0 < x_0 < 1$.

In order to find a criterion let us transform Eq. (18.31) via

$$x = \frac{1 + w}{1 - w}. \quad (18.42)$$

This transformation maps a unit circle on a complex plane $|x| < 1$ into $\text{Re } w < 0$. One gets

$$w^3 + \frac{13m^2 + \rho m^2 - 1 + 3\rho}{7m^2 - 3\rho m^2 + 5 - \rho} w^2 + \frac{5m^2 + 3\rho m^2 - 5 - 3\rho}{7m^2 - 3\rho m^2 + 5 - \rho} w + \frac{\rho + 1 - m^2 - \rho m^2}{7m^2 - 3\rho m^2 + 5 - \rho} = 0. \quad (18.43)$$

It is easy to see that the map (18.42) transforms roots of Eq. (18.31) into roots of Eq. (18.43) and real roots into real roots. Especially roots inside the unit circle into roots with negative real part. Thus if Eq. (18.31) possesses only one real root inside the unit circle then Eq. (18.43) possesses only one real root with the negative real part. Let w_0 be such a root of Eq. (18.43) and z and $\bar{z}$ be the remaining roots. One gets

$$(w - w_0)(w - z)(w - \bar{z}) = w^3 - (2\text{Re}(z) + w_0)w^2 + (|z|^2 + 2\text{Re}(z)w_0)w - w_0|z|^2 = 0. \quad (18.44)$$

Thus we only need

$$-w_0|z|^2 > 0, \quad (18.45)$$

$$x_0 = \frac{1 + w_0}{1 - w_0}, \quad (18.46)$$

which means

$$\frac{\rho + 1 - m^2 - \rho m^2}{7m^2 - 3\rho m^2 + 5 - \rho} > 0. \quad (18.47)$$

The last condition means also that Eq. (18.31) has a real root inside the unit circle, i.e. x_0 . Now we only need to satisfy $x_0 > 0$, which simply means

$$\frac{\rho}{3m^2} < 0 \quad (18.48)$$

or (if $m^2 > 0$)

$$\rho < 0. \quad (18.49)$$

In this way we get conditions

$$(\rho + 1 - m^2 - \rho m^2)(7m^2 - 3\rho m^2 + 5 - \rho) > 0, \quad (18.50)$$

$$\rho < 0. \quad (18.51)$$

From (18.50–51) we get two solutions:

$$\text{I} \quad 0 > \rho > -1, \quad 1 > m^2 > \frac{5 - \rho}{7 - 3\rho} \quad (18.52)$$

$$\text{II} \quad \rho < -1, \quad 0 < m^2 < \frac{5 - \rho}{7 - 3\rho}. \quad (18.53)$$

Simultaneously we have

$$D(\rho, m) > 0. \quad (18.54)$$

Under these conditions our solution given by Eq. (18.41) is non-negative with modulus smaller than one.

In this way we have

$$n_s(K) = 1 \quad (18.55)$$

for

$$\ln K = \frac{v_{\pm} + 2K(m^2)l}{C} + K'_0 \quad (18.56)$$

$$K = K_0 \exp\left(\frac{v_{\pm} + 2K(m^2)l}{C}\right) \quad (18.57)$$

where $K'_0 = \ln K_0$.

Moreover we are interested in a deviation from a flat power spectrum. Thus we consider

$$n_s(K) = 1 \pm \frac{dn_s(K)}{d \ln K} \Delta \ln K \quad (18.58)$$

where $\Delta \ln K \sim 10$ and $\frac{dn_s(K)}{d \ln K}$ is calculated for $\ln K$ given by the formula (18.55). According to the formula (18.15) $\frac{dn_s(K)}{d \ln K}$ is proportional to C^2 .

Thus using (18.26) we get

$$\frac{dn_s(K)}{d \ln K} \Delta \ln K \sim 10 \left(\frac{K(m^2)}{10}\right)^2 \sim 0.1 \quad (18.59)$$

which gives us rough estimate of a deviation from a flat power spectrum.

For K given by Eq. (18.57) one gets

$$\begin{aligned} \frac{dn_s(K)}{d \ln K} = 2C^2 & \left[\varepsilon \left(\varepsilon_0 x_0 \sqrt{1 - m^2 x_0} - (1 + x_0(m^2 - 1)) \sqrt{1 - x_0} \right) \right. \\ & \times \frac{\sqrt{1 - m^2 x_0} + \varepsilon \varepsilon_0 m^2 \sqrt{x_0(1 - x_0)}}{\sqrt{x_0(1 - x_0)}(1 - m^2 x_0)} \\ & \left. + \frac{1 + 3m^2 x_0^2 - x_0(2m^2 + 1)}{\rho + x_0} + \frac{4x_0(1 - x_0)(1 - m^2 x_0)}{(\rho + x_0)^2} \right] \end{aligned} \quad (18.60)$$

where

$$\varepsilon_0^2 = \varepsilon^2 = 1. \quad (18.61)$$

It is interesting to estimate a range of K . One can consider

$$K_0 \exp\left(\frac{v_+ - K(m^2)}{C}\right) \leq K \leq K_0 \exp\left(\frac{v_+ + K(m^2)}{C}\right). \quad (18.62)$$

Let us consider Eq. (18.58). If we find conditions for $\frac{dn_s(K)}{d \ln K}$ to be zero, we get a local flat spectrum (around K given by Eq. (18.57)). One gets using (18.29) and (18.60)

$$\begin{aligned} & \left(\varepsilon_0 x_0 \sqrt{1 - m^2 x_0} - (1 + x_0(m^2 - 1)) \sqrt{1 - x_0} \right) \\ & \times \frac{\sqrt{1 - m^2 x_0} + \varepsilon_0 m^2 \sqrt{x_0(1 - x_0)}}{\sqrt{x_0(1 - x_0)(1 - m^2 x_0)}} \\ & = \frac{x_0(2m^2 + 1) - 1 - 3m^2 x_0^2}{\rho + x_0} + \frac{4x_0(x_0 - 1)(1 - m^2 x_0)}{(\rho + x_0)^2}. \end{aligned} \quad (18.63)$$

For $x_0 = x_0(m, \rho)$ (see Eq. (18.41)) is a known function of m and ρ , the condition (18.63) can be considered as a constraint for m and ρ . Together with (18.52), (18.53) and (18.54) it can give an interesting range for parameters m and ρ . In this range the spectrum function (18.10) looks flat (around K given by Eq. (18.57)).

Let us consider a formula for an evolution of Higgs' field given by Eq. (14.369):

$$\Phi(t) = \frac{1}{2\alpha_s} \left(1 \pm \sqrt{\frac{5}{1 - e^{-5b(t-t_0)}}} \right) \quad (18.64)$$

where $b = \frac{e^{n\Psi_1} A}{6r^2 H_0} > 0$,

$$\lim_{t \rightarrow \infty} \Phi(t) = \frac{1}{\alpha_s} \left(\frac{\pm \sqrt{5} + 1}{2} \right).$$

It is easy to see that for

$$t_{\max} - t_0 = \frac{2}{b}, \quad (18.66)$$

$$\Phi\left(\frac{2}{b} + t_0\right) \simeq \frac{1}{\alpha_s} \left(\frac{1 \pm \sqrt{5}}{2} \right). \quad (18.67)$$

Thus we can consider from practical point of view that an inflation has been completed for $t = t_0 + \frac{2}{b}$. The inflation starts for

$$t_{\text{initial}} = t_0 + \frac{1}{5b} \ln \frac{5}{4}. \quad (18.68)$$

Let us calculate $P_R(K)$, a spectral function for this kind of evolution of Higgs' field. One gets from Eqs (18.8) and (18.9)

$$P_R(K) = P_0 \frac{\left(\frac{K}{K_0}\right)^{5\bar{\mu}}}{\left(\left(\frac{K}{K_0}\right)^{\bar{\mu}} - 1\right)^3}, \quad (18.69)$$

$$P_0 = \frac{4H_0^4 \alpha_s^2 n_1}{125\pi^2 b^2} \quad (18.70)$$

$$K_0 = R_0 H_0 e^{t_0 H_0}. \quad (18.71)$$

For a spectral index we get

$$\frac{d \ln P_R(K)}{d \ln K} = n_s(K) - 1 = \bar{\mu} \frac{2 \left(\frac{K}{K_0} \right)^{\bar{\mu}} - 5}{\left(\frac{K}{K_0} \right)^{\bar{\mu}} - 1} \quad (18.72)$$

$$\bar{\mu} = \frac{5b}{H_0}. \quad (18.73)$$

For the first derivative of n_s

$$\frac{dn_s}{d \ln K} = \frac{8\bar{\mu}^2 \left(\frac{K}{K_0} \right)^{\bar{\mu}}}{\left(\left(\frac{K}{K_0} \right)^{\bar{\mu}} - 1 \right)^2}. \quad (18.74)$$

An interesting question is to find the range of K . Taking

$$t_{\text{initial}}^* = \frac{1}{H_0} \left(\frac{K_{\text{initial}}}{R_0 H_0} \right) = t_0 + \frac{1}{5b} \ln \frac{5}{4} = t_{\text{initial}} \quad (18.75)$$

one gets

$$K_{\text{initial}} = K_0 \left(\frac{5}{4} \right)^{1/\bar{\mu}} \quad (18.76)$$

$$t_{\text{max}}^* = \frac{1}{H_0} \left(\frac{K_{\text{max}}}{R_0 H_0} \right) = t_0 + \frac{2}{b} = t_{\text{max}} \quad (18.77)$$

$$K_{\text{max}} = K_0 e^{2H_0/b} = K_0 e^{10/\bar{\mu}} \quad (18.78)$$

and finally

$$K_0 \left(\frac{5}{4} \right)^{1/\bar{\mu}} < K < K_0 e^{10/\bar{\mu}}. \quad (18.79)$$

Let us find $\tilde{K}$ such that

$$n_s(\tilde{K}) = 1. \quad (18.80)$$

One gets

$$\tilde{K} = K_0 \left(\frac{5}{2} \right)^{1/\bar{\mu}} \quad (18.81)$$

$$K_0 \left(\frac{5}{4} \right)^{1/\bar{\mu}} < \tilde{K} < K_0 e^{10/\bar{\mu}}. \quad (18.82)$$

For such a $\tilde{K}$ one gets

$$\frac{dn_s(\tilde{K})}{d \ln K} = 5\bar{\mu}^2 \quad (18.83)$$

$$n_s(\tilde{K}) \cong 1 \pm 5\bar{\mu}^2 \Delta \ln K. \quad (18.84)$$

If we take as usual $\Delta \ln K \simeq 10$, we get

$$n_s(K) \cong 1 \pm \frac{1250}{H_0^2} b^2 = 1 \pm \frac{625}{18r^4 H_0^4} e^{2n_{\Psi_1}} A^2. \quad (18.85)$$

We can try to make it sufficiently close to 1 taking a big H_0 .

In this case we have parametrized a spectral function and a spectral index by only three parameters $\bar{\mu}$, P_0 and K_0 , from which only $\bar{\mu}$ matters. Taking K from the range

$$1.2^{1/\bar{\mu}} < \frac{K}{K_0} < 10^{4.3/\bar{\mu}} \quad (18.86)$$

we can calculate in principle all CMB characteristics (see Ref. [128]) and to fit to observational data (as good as possible) parameters $\bar{\mu}$ (and K_0). This will be a subject of future investigations.

Finally (in order to give a complete set of spectral functions) let us come back to some formulae (see Eqs (14.529–531)):

$$P_R(K) = P_0 g\left(\frac{K}{R_0 H_0}\right) \quad (18.87)$$

$$g(x) = \frac{1}{x^2} \cdot \frac{2}{\left(\text{sd}(u, \frac{1}{2}) + \sqrt{2} \text{cd}(u, \frac{1}{2}) \text{nd}(u, \frac{1}{2})\right)^2}$$

$$u = \sqrt{2}C_1 x + K\left(\frac{1}{2}\right) - C_1 \sqrt{2} - \int_0^{\arccos(\sqrt{5}/C_1)} \frac{d\varphi}{\sqrt{1 - \frac{1}{2} \sin^2 \varphi}} \quad (18.88)$$

$$P_0 = \left(\frac{H_0}{2\pi}\right)^2 \cdot \frac{4\alpha_s^2 n_1}{5C_1^2} \quad (18.89)$$

$$n_s(K) - 1 = -2 - C_1 x \cdot \frac{\sqrt{2} \text{cd}(u, \frac{1}{2}) - \text{sd}^2(u, \frac{1}{2})}{\text{sd}(u, \frac{1}{2}) + \sqrt{2} \text{cd}(u, \frac{1}{2}) \text{nd}(u, \frac{1}{2})} \quad (18.90)$$

$$\frac{dn_s(K)}{d \ln K} = -2C_1 x \cdot \frac{\sqrt{2} \text{cd}(u, \frac{1}{2}) - \text{sd}^2(u, \frac{1}{2})}{\text{sd}(u, \frac{1}{2}) + \sqrt{2} \text{cd}(u, \frac{1}{2}) \text{nd}(u, \frac{1}{2})} - 2C_1^2 x^2 \text{sd}^2(u, \frac{1}{2})$$

$$- 2C_1 x^2 \cdot \frac{\left(\sqrt{2} \text{cd}(u, \frac{1}{2}) - \text{sd}^3(u, \frac{1}{2})\right)^2}{\left(\text{sd}(u, \frac{1}{2}) + \sqrt{2} \text{cd}(u, \frac{1}{2}) \text{nd}(u, \frac{1}{2})\right)^2} \quad (18.91)$$

In this case independent parameters are C_1 and $K_0 = R_0 H_0$. The amount of inflation in three cases can be calculated. In the first case (function (18.18)) it is given

by the corrected formulae (14.357–358), i.e.

$$\overline{N}_0 = H_0 \sqrt{\frac{2r^2}{5Aae^{n\Psi_1}}} \int_{\varphi_2}^{\varphi_1} \frac{d\theta}{\sqrt{1 - \left(\frac{4a+5}{10}\right) \sin^2 \theta}} \quad (18.92)$$

or

$$\overline{N}_0 = \frac{H_0}{\sqrt{5\alpha_s}} \sqrt[4]{\frac{4r^2}{ABe^{n\Psi_1}}} \int_{\varphi_2}^{\varphi_1} \frac{d\theta}{\sqrt{1 - \left(\frac{4a+5}{10}\right) \sin^2 \theta}} \quad (18.93)$$

$$\cos^2 \varphi_1 = \frac{1}{5+4a}, \quad \cos^2 \varphi_2 = \frac{5}{5+4a}. \quad (18.94)$$

It is interesting to notice that we can calculate $\overline{N}_0$ from Eq. (18.92) for simplified Weinberg-Salam model (see the first point of Ref. [75] in the nonsymmetric version). One gets

$$\begin{aligned} \overline{N}_0 = & J \cdot \sqrt{\frac{2}{15}} \left(\frac{m_{\tilde{A}}}{m_{\text{pl}}\alpha_s} \right)^{7/2} 2^{-27/4} \\ & \times \frac{\left(7g(\zeta, \mu) + \sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} \right)^{5/4}}{(2\mu^3 + 7\mu^2 + 5\mu + 20)^{7/4} |\zeta|^{1/4} (\mu^2 + 4)^{1/2}} \\ & \times \frac{(1 + \zeta^2)^{1/4} \left(49g^2(\zeta, \mu) + g(\zeta, \mu)(\mu^2 + 4) \sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} \right)^{1/2}}{\left(\ln \left(|\zeta| + \sqrt{\zeta^2 + 1} \right) + 2\zeta^2 + 1 \right)^{3/2} ((5\zeta^2 + 4) \ln 2 - 2\mu^2(1 + \zeta^2))^{1/4}} \end{aligned} \quad (18.92a)$$

where

$$\begin{aligned} f(\zeta) = & \frac{16|\zeta|^3(\zeta^2 + 1)}{3(2\zeta^2 + 1)(1 + \zeta^2)^{5/2}} \left(\zeta^2 E \left(\frac{|\zeta|}{\sqrt{\zeta^2 + 1}} \right) - (2\zeta^2 + 1) K \left(\frac{|\zeta|}{\sqrt{1 + \zeta^2}} \right) \right) \\ & + 8 \ln \left(|\zeta| \sqrt{\zeta^2 + 1} \right) + \frac{4(1 + 9\zeta^2 - 8\zeta^4)|\zeta|^3}{3(1 + \zeta^2)^{3/2}} \end{aligned} \quad (18.95)$$

$$g(\zeta, \mu) = -f(\zeta)(\mu^2 + 4) \quad (18.96)$$

$$\begin{aligned} h(\zeta, \mu) = & |\zeta|(2\mu^3 + 7\mu^2 + 5\mu + 20)(1 + \zeta^2)^{-1} \\ & \times ((5\zeta^2 + 4) \ln 2 - 2\mu^2(1 + \zeta^2)) \left(\ln \left(|\zeta| + \sqrt{\zeta^2 + 1} \right) + 2\zeta^2 + 1 \right) \end{aligned} \quad (18.97)$$

$$J = \int_{\varphi_2}^{\varphi_1} \frac{d\theta}{\sqrt{1 - \left(\frac{4a+5}{10}\right) \sin^2 \theta}} \quad (18.98)$$

$$\begin{aligned}
a = & \left(\frac{m_{\text{pl}}}{m_{\tilde{A}}} \right)^7 2^{35/2} \alpha_s^{16} B^{1/2} \frac{1}{m_{\tilde{A}}} \\
& \times \left(\frac{(2\mu^3 + 7\mu^2 + 5\mu + 20)}{(\mu^2 + 4) \left(7g(\zeta, \mu) + \sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} \right)} \right)^{7/2} \\
& \times \frac{(1 + \zeta^2)^{1/2} \left(\ln \left(|\zeta| + \sqrt{\zeta^2 + 1} \right) + 2\zeta^2 + 1 \right)^4}{|\zeta|^{1/2} ((5\zeta^2 + 4) \ln 2 - 2\mu^2(1 + \zeta^2))^{1/2}} \quad (18.99)
\end{aligned}$$

and a satisfies the condition (18.6), which is a condition for an integration constant B , $B > 0$. All of these formulae are subject to conditions

$$\mu > \mu_0 = -3.581552661 \dots \quad (\tilde{R}(\tilde{\Gamma}) > 0) \quad (18.100a)$$

$$|\zeta| > \zeta_0 = 1.36 \dots \quad (\tilde{P}(\zeta) < 0) \quad (18.100b)$$

$$\mu^2 < \frac{(5\zeta^2 + 4) \ln 2}{2(1 + \zeta^2)} \quad (A > 0). \quad (18.100c)$$

In the second case considered here the amount of an inflation is infinite (in principle). Moreover, it can be given by a finite formula

$$\begin{aligned}
N &= H_0 (t_{\text{end}} - t_{\text{initial}}) = H_0 (t_{\text{max}} - t_{\text{initial}}) \\
&= H_0 \left(\frac{2}{b} - \frac{1}{5b} \ln \frac{5}{4} \right) = \frac{H_0}{b} \left(2 - \frac{1}{5} \ln \frac{5}{4} \right) \cong 1.96 \frac{H_0}{b}. \quad (18.101)
\end{aligned}$$

In the third case the amount of an inflation is given by the formula (14.520)

$$\sqrt{2}C_2 - K \left(\frac{1}{2} \right) = \int_0^{\arccos\left(\frac{1}{2C_1 e^{N_0}}\right)} \frac{d\theta}{\sqrt{1 - \frac{1}{2} \sin^2 \theta}} - \sqrt{2}C_1 e^{N_0}, \quad (18.102)$$

which is a transcendental equation for N_0 .

All of those $P_R(K)$ functions will be treated by numerical methods from Ref. [128] and afterwards we compare the results with observational data. This will be the first stage of an application of our theory in cosmology. Afterwards we try to estimate intrinsic parameters of our theory and to start a multicomponent inflation.

Now we calculate a total amount of an inflation during two de Sitter phases in our cosmological models and corresponding masses of quintessence particles.

We consider a total amount of an inflation in cosmological models. In our approach we get two de Sitter phases governed by two different Hubble constants H_0 and H_1 . We underline that we have to do with phase transitions: of the second order in the configuration of Higgs' field and of the first order in the evolution of the Universe.

The Hubble constant in the first phase is given by the formula ($\hbar = c = 1$)

$$H_0 = \frac{\left(-n\gamma + \sqrt{n^2\gamma^2 + 4(n^2 - 4)\alpha_1\beta}\right)^{(n-2)/4}}{2^{(n+2)/4}(n+2)^{(n+2)/4}\sqrt{3}\beta^{n/4}} \times \left(n^2\gamma^2 - \gamma\sqrt{n^2\gamma^2 + 4(n^2 - 4)\alpha_1\beta} + 4\alpha_1\beta(n+2)\right)^{1/2}, \quad (18.103)$$

where

$$\gamma = \frac{m_{\tilde{A}}^2}{\alpha_s^2} \tilde{P} = \frac{1}{r^2} \tilde{P} \quad (18.104)$$

$$\beta = \frac{\alpha_s^2}{l_{\text{pl}}^2} \tilde{R}(\tilde{\Gamma}) = \alpha_s^2 m_{\text{pl}}^2 \tilde{R}(\tilde{\Gamma}) \quad (18.105)$$

$$\alpha_1 = \frac{\alpha_s^2}{r^2} \left(\frac{\alpha_s l_{\text{pl}}}{r}\right)^2 \left(\frac{A}{\alpha_s^6}\right) = m_{\tilde{A}}^2 \left(\frac{m_{\tilde{A}}}{m_{\text{pl}}}\right)^2 \left(\frac{A}{\alpha_s^6}\right). \quad (18.106)$$

α_s is a coupling constant, m_{pl} and l_{pl} are Planck's mass and Planck's length, respectively, $m_{\tilde{A}}$ and r are a scale of an energy (a scale of a mass of broken gauge bosons).

$\tilde{R}(\tilde{\Gamma})$ is a scalar of a curvature of a connection on a group manifold H .

$$\tilde{P} = \frac{1}{V_2} \int_M R(\hat{\Gamma}) \sqrt{|\tilde{g}|} d^{n_1} x, \quad (18.107)$$

where V_2 is a measure of a manifold M (a vacuum states manifold), $R(\hat{\Gamma})$ is a scalar of a curvature for a connection defined on M , $\tilde{g} = \det(g_{\tilde{a}\tilde{b}})$.

$$A = \frac{1}{V_2} \int_M \left(l_{ab} \left(g^{[\tilde{m}\tilde{n}]} g^{[\tilde{a}\tilde{b}]} C_{\tilde{a}\tilde{b}}^b C_{\tilde{m}\tilde{n}}^a - C_{\tilde{m}\tilde{n}}^b g^{\tilde{a}\tilde{n}} g^{\tilde{b}\tilde{m}} \tilde{L}_{\tilde{a}\tilde{b}}^a \right) \right) \sqrt{\tilde{g}} d^{n_1} x, \quad (18.108)$$

$$l_{dc} g_{\tilde{m}\tilde{b}} g^{\tilde{c}\tilde{b}} \tilde{L}_{\tilde{c}\tilde{a}}^d + l_{cd} g_{\tilde{a}\tilde{m}} g^{\tilde{m}\tilde{c}} \tilde{L}_{\tilde{b}\tilde{c}}^d = 2l_{cd} g^{\tilde{a}\tilde{m}} g^{\tilde{m}\tilde{c}} C_{\tilde{b}\tilde{c}}^d. \quad (18.109)$$

$C_{\tilde{b}\tilde{c}}^d$ are structure constants for a Lie algebra $\mathfrak{h}$ (a Lie algebra of a group H).

Using Eqs (18.104–106) one gets

$$H_0 = \left(\frac{m_{\tilde{A}}}{\alpha_s}\right) \left(\frac{m_{\tilde{A}}}{\alpha_s^2 m_{\text{pl}}}\right)^{n/2} \frac{\left(-n\tilde{P} + \sqrt{n^2\tilde{P}^2 + 4(n^2 - 4)\tilde{R}(\tilde{\Gamma})A}\right)^{(n-2)/4}}{2^{(n+2)/4}(n+2)^{(n+2)/4}\sqrt{3}(\tilde{R}(\tilde{\Gamma}))^{n/4}} \times \left(n^2\tilde{P}^2 - \tilde{P}\sqrt{n^2\tilde{P}^2 + 4(n^2 - 4)A\tilde{R}(\tilde{\Gamma})} + 4(n+2)A\tilde{R}(\tilde{\Gamma})\right)^{1/2}, \quad (18.103a)$$

In the case of more general models (two different types of critical points for Higgs' field configuration) we take for A

$$A = 4\alpha_s^2 V(\Phi_{\text{crt}}^1) \quad (18.110)$$

where $V(\Phi_{\text{crt}}^1)$ is a value of Higgs' potential for Φ_{crt}^1 (a metastable state Higgs' field configuration).

In the case of a Hubble constant H_1 we get

$$H_1 = \frac{|\gamma|^{(n+2)/4} n^{n/4}}{\sqrt{3} \beta^{n/4} (n+2)^{(n+2)/4}} \quad (18.111)$$

or

$$H_1 = \left(\frac{m_{\tilde{A}}}{m_{\text{pl}}} \right)^{n/2} \frac{m_{\tilde{A}}}{\alpha_s^{n+1}} \frac{|\tilde{P}|^{(n+2)/4} n^{n/4}}{\sqrt{3} (n+2)^{(n+2)/4} (\tilde{R}(\tilde{I}))^{n/4}} \quad (18.111a)$$

In this case we suppose that $\gamma < 0$ and $\alpha_1, \beta > 0$. This is possible because γ is a function of a constant ζ and can have any sign. In the case of β it is the same. β is a function of a constant ξ (μ here). This is because of properties of $\tilde{R}(\tilde{I})$ and $\tilde{P}$.

We consider here several consequences of Higgs' field dynamics on primordial fluctuations (perturbations) spectrum. We find three different functions and examine their properties. We write down also an equation for an amount of an inflation in the first de Sitter phase for these functions. They are calculable for functions (18.10) and (18.87). In the case of a function (18.69) we get under some practical assumption

$$N_0 = 1.96 \frac{H_0}{b} = 1.96 \frac{6r^2 H_0^2}{e^{n\Psi_1} A} = \frac{1.96}{\alpha_s^4} \left(\frac{m_{\tilde{A}}}{m_{\text{pl}}} \right)^2 F(\beta, \gamma, \alpha_1) \quad (18.112)$$

where $F(\beta, \gamma, \alpha_1)$ is given by the formula

$$F(\beta, \gamma, \alpha_1) = \frac{\left(n^2 \gamma^2 - \gamma \sqrt{n^2 \gamma^2 + 4(n^2 - 4)\alpha_1 \beta} + 4\alpha_1 \beta (n+2) \right)}{(n+2)\alpha_1 \left(-n\gamma + \sqrt{n^2 \gamma^2 + 4(n^2 - 4)\alpha_1 \beta} \right)} \quad (18.113)$$

or

$$N_0 = 1.96 \left(\frac{m_{\tilde{A}}}{m_{\text{pl}}} \right)^4 F(\tilde{R}(\tilde{I}), \tilde{P}, A). \quad (18.113a)$$

Now let us come to the second de Sitter phase. During this phase the inflation is driven by the field Ψ which changes (slowly $\dot{\Psi} \approx 0$) from Ψ_0 to $\frac{1}{2} \ln \left(\frac{\beta}{|\gamma|} \right)$. We develop in Sec. 14 two approximation schemes for an evolution of Ψ . In the case of a harmonic oscillation we calculate an amount of an inflation. We find also a different scheme—the so called slow-roll approximation which offers an infinite in time evolution of Ψ .

Moreover from practical point of view we can suppose that an evolution starts if Ψ is very close to $\frac{1}{2} \ln \left(\frac{n|\gamma|}{(n+2)\beta} \right)$. In term of a variable

$$y = \sqrt{\frac{\beta}{|\gamma|}} e^{\Psi} \quad (18.114)$$

y is close to $\sqrt{\frac{n}{n+2}}$. Let y be

$$y = \sqrt{\frac{n}{n+2}} + \varepsilon \quad (18.115)$$

where $\varepsilon > 0$ is very small. Let us take

$$\varepsilon = \frac{1}{n(n+1)(n+2)}. \quad (18.116)$$

We remind that $n \geq 14$. Thus

$$\varepsilon < 3 \cdot 10^{-4}. \quad (18.117)$$

Let us consider the formula (14.416) and let us calculate the limit

$$\lim_{y \rightarrow \sqrt{\frac{n}{n+2}} + \varepsilon} I. \quad (18.118)$$

One gets after some simplifications (taking under consideration the fact that $n > 14$)

$$\lim_{y \rightarrow \sqrt{\frac{n}{n+2}} + \varepsilon} I = -\frac{\pi}{2} - \frac{1}{2\sqrt{2n}} \ln 2. \quad (18.119)$$

From the other side we have

$$\lim_{y \rightarrow 1} I = -\frac{\pi}{2} + \frac{1}{\sqrt{2n}} \ln \left(n + 1 + \sqrt{n(n+2)} \right). \quad (18.120)$$

We have

$$I = \overline{B}(t - t_0) \quad (18.121)$$

where

$$\overline{B} = 4\sqrt{2\pi}(n+2)|\gamma| \left(\frac{n}{n+2} \right)^{n/2} \left(\frac{|\gamma|}{\beta} \right)^{n/2} \frac{1}{m_{\text{pl}}} \sqrt{M} \quad (18.122)$$

or

$$\overline{B} = 4\sqrt{2\pi} \left(\frac{m_{\tilde{A}}}{m_{\text{pl}}} \right)^{n+1} (n+2) \left(\frac{n}{n+2} \right)^{n/2} \frac{m_{\tilde{A}}}{\alpha_s^{2(n+1)} |\tilde{P}|} \left(\frac{|\tilde{P}|}{\tilde{R}(\tilde{I})} \right)^{n/2} \frac{1}{\sqrt{M}}. \quad (18.122a)$$

Writing

$$I(1) = \overline{B}(t_{\text{end}}^{\text{II}} - t_0) \quad (18.123)$$

$$I \left(\sqrt{\frac{n}{n+2}} + \varepsilon \right) = \overline{B}(t_{\text{initial}}^{\text{II}} - t_0) \quad (18.124)$$

where $t_{\text{initial}}^{\text{II}}$ and $t_{\text{end}}^{\text{II}}$ mean an initial and end time of the second de Sitter phase, one gets

$$\Delta t^{\text{II}} = t_{\text{end}}^{\text{II}} - t_{\text{initial}}^{\text{II}} = \frac{1}{\overline{B}} \frac{1}{\sqrt{2n}} \ln \left(\sqrt{2} \left(n + 1 + \sqrt{n(n+2)} \right) \right) \quad (18.125)$$

where Δt^{II} means a period of time of the second de Sitter phase. Thus the amount of inflation for this phase is

$$N_1 = H_1 \Delta t^{\text{II}} = \frac{H_1}{B} \frac{\ln \left(\sqrt{2} \left(n + 1 + \sqrt{n(n+2)} \right) \right)}{\sqrt{2n}}. \quad (18.126)$$

Using formulae (18.122) and (18.111) one gets

$$N_1 = \sqrt{2\pi} \left(\frac{\beta}{|\gamma|} \right)^{n/4} \left(\frac{n+2}{n} \right)^{n/4} \frac{\ln \left(\sqrt{2} \left(n + 1 + \sqrt{n(n+2)} \right) \right)}{\sqrt{6n|\gamma|}(n+2)^3} \cdot \frac{m_{\text{pl}}}{\sqrt{M}} \quad (18.127)$$

or

$$N_1 = \left(\frac{2\sqrt{2\pi}}{\sqrt{M}} \right) \left(\frac{m_{\text{pl}}}{m_{\tilde{A}}} \right)^{(n+2)/2} \left(\frac{\tilde{R}(\tilde{\Gamma})}{|\tilde{P}|} \right)^{n/4} \left(\frac{n+2}{n} \right)^{n/4} \times \frac{\ln \left(\sqrt{2} \left(n + 1 + \sqrt{n(n+2)} \right) \right)}{2\sqrt{6}\alpha_s^{n-1}|\tilde{P}|^{1/2}(n+2)^3}. \quad (18.127a)$$

For large n one finds

$$N_1 \simeq \sqrt{2\pi} \sqrt{\frac{e}{6}} \left(\frac{\beta}{|\gamma|} \right)^{n/4} \frac{\ln n}{|\gamma|^{1/2} n^{7/2}} \cdot \frac{m_{\text{pl}}}{\sqrt{M}} \quad (18.128)$$

or

$$N_1 \simeq \sqrt{\frac{e}{6}} \left(\frac{\tilde{R}(\tilde{\Gamma})}{|\tilde{P}|} \right)^{n/4} \left(\frac{m_{\text{pl}}}{m_{\tilde{A}}} \right)^{(n+2)/2} \left(\frac{\sqrt{2\pi}}{\sqrt{M}} \right) \frac{\ln n}{\alpha_s^{n-1}|\tilde{P}|^{1/2} n^{7/2}}. \quad (18.128a)$$

The total amount of an inflation considered in the paper is

$$N_{\text{tot}} = N_0 + N_1 \quad (18.129)$$

and should be fixed to ~ 60 .

In order to give an example of these calculations let us consider a six-dimensional Weinberg-Salam model (see Ref. [75], the first point). It is of course a bosonic part of this model. In this case $H = G2$, $\dim H = 14$, $M = S^2$ (two-dimensional sphere). This will be of course the model in Nonsymmetric Kaluza–Klein (Jordan–Thiry) Theory. In order to simplify the calculations we take

$$\tilde{R}(\tilde{\Gamma}) = \frac{2(2\mu^3 + 7\mu^2 + 5\mu + 20)}{(\mu^2 + 4)^2} \quad (18.130)$$

(see formula (7.21) from the fourth point of Ref. [18]).

The scalar curvature has been calculated here for $H = SO(3) \simeq SU(2)$. Moreover it gives a taste of the full theory

$$\begin{aligned} \tilde{\underline{P}} &= \tilde{\underline{P}}(\zeta) \\ &= \left\{ \frac{16|\zeta|^3(\zeta^2+1)}{3(2\zeta^2+1)(1+\zeta^2)^{5/2}} \left(\zeta^2 E \left(\frac{|\zeta|}{\sqrt{\zeta^2+1}} \right) - (2\zeta^2+1) K \left(\frac{|\zeta|}{\sqrt{\zeta^2+1}} \right) \right) \right. \\ &\quad \left. + 8 \ln \left(|\zeta| \sqrt{\zeta^2+1} \right) + \frac{4(1+9\zeta^2-8\zeta^4)|\zeta|^3}{3(1+\zeta^2)^{3/2}} \right\} / \left(\ln \left(|\zeta| + \sqrt{\zeta^2+1} \right) + 2\zeta^2+1 \right) \end{aligned} \quad (18.131)$$

(see formula (8.64)).

For V_2 one gets

$$V_2 = \frac{2\pi}{|\zeta|} \left(\ln \left(|\zeta| + \sqrt{\zeta^2+1} \right) + 2\zeta^2+1 \right) \quad (18.132)$$

(see formula (8.65)), where

$$K(k) = \int_0^{\pi/2} \frac{d\theta}{\sqrt{1-k^2 \sin^2 \theta}}, \quad (18.133)$$

$$E(k) = \int_0^{\pi/2} \sqrt{1-k^2 \sin^2 \theta} d\theta. \quad (18.134)$$

$\tilde{\underline{P}}$ has been calculated for an Einstein-Kaufmann connection defined on S^2 . We need $\tilde{\underline{P}} < 0$, thus in our case

$$|\zeta| > \zeta_0 = 1.36 \dots \quad (18.135)$$

(see Fig. 6).

In order to write the formula in this case we need to calculate A from Eq. (18.108). We should calculate $\tilde{L}_{\tilde{b}\tilde{c}}^d$ from Eq. (18.109). In our case this is quite easy for S^2 is 2-dimensional and $\tilde{a}, \tilde{b} = 1, 2$ ($\tilde{a}, \tilde{b} = 5, 6$ if we embed S^2 as a vacuum state manifold in the full theory).

In this case one easily finds

$$\tilde{L}_{\tilde{a}\tilde{b}}^c = h^{ce} l_{ed} C_{\tilde{a}\tilde{b}}^d \quad (18.136)$$

where $C_{\tilde{a}\tilde{b}}^d$ are structure constants of the group H in such a way that $\tilde{a}, \tilde{b}$ correspond to the complement $\mathfrak{m}$, $\mathfrak{g} = \mathfrak{g}_0 \dot{+} \mathfrak{m}$,

$$\mathfrak{g} \subset \mathfrak{h} \quad (18.137)$$

where $\mathfrak{g} = A_1$, $l_{ab} = h_{ab} + \mu k_{ab}$, $h^{ef} h_{fd} = \delta_d^e$. Using the exact form of the nonsymmetric tensor on $SO(3)$ and the nonsymmetric tensor on S^2 (see formulae (2.24a), (6.31)) one

gets

$$\begin{aligned}\bar{A} &= l_{ab} \left(g^{[\tilde{m}\tilde{n}]} g^{[\tilde{a}\tilde{b}]} C_{\tilde{m}\tilde{n}}^a C_{\tilde{a}\tilde{b}}^b - C_{\tilde{m}\tilde{n}}^b g^{\tilde{a}\tilde{n}} g^{\tilde{b}\tilde{m}} \tilde{L}_{\tilde{a}\tilde{b}}^a \right) \\ &= - \left(\frac{\zeta^2}{(1+\zeta^2)^2 \sin^2 \theta} + \frac{4 + \mu^2 \sin^2 \theta}{(1+\zeta^2) \sin^2 \theta} \right)\end{aligned}\quad (18.138)$$

and finally

$$A = \frac{1}{V_2} \int_{S^2} \bar{A} \sqrt{\tilde{g}} d\theta d\varphi = \frac{1}{V_2} \int_0^{2\pi} \int_0^\pi d\varphi d\theta \bar{A} \sqrt{\tilde{g}}, \quad (18.139)$$

$$A = \frac{|\zeta| [(5\zeta^2 + 4) \ln 2 - 2\mu^2(1 + \zeta^2)]}{(1 + \zeta^2) \left(\ln \left(|\zeta| + \sqrt{\zeta^2 + 1} \right) + 2\zeta^2 + 1 \right)} \quad (18.140)$$

where the integral $\int_0^\pi \frac{d\theta}{\sin \theta}$ has been calculated in the sense of a principal value:

$$\int_0^\pi \frac{d\theta}{\sin \theta} = \lim_{\varepsilon \rightarrow 0^+} \int_\varepsilon^{\pi-\varepsilon} \frac{d\theta}{\sin \theta} = -\ln 2. \quad (18.141)$$

Using Eqs (18.130–131), (18.140) one writes the formula (18.103a) in the form ($n = 14$)

$$\begin{aligned}H_0 &= \frac{1}{2^{19} \sqrt{3}} \left(\frac{m_{\tilde{A}}}{\alpha_s^2 m_{\text{pl}}} \right)^7 \cdot \frac{m_{\tilde{A}}}{\alpha_s} \\ &\times \frac{\left(7g(\zeta, \mu) + \sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} \right)^3 (\mu^2 + 4)^3}{\left(\ln \left(|\zeta| + \sqrt{\zeta^2 + 1} \right) + 2\zeta^2 + 1 \right) (2\mu^3 + 7\mu^2 + 5\mu + 20)^{7/2}} \\ &\times \left(98g^2(\zeta, \mu) + (\mu^2 + 4)g(\zeta, \mu)\sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} + 32H(\zeta, \mu) \right)^{1/2}\end{aligned}\quad (18.142)$$

where

$$g(\zeta, \mu) = -f(\zeta)(\mu^2 + 4) \quad (18.143)$$

$$\begin{aligned}h(\zeta, \mu) &= |\zeta|(2\mu^3 + 7\mu^2 + 5\mu + 20) \left((5\zeta^2 + 4) \ln 2 - 2\mu^2(1 + \zeta^2) \right) \\ &\times \frac{\left(\ln \left(|\zeta| + \sqrt{\zeta^2 + 1} \right) + 2\zeta^2 + 1 \right)}{(1 + \zeta^2)}\end{aligned}\quad (18.144a)$$

$$H(\zeta, \mu) = h(\zeta, \mu) \left(\ln \left(|\zeta| + \sqrt{\zeta^2 + 1} \right) + 2\zeta^2 + 1 \right) \quad (18.144b)$$

$$\begin{aligned}f(\zeta) &= \frac{16|\zeta|^3(\zeta^2 + 1)}{3(2\zeta^2 + 1)(1 + \zeta^2)^{5/2}} \left(\zeta^2 E \left(\frac{|\zeta|}{\sqrt{\zeta^2 + 1}} \right) - (2\zeta^2 + 1) K \left(\frac{|\zeta|}{\sqrt{\zeta^2 + 1}} \right) \right) \\ &+ 8 \ln \left(|\zeta| \sqrt{\zeta^2 + 1} \right) + \frac{4(1 + 9\zeta^2 - 8\zeta^4)|\zeta|^3}{3(1 + \zeta^2)^{3/2}}.\end{aligned}\quad (18.145)$$

From Eq. (18.111a) one gets

$$H_1 = \sqrt{\frac{2}{3}} \left(\frac{m_{\tilde{A}}}{m_{\text{pl}}} \right)^7 \frac{m_{\tilde{A}}}{\alpha_s^{15}} \cdot \frac{7^7}{2^{13}} \times \frac{g^4(\zeta, \mu)(\mu^2 + 4)^4}{(2\mu^3 + 7\mu^2 + 5\mu + 20)^{7/2}} \cdot \left(\ln \left(|\zeta| + \sqrt{\zeta^2 + 1} \right) + 2\zeta^2 + 1 \right)^{-4}. \quad (18.146)$$

From Eq. (18.113a) one finds

$$N_0 = 0.12 \left(\frac{m_{\tilde{A}}}{m_{\text{pl}}} \right)^4 \times \frac{\left(7g(\zeta, \mu) + \sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} \right)^{-1} (\mu^2 + 4)^2 (1 + \zeta^2)}{|\zeta| ((5\zeta^2 + 4) \ln 2 - 2\mu^2 (1 + \zeta^2))} \times \left(98g^2(\zeta, \mu) + (\mu^2 + 4)g(\zeta, \mu)\sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} + 32H(\zeta, \mu) \right). \quad (18.147)$$

Equation (18.122a) is transformed into

$$\overline{B} = \left(\frac{m_{\tilde{A}}}{m_{\text{pl}}} \right)^{15} \left(\frac{7^7 m_{\tilde{A}}}{2^{16} \alpha_s^{30}} \right) \left(\frac{|f(\zeta)|}{2\mu^3 + 7\mu^2 + 5\mu + 20} \right)^7 \cdot \frac{1}{\sqrt{|\overline{M}|}} \times \frac{|f(\zeta)|(\mu^2 + 4)^6}{\left(\ln \left(|\zeta| + \sqrt{\zeta^2 + 1} \right) + 2\zeta^2 + 1 \right)^8}. \quad (18.148)$$

And finally Eq. (18.127a) gives

$$N_1 \simeq 5.47 \cdot 10^4 \left(\frac{m_{\text{pl}}}{m_{\tilde{A}}} \right)^9 \left(\frac{|f(\zeta)|}{2\mu^3 + 7\mu^2 + 5\mu + 20} \right)^{7/2} \cdot \frac{\alpha_s}{\sqrt{|\overline{M}|}} \times \frac{\left(\ln \left(|\zeta| + \sqrt{\zeta^2 + 1} \right) + 2\zeta^2 + 1 \right)^{3/2}}{|f(\zeta)|^{1/2} (\mu^2 + 4)^3}. \quad (18.149)$$

Let us notice that according to our assumption $f(\zeta) < 0$ ($\tilde{P} < 0$), $2\mu^3 + 7\mu^2 + 5\mu + 20 > 0$ ($\tilde{R}(\tilde{T}) > 0$), and

$$\frac{(5\zeta^2 + 4) \ln 2}{2(1 + \zeta^2)} > \mu^2 \quad (18.150)$$

(from $A > 0$).

The function on the left hand side of the inequality (18.150) is rising for $\zeta > 0$ and falling for $\zeta < 0$, having a minimum at $\zeta = 0$ equal to $2 \ln 2$. For $\zeta \rightarrow \pm\infty$ it is going to $\frac{5}{2} \ln 2$. For a specific value $\zeta = \zeta_0 = \pm 1.36$ (see Eq. (18.135)), it is equal to 1.61.

Thus we get

$$|\mu| \leq \sqrt{2 \ln 2} \approx 1.17741. \quad (18.151)$$

Moreover the polynomial $W(\mu) = 2\mu^3 + 7\mu^2 + 5\mu + 20$ possesses one real root

$$\mu_0 = -\frac{\sqrt[3]{1108 + 3\sqrt{135645}}}{6} - \frac{7}{6} - \frac{19}{6 \cdot \sqrt[3]{1108 + 3\sqrt{135645}}} = -3.581552661 \dots$$

Thus we can have $\tilde{R}(\tilde{\Gamma}) > 0$ for $\mu > \mu_0$ and in the region given by (18.151). Let us notice that taking sufficiently big $|\zeta|$ we can make N_0 in (18.147) arbitrarily big, i.e. about 60. From the other side if we take $|\zeta|$ sufficiently big, N_1 can also be arbitrarily large (i.e. ~ 60). Thus it seems that in this simple example it is enough to consider arbitrarily big ζ in order to get large amount of an inflation.

It is interesting to find in this simplified model a constant a (see Eq. (18.6)). One gets after some algebra

$$\begin{aligned} a^2 &= \frac{\alpha_s^4 B}{A m_{\tilde{A}}^2 e^{n\Psi_1}} = \left(\frac{m_{\text{pl}}}{m_{\tilde{A}}} \right)^{14} 2^{35} \alpha_s^{32} \left(\frac{B}{m_{\tilde{A}}^2} \right) \\ &\times \left(\frac{2\mu^3 + 7\mu^2 + 5\mu + 20}{(\mu^2 + 4) \left(7g(\zeta, \mu) + \sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} \right)} \right)^7 \\ &\times \frac{(1 + \zeta^2) \left(\ln \left(|\zeta| + \sqrt{\zeta^2 + 1} \right) + 2\zeta^2 + 1 \right)^8}{|\zeta| ((5\zeta^2 + 4) \ln 2 - 2\mu^2(1 + \zeta^2))} \end{aligned} \quad (18.153)$$

(we put $n = 14$).

The condition

$$0 < a^2 < 0.09703 \quad (18.154)$$

gives a constraint on an integration constant B . In this case $A > 0$. However we consider in Sec. 14 a special type of a dynamics of Higgs' field with $A < 0$. In this case we have the following constraint imposed on the constant $\bar{a}$. One gets after some algebra

$$\begin{aligned} \bar{a} &= \frac{e^{n\Psi_1} m_{\tilde{A}}^2 A}{2H_0^2 \alpha_s^2} \\ &= \frac{192 \left(7g(\zeta, \mu) + \sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} \right) \left(\ln \left(|\zeta| + \sqrt{\zeta^2 + 1} \right) + 2\zeta^2 + 1 \right)}{98g^2(\zeta, \mu) + (\mu^2 + 4)g(\zeta, \mu)\sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} + 32H(\zeta, \mu)} \\ &\times \frac{|\zeta|(\mu^2 + 4) ((5\zeta^2 + 4) \ln 2 - 2\mu^2(1 + \zeta^2))}{1 + \zeta^2}. \end{aligned} \quad (18.155)$$

Moreover in this case we should put

$$\bar{a} = -\frac{4}{15}, \quad (18.156)$$

i.e. $A < 0$ and we get

$$\begin{aligned} & 720 \left(7g(\zeta, \mu) + \sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} \right) \left(\ln \left(|\zeta| + \sqrt{\zeta^2 + 1} \right) + 2\zeta^2 + 1 \right) \\ & \times |\zeta|(\mu^2 + 4) (2\mu^2(1 + \zeta^2) - (5\zeta^2 + 4) \ln 2) \\ & = \left(98g^2(\zeta, \mu) + (\mu^2 + 4)g(\zeta, \mu)\sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} + 32H(\zeta, \mu) \right) (1 + \zeta^2). \end{aligned} \quad (18.157)$$

Eq. (18.157) gives a constraint among parameters of the theory. If Eq. (18.157) is satisfied, we have the dynamics of Higgs' field described by Eq. (14.381).

In this way we can consider the spectral function (14.107) and all the consequences coming from it. We need of course some supplementary conditions

$$\frac{1}{2} \cdot \frac{5\zeta^2 + 4}{1 + \zeta^2} \cdot \ln 2 < \mu^2 \quad (18.158)$$

and as usual

$$f(\zeta) < 0 \quad (18.159)$$

$$2\mu^3 + 7\mu^2 + 5\mu + 20 > 0. \quad (18.160)$$

In the case when condition (18.154) is satisfied we can consider a different dynamics of Higgs' field described by Eq. (18.1) leading to the spectral function (18.10) with all the consequences of this function. In this case we have condition (18.150) and conditions (18.159) and (18.160). They are easily satisfied.

The programme of research given here will give an additional constraint and could (in principle) lead to the realistic theory with more complicated groups and patterns of symmetry breaking.

Let us notice that in our theory we get cosmological terms. These terms are described by constants β, γ and α_1 which are proportional to $\tilde{R}(\tilde{I})$, $\tilde{P}$ and A . The importance of a cosmological constant is now obvious. Thus it is necessary to control these terms. They depend on constants ζ and μ . The first step in order to control them is to find conditions when they are equal to zero. In the simplified model we have found these conditions:

$$\beta(\mu_0 = -3.581552661 \dots) = 0 \quad (18.161)$$

$$\gamma(\zeta = \pm 1.36 \dots) = 0 \quad (18.162)$$

and

$$\alpha_1 = 0 \quad (18.163)$$

for

$$\mu = \pm \sqrt{\frac{\ln 2(5\zeta^2 + 4)}{2(\zeta^2 + 1)}}. \quad (18.164)$$

In this way we control the sign of A , $\tilde{P}$ and $\tilde{R}(\tilde{\Gamma})$ playing with constants ζ and μ :

$$\tilde{R}(\tilde{\Gamma}) > 0 \quad \text{for } \mu > \mu_0 = -3.581552661\dots \quad (18.165)$$

$$\tilde{P} < 0 \quad \text{for } |\zeta| > 1.36. \quad (18.166)$$

The sign of A is also controllable. All of these results give us interesting cosmological consequences.

It is interesting to find some conditions on stability of the first de Sitter evolution of the Universe. In Sec. 14 we find a criterion

$$M > M_0 \quad (18.167)$$

where

$$M_0 = \frac{4}{3} \times \frac{(n+2)\gamma\sqrt{n^2\gamma^2 + 4(n^2-4)\alpha_1\beta} - 4(n+2)^2(n-1)\alpha_1\beta - n^2(n+2)\gamma^2}{n\gamma^2 + 4(n+2)\alpha_1\beta - \gamma\sqrt{n^2\gamma^2 + 4(n^2-4)\alpha_1\beta}} \quad (18.168)$$

or

$$M_0 = \frac{4}{3} \times \frac{(n+2)\tilde{P}\sqrt{n^2\tilde{P}^2 + 4(n^2-4)A\tilde{R}(\tilde{\Gamma})} - 4(n+2)^2(n-1)A\tilde{R}(\tilde{\Gamma}) - n^2(n+2)\tilde{P}^2}{n\tilde{P}^2 + 4(n+2)A\tilde{R}(\tilde{\Gamma}) - \tilde{P}\sqrt{n^2\tilde{P}^2 + 4(n^2-4)A\tilde{R}(\tilde{\Gamma})}}, \quad (18.168a)$$

If

$$\tilde{P} > 0, \quad \tilde{R}(\tilde{\Gamma}) > 0, \quad A > 0, \quad (18.169)$$

then

$$M_0 > 0. \quad (18.170)$$

Moreover if

$$\tilde{P} < 0, \quad \tilde{R}(\tilde{\Gamma}) > 0, \quad A > 0, \quad (18.171)$$

the stability condition is ($M_0 < 0$)

$$M < 0. \quad (18.172)$$

Moreover for $A < 0$ the condition $M_0 < 0$ is not trivial and we should have

$$4(n+2)^2(n-1)|A|\tilde{R}(\tilde{\Gamma}) > (n+2)|\tilde{P}| \left(n^2|\tilde{P}| + \sqrt{n^2\tilde{P}^2 - 4(n^2-4)|A|\tilde{R}(\tilde{\Gamma})} \right). \quad (18.173)$$

In the case of $SO(3)$ group and S^2 vacuum states manifold (with $n = 14$) we have

$$M(SO(3)) = -\frac{2(36 + 7\mu^2)}{(4 + \mu^2)M_{\text{pl}}^2} < 0 \quad (18.174)$$

(see the equation on p. 260 of the second point of Ref. [18]).

Thus if $M_0 < 0$ the first de Sitter evolution of the Universe is stable. It means it happens if condition (18.173) is satisfied. One finds:

$$\begin{aligned} & \frac{416|\zeta|}{1 + \zeta^2} (2\mu^2(1 + \zeta^2) - (5\zeta^2 + 4) \ln 2) \cdot (2\mu^3 + 7\mu^2 + 5\mu + 20) \\ & > \frac{g(\zeta, \mu) \left(49g(\zeta, \mu) + \sqrt{49g^2(\zeta, \mu) - 384|h(\zeta, \mu)|} \right)}{\ln \left(\left(|\zeta| + \sqrt{\zeta^2 + 1} \right) + 2\zeta^2 + 1 \right)}. \end{aligned} \quad (18.175)$$

We have also supplementary conditions

$$|\zeta| > |\zeta_0| = 1.36 \dots \quad (18.176)$$

$$\mu > \mu_0 = -3.581552661 \dots \quad (18.177)$$

$$\mu^2 > \frac{\ln 2}{2} \cdot \frac{5\zeta^2 + 4}{\zeta^2 + 1}. \quad (18.178)$$

During the second de Sitter phase the evolution of the Universe is unstable against small perturbation of initial data. Moreover it can be made stable if $M < 0$. However, the second de Sitter phase in our model should be unstable because it ends with a radiation epoch. The instability of this phase in this particular case is caused by some physical processes beyond cosmological models.

The interesting point in the theory is a mass of the scalar field (a scalar particle) during both de Sitter phases. However, in that case we should consider a quintessence field Q

$$Q = \sqrt{|\overline{M}|} \frac{\Psi}{M_{\text{pl}}}. \quad (18.179)$$

In both de Sitter phases we can consider small oscillations q_k of the quintessence field around an equilibrium

$$Q = Q_k + q_k, \quad k = 0, 1. \quad (18.180)$$

For these small oscillations one finds

$$\overline{m}_k^2 = -\frac{1}{2} \frac{d^2 \lambda_{ck}}{d\Psi^2}(\Psi_k) \quad (18.181)$$

or

$$\begin{aligned}
\overline{m}_1^2 &= \frac{-\left(-n\gamma + \sqrt{n^2\gamma^2 + 4(n^2 - 4)\alpha_1\beta}\right)^{(n-2)/2}}{2^{(n+4)/2}(n+2)^{n/2}\beta^{n/2}} \\
&\quad \times \left(n\gamma\sqrt{n^2\gamma^2 + 4(n^2 - 4)\alpha_1\beta} - 2\gamma^2n^2 - 4(n+2)(n-3)\alpha_1\beta\right) \\
&= \left(\frac{m_{\tilde{A}}}{\alpha_s m_{\text{pl}}}\right)^n \left(\frac{m_{\tilde{A}}}{\alpha_s}\right)^2 (n+2)^{-n/2} 2^{-(n+4)/2} \\
&\quad \times \left(-\tilde{P} + \sqrt{n^2\tilde{P}^2 + 4(n^2 - 4)A\tilde{R}(\tilde{\Gamma})}\right)^{(n-2)/2} \cdot \frac{1}{\left(\tilde{R}(\tilde{\Gamma})\right)^{n/2}} \\
&\quad \times \left(2n^2\tilde{P}^2 + 4(n+2)(n-3)A\tilde{R}(\tilde{\Gamma}) - n\tilde{P}\sqrt{n^2\tilde{P}^2 + 4(n^2 - 4)A\tilde{R}(\tilde{\Gamma})}\right)
\end{aligned} \tag{18.182}$$

$$\begin{aligned}
m_0^2 &= \frac{1}{2}n|\gamma| \left(\frac{n}{n+2}\right)^{n/2} \left(\frac{|\gamma|}{\beta}\right)^{n/2} \\
&= \frac{n}{2} \cdot \left(\frac{m_{\tilde{A}}}{\alpha_s^2 m_{\text{pl}}}\right)^n \left(\frac{n}{n+2}\right)^{n/2} \left(\frac{m_{\tilde{A}}}{\alpha_s}\right)^2 \cdot |\tilde{P}| \cdot \left(\frac{|\tilde{P}|}{\tilde{R}(\tilde{\Gamma})}\right)^{n/2}.
\end{aligned} \tag{18.183}$$

Using our simplified model for $\tilde{R}(\tilde{\Gamma})$, $\tilde{P}$ and A one gets

$$\begin{aligned}
m_1^2 &= \left(\frac{m_{\tilde{A}}}{\alpha_s}\right)^2 \left(\frac{m_{\tilde{A}}}{\alpha_s^2 m_{\text{pl}}}\right)^{14} \cdot 2^{-36} \\
&\quad \times \frac{\left(7g(\zeta, \mu) + \sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)}\right)^6 (\mu^2 + 4)^8}{\left(\ln(|\zeta| + \sqrt{\zeta^2 + 1}) + 2\zeta^2 + 1\right)^6 (2\mu^3 + 7\mu^2 + 5\mu + 20)^7} \\
&\quad \times \left(7g(\zeta, \mu)\sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} - 98g^2(\zeta, \mu) - 352h(\zeta, \mu)\right).
\end{aligned} \tag{18.184}$$

Taking $m_{\tilde{A}} \simeq m_{\text{EW}} = 80 \text{ GeV}$ (an electro-weak energy scale) and $m_{\text{pl}} \simeq 2.4 \cdot 10^{18} \text{ GeV}$, $\alpha_s^2 = \alpha_{\text{em}} = \frac{1}{137}$, one gets

$$m_1 \simeq 3.34 \cdot 10^{-105} m_{\text{EW}} G(\zeta, \mu) \tag{18.185}$$

where

$$\begin{aligned}
G(\zeta, \mu) &= \frac{\left(7g(\zeta, \mu) + \sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)}\right)^3 (\mu^2 + 4)^4}{\left(\ln(|\zeta| + \sqrt{\zeta^2 + 1}) + 2\zeta^2 + 1\right)^3 (2\mu^3 + 7\mu^2 + 5\mu + 20)^{7/2}} \\
&\quad \times \left(7g(\zeta, \mu)\sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} - 98g^2(\zeta, \mu) - 352h(\zeta, \mu)\right)^{1/2}.
\end{aligned} \tag{18.186}$$

In the second de Sitter phase one gets

$$m_0^2 = \frac{1}{4} \left(\frac{7}{4} \right)^8 \left(\frac{m_{\tilde{A}}}{\alpha_s^2 m_{\text{pl}}} \right)^{14} \left(\frac{m_{\tilde{A}}^2}{\alpha_s^2} \right) \left(\frac{\mu^2 + 4}{2\mu^3 + 7\mu^2 + 5\mu + 20} \right)^7 \times \left(\frac{g(\zeta, \mu)}{\ln(|\zeta| + \sqrt{\zeta^2 + 1}) + 2\zeta^2 + 1} \right)^8. \quad (18.187)$$

Taking as before for $m_{\tilde{A}}$ an electro-weak energy scale and for $\alpha_s^2 = \frac{1}{137}$, one gets

$$m_0 \cong 10^{-102} m_{\text{EW}} F(\zeta, \mu) \quad (18.188)$$

where

$$F(\zeta, \mu) = \frac{(\mu^2 + 4)^{7/2} (g(\zeta, \mu))^4}{(2\mu^3 + 7\mu^2 + 5\mu + 20)^{7/2} \left(\ln(|\zeta| + \sqrt{\zeta^2 + 1}) + 2\zeta^2 + 1 \right)^4}. \quad (18.189)$$

According to modern ideas a mass of scalar-quintessence particle should be smaller than 10^{-3} eV. For the cosmological constant from the second de Sitter phase is the same for our contemporary epoch, m_0 is also a mass of a scalar quintessence particle for our epoch.

It is interesting to ask what is a number of scalar-quintessence particle per unit volume during both de Sitter phases of the evolution of the Universe (if a quintessence has been deposited as particles which is not so obvious). (It can be deposited in a form of a classical scalar field.) One gets

$$\begin{aligned} n_1 &= \frac{\rho_Q^1}{m_1} = \frac{6H_0^2}{m_1} \\ &= \left(\frac{m_{\tilde{A}}}{\alpha_s} \right) \left(\frac{m_{\tilde{A}}}{\alpha_s m_{\text{pl}}} \right)^{n/2} (2(n+2))^{-(n+4)/4} \cdot 2^5 \\ &\times \frac{n^2 \tilde{P}^2 - \tilde{P} \sqrt{n^2 \tilde{P}^2 + 4(n^2 - 4)A\tilde{R}(\tilde{\Gamma})} + 4(n+2)A\tilde{R}(\tilde{\Gamma})}{\left(2n^2 \tilde{P}^2 + 4(n+2)(n-3)A\tilde{R}(\tilde{\Gamma}) - n\tilde{P} \sqrt{n^2 \tilde{P}^2 + 4(n^2 - 4)A\tilde{R}(\tilde{\Gamma})} \right)^{1/2}} \\ &\times \left(-n\tilde{P} + \sqrt{n^2 \tilde{P}^2 + 4(n^2 - 4)A\tilde{R}(\tilde{\Gamma})} \right)^{(n-2)/4}. \end{aligned} \quad (18.190)$$

In the second de Sitter phase one gets

$$\begin{aligned} n_0 &= \frac{\rho_Q}{m_0} = \frac{6H_1^2}{m_0} \\ &= \left(\frac{m_{\tilde{A}}}{m_{\text{pl}}} \right)^{n/2} \left(\frac{m_{\tilde{A}}}{\alpha_s^n} \right) \left(\frac{n}{n+2} \right)^{(n+4)/4} |\tilde{P}|^{1/2} \left(\frac{|\tilde{P}|}{\tilde{R}(\tilde{\Gamma})} \right)^{n/4}. \end{aligned} \quad (18.191)$$

Using our simplified model with $n = 14$ one gets from (18.190)

$$\begin{aligned}
n_1 = & 2^{-19} \left(\frac{m_{\tilde{A}}}{\alpha_s m_{\text{pl}}} \right)^7 \left(\frac{m_{\tilde{A}}}{\alpha^s} \right) \frac{\left(7g(\zeta, \mu) + \sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} \right)^3}{(2\mu^3 + 7\mu^2 + 5\mu + 20)^{7/2}} \\
& \times \frac{(\mu^2 + 4)^3 \left(98g^2(\zeta, \mu) + (\mu^2 + 4)\sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} + 32H(\zeta, \mu) \right)}{\left(7g(\zeta, \mu)\sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} - 98g^2(\zeta, \mu) - 352h(\zeta, \mu) \right)^{1/2}} \quad (18.192) \\
& \times \left(\ln \left(|\zeta| + \sqrt{\zeta^2 + 1} \right) + 2\zeta^2 + 1 \right)^{-1}
\end{aligned}$$

and from the formula (18.191)

$$\begin{aligned}
n_0 = & \left(\frac{m_{\tilde{A}}}{m_{\text{pl}}} \right)^7 \left(\frac{m_{\tilde{A}}}{\alpha_s^{14}} \right) \left(\frac{7^{9/2}}{2^{17}} \right) \\
& \times \frac{|f(\zeta)|^{1/2}}{\left(\ln \left(|\zeta| + \sqrt{\zeta^2 + 1} \right) + 2\zeta^2 + 1 \right)^{15/2}} \cdot \left(\frac{|f(\zeta)|(\mu^2 + 4)^2}{2\mu^3 + 7\mu^2 + 5\mu + 20} \right)^7. \quad (18.193)
\end{aligned}$$

For the cosmological constant from the second de Sitter phase is the same for our contemporary epoch, n_0 is also a number of scalar quintessence particles for our epoch. Taking as usual $m_{\tilde{A}} = m_{\text{EW}}$ and $\alpha_s^2 = \frac{1}{137}$ one gets

$$n_1 \simeq 9 \cdot 10^{-106} m_{\text{EW}} \cdot K(\zeta, \mu) \quad (18.194)$$

$$n_0 \simeq 6.2 \cdot 10^{-96} m_{\text{EW}} \cdot L(\zeta, \mu), \quad (18.195)$$

where

$$\begin{aligned}
K(\zeta, \mu) = & \frac{\left(7g(\zeta, \mu) + \sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} \right)^3 (\mu^2 + 4)^3}{(2\mu^3 + 7\mu^2 + 5\mu + 20)^{7/2}} \\
& \times \frac{\left(98g^2(\zeta, \mu) + (\mu^2 + 4)\sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} + 32H(\zeta, \mu) \right)}{\left(7g(\zeta, \mu)\sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} - 98g^2(\zeta, \mu) - 352h(\zeta, \mu) \right)^{1/2}}, \quad (18.196)
\end{aligned}$$

$$L(\zeta, \mu) = \frac{|f(\zeta)|^{1/2}}{\left(\ln \left(|\zeta| + \sqrt{\zeta^2 + 1} \right) + 2\zeta^2 + 1 \right)^{15/2}} \cdot \left(\frac{|f(\zeta)|(\mu^2 + 4)^2}{2\mu^3 + 7\mu^2 + 5\mu + 20} \right)^7, \quad (18.197)$$

$f(\zeta)$, $g(\zeta, \mu)$, $h(\zeta, \mu)$, $H(\zeta, \mu)$ are given by the formulae (18.143–145). In this way the masses of scalar particles and their numbers per unit volume in both de Sitter phases and for our contemporary epoch depend on geometric parameters in our theory.

We can connect $\bar{\mu}$ and N from the formula (18.101):

$$\bar{\mu} = \frac{9.8}{N}. \quad (18.198)$$

If $N \simeq 60$, one gets

$$\bar{\mu} = 0.16 \quad (18.199)$$

and

$$5\bar{\mu}^2 = 0.13. \quad (18.200)$$

Using Eq. (18.83) one finds:

$$n_s(K) = 1 \pm 0.13 \Delta \ln K. \quad (18.201)$$

We can also find $\bar{\mu}$ using the formulae (18.113) and (18.113a). One gets

$$\bar{\mu} = \left(\frac{m_{\text{pl}}}{m_{\tilde{A}}} \right)^2 \alpha_s^4 F(\beta, \gamma, \alpha_1) = \left(\frac{m_{\text{pl}}}{m_{\tilde{A}}} \right)^4 F(\tilde{R}(\tilde{\Gamma}), \tilde{\underline{P}}, A). \quad (18.202)$$

Using Eq. (18.83) one also gets

$$n_s(K) = 1 \pm 5 \left(\frac{m_{\text{pl}}}{m_{\tilde{A}}} \right)^8 F^2(\tilde{R}(\tilde{\Gamma}), \tilde{\underline{P}}, A) \Delta \ln K. \quad (18.203)$$

Using our simplified model one gets

$$\begin{aligned} \bar{\mu} = 81.67 \left(\frac{m_{\text{pl}}}{m_{\tilde{A}}} \right)^4 & \frac{7g(\zeta, \mu) + \sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)}}{(\mu^2 + 4)^2(1 + \zeta^2)} \\ & \times \frac{|\zeta| ((5\zeta^2 + 4) \ln 2 - 2\mu^2(1 + \zeta^2))}{\left(98g^2(\zeta, \mu) + (\mu^2 + 4)\sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} + 32H(\zeta, \mu) \right)} \end{aligned} \quad (18.204)$$

and respectively

$$\begin{aligned} n_s(K) = 1 \pm 408.35 & \frac{\left(7g(\zeta, \mu) + \sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} \right)^2}{(\mu^2 + 4)^4(1 + \zeta^2)^2} \\ & \times \frac{|\zeta|^2 ((5\zeta^2 + 4) \ln 2 - 2\mu^2(1 + \zeta^2))^2 \Delta \ln K}{\left(98g^2(\zeta, \mu) + (\mu^2 + 4)\sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} + 32H(\zeta, \mu) \right)^2}. \end{aligned} \quad (18.205)$$

It is easy to see that we can get $n_s(K)$ close to 1 (a flat power spectral function) using parameters μ and ζ in such a way that

$$(5\zeta^2 + 4) \ln 2 - 2\mu^2(1 + \zeta^2)$$

is close to zero. One can express $\cos \varphi$ (see Eq. (18.5)) in terms of parameters of the theory and get

$$\begin{aligned} \cos \varphi = & - \left(\frac{m_{\tilde{A}}}{\alpha_s^2 m_{\text{pl}}} \right)^{n/2} \frac{P_1^{n/4}}{3 \cdot 2^{(n+8)/4} (n+2)^{n/4} \left(\tilde{R}(\tilde{\Gamma}) \right)^{n/4}} \\ & \times \frac{40A \left(\frac{m_{\tilde{A}}}{\alpha_s^2 m_{\text{pl}}} \right)^n m_{\tilde{A}}^2 P_1^{n/2} + 9B\alpha_s^4 2^{n/2} (n+2)^{n/2} \left(\tilde{R}(\tilde{\Gamma}) \right)^{n/2}}{5A \left(\frac{m_{\tilde{A}}}{\alpha_s^2 m_{\text{pl}}} \right)^n m_{\tilde{A}}^2 P_1^{n/2} - 3\alpha_s^4 B 2^{n/2} (n+2)^{n/2} \left(\tilde{R}(\tilde{\Gamma}) \right)^{n/2}}, \end{aligned} \quad (18.206)$$

where

$$P_1 = -n\tilde{\underline{P}} + \sqrt{n^2\tilde{\underline{P}}^2 + 4(n^2 - 4)A\tilde{R}(\tilde{\Gamma})}. \quad (18.207)$$

For the constant a (see Eq. (18.6)) one gets

$$a = \alpha_s^2 \left(\frac{\sqrt{B}}{m_{\tilde{A}}} \right) \left(\frac{\alpha_s^2 m_{\text{pl}}}{m_{\tilde{A}}} \right)^{n/2} \frac{2^{n/4} (n+2)^{n/4} \left(\tilde{R}(\tilde{\Gamma}) \right)^{n/4}}{\sqrt{A} \left(-n\tilde{\underline{P}} + \sqrt{n^2\tilde{\underline{P}}^2 + 4(n^2 - 4)A\tilde{R}(\tilde{\Gamma})} \right)^{n/4}} \quad (18.208)$$

(see Refs [129], [130]).

19. Properties of a Quintessence

In this section we consider several consequences of the Nonsymmetric Kaluza–Klein (Jordan–Thiry) Theory. We consider a value of a mass of quintessence particle, several interesting relations among energy scales, radiation density in the second de Sitter phase. We find a spatial dependence of a quintessence field (and an effective gravitational constant G_{eff}) in a case of spherical-statical symmetry, cylindrical-statical symmetry, flat-static symmetry. We find a time dependences of a quintessence field (with no spatial dependence). We get a solution for a quintessence field (a travelling wave) and two-dimensional wave solution applying those solutions for G_{eff} . We propose some kind of statistical approach to our results. We calculate a speed of sound in a quintessence.

Let us consider a value of mass of a quintessence particle (a scalar particle) (18.183) obtained from Nonsymmetric Kaluza–Klein (Jordan–Thiry) Theory:

$$m_0 = \sqrt{\frac{n}{2}} \left(\frac{n}{n+2} \right)^{n/4} |\gamma|^{1/2} \left(\frac{|\gamma|}{\beta} \right)^{n/4}. \quad (19.1)$$

The value of this mass has been obtained by this particle during the second de Sitter phase. Moreover during our contemporary epoch it is the same. The parameters n , γ and β are defined in Sec. 14 (Eq. (14.33)). During the second de Sitter phase a cosmological constant has been calculated and one finds

$$\lambda_{c0} = 6H_1^2 = \frac{2|\gamma|^{(n+2)/2} n^{n/2}}{\beta^{n/2} (n+2)^{(n+2)/2}}. \quad (19.2)$$

The value of a cosmological constant remains the same during our contemporary epoch. H_1 is a Hubble constant during the second de Sitter phase (see Eq. (14.79)).

Thus for our contemporary epoch one gets

$$m_0 = \frac{1}{2} \sqrt{n(n+2)} \sqrt{\lambda_{c0}}. \quad (19.3)$$

In this way we can connect the value of a cosmological constant of our contemporary epoch to the value of a mass of a quintessence particle. From recent observational data we get

$$\lambda_{c0} = \Lambda = 10^{-52} \frac{1}{\text{m}^2}. \quad (19.4)$$

Moreover in order to get a correct dimension for a mass we should add a factor with $\hbar$ and c (now we abandon the system of units with $\hbar = c = 1$).

One gets

$$m_0 = \frac{1}{2} \frac{\hbar}{c} \sqrt{n(n+2)} \sqrt{\lambda_{c0}}. \quad (19.3a)$$

Using the value of λ_{c0} one finally gets

$$m_0 \simeq \sqrt{n(n+2)} \cdot 0.17 \cdot 10^{-39} \text{ g} \quad (19.5)$$

or

$$m_0 \simeq \sqrt{n(n+2)} \cdot 0.95 \cdot 10^{-5} \text{ eV}. \quad (19.5a)$$

For example, if we take $n = 14 (= \dim G_2)$, one finally gets

$$m_0 \simeq 14.2 \cdot 10^{-5} \text{ eV}. \quad (19.6)$$

This value is bigger than that considered by different authors. Moreover, still sufficiently small. The particle interacts only gravitationally and because of this it is undetectable by using known experimental methods. Taking a density of dark energy as 0.7 of a critical density,

$$\rho_c = 1.88 h^2 \cdot 10^{-29} \frac{\text{g}}{\text{cm}^3}, \quad (19.7)$$

one gets a number of quintessence particles per unit volume

$$\bar{n} = \frac{h^2}{\sqrt{n(n+2)}} \cdot 1.31 \cdot 10^{10} \frac{1}{\text{cm}^3} \quad (19.8)$$

where h is a dimensionless Hubble constant $0.7 < h < 1$. Taking $n = 14$ and $h = 0.7$ one finally gets

$$\bar{n} = 4 \cdot 10^8 \frac{1}{\text{cm}^3} \quad (19.8a)$$

which is many orders of magnitude smaller than Loschmidt number. Thus a gas of quintessence particles is not so dense from the point of view of our earth conditions. However, if this number of particles per unit volume is considered in a container of size 200 Mpc, the gas can be considered as extremely dense.

In order to settle—is this gas dense or not—we should calculate a mean scattering length. The scattering cross-section for a quintessence particle

$$\sigma = \frac{1}{\lambda_{c0}} = 10^{52} \text{ m}^2. \quad (*)$$

A mean scattering length

$$l = \frac{1}{\sigma n} \quad (**)$$

where n is a number of quintessence particles per unit volume (Eq. (19.8a)).

One gets

$$l = 10^{-60} \text{ m}. \quad (***)$$

It means that a gas of quintessence particles is extremely dense (if we apply the Knudsen criterion—a gas is dense if $l \ll L$, where L is the size of the container) even in the Solar System.

Let us consider the Eq. (16.170) which connects several scales of energy and gives an account of the smallness of gravitational interactions in our theory. We rewrite this equation in the form

$$\left(\frac{m_{\text{pl}}}{m_{\text{EW}}}\right) \left(\frac{m_{\tilde{A}}}{m_{\text{EW}}}\right)^{n_1} = \left(\frac{n|\gamma|}{(n+2)\beta}\right)^{(n+2)(n_1+2)/2} \quad (19.9)$$

where m_{EW} is an electro-weak interactions energy scale. Taking

$$\begin{aligned} m_{\tilde{A}} &= m_{\text{EW}} \\ n_1 &= 2 \quad (M = S^2) \\ n &= 14 \quad (H = G2) \end{aligned}$$

one gets

$$\frac{m_{\text{pl}}}{m_{\text{EW}}} = \left(\frac{7}{8}\kappa\right)^{24} \quad (19.10)$$

where

$$\kappa = \frac{|\gamma|}{\beta} \quad (19.11)$$

and eventually one gets

$$\kappa = \left(\frac{m_{\text{pl}}}{m_{\text{EW}}}\right)^{1/24} \cdot \frac{8}{7}. \quad (19.12)$$

Taking $m_{\text{EW}} \simeq 80 \text{ GeV}$ and $m_{\text{pl}} \simeq 2.4 \cdot 10^{18} \text{ GeV}$ one gets

$$\kappa \simeq 6.19 \quad (19.13)$$

which is very reasonable for it establishes a relation between two cosmological terms γ and β as of the same order. Simultaneously this is a consistency condition for our model with energy scales (the 6-dimensional Planck's mass is equal to m_{EW}). In this way we can calculate a mass of a quintessence particle for our contemporary epoch from Eq. (19.1)

$$m_0 = 2 \cdot 10^{15} \text{ eV} \cdot |\underline{\tilde{P}}|^{1/2}. \quad (19.14)$$

This gives us an estimation for $|\underline{\tilde{P}}|$ and $\tilde{R}(\tilde{\Gamma})$:

$$\tilde{R}(\tilde{\Gamma}) = \frac{1}{\kappa} \left(\frac{m_{\text{EW}}}{m_{\text{pl}}\alpha_{\text{em}}}\right)^2 |\underline{\tilde{P}}| \quad (19.15)$$

where $\alpha_{\text{em}} = \frac{1}{137}$ is a fine coupling constant. Using Eq. (19.13) and values of m_{pl} and m_{EW} one gets

$$\tilde{R}(\tilde{\Gamma}) = 24.75 \cdot 10^{-31} |\underline{\tilde{P}}|. \quad (19.16)$$

From Eq. (19.14) and Eq. (19.6) one gets

$$|\underline{\tilde{P}}| = 5 \cdot 10^{-39} \quad (19.17)$$

and

$$\tilde{R}(\tilde{\Gamma}) \cong 1.2 \cdot 10^{-68}. \quad (19.18)$$

Moreover in our simplified theory we have

$$\tilde{R}(\tilde{\Gamma}) = \frac{2(2\mu^3 + 7\mu^2 + 5\mu + 20)}{(\mu^2 + 4)^2} \quad (19.19)$$

and μ should be very close to the root of the polynomial

$$W(\mu) = 2\mu^3 + 7\mu^2 + 5\mu + 20. \quad (19.20)$$

From (19.18) and (19.19) one gets

$$2\mu^3 + 7\mu^2 + 5\mu + 20 \cong 1.3 \cdot 10^{-66}. \quad (19.21)$$

Thus μ is very close to the 70-digit approximation of the root of the polynomial (19.20) ($W(\bar{\mu}) = 0.1 \cdot 10^{-67}$). Due to this we can control the cosmological terms.

In the case of $|\underline{\tilde{P}}|$ we have the formula (8.64) and one gets

$$\underline{\tilde{P}}(\zeta_0 + \varepsilon) \cong \underline{\tilde{P}}(\zeta_0) + \frac{d\underline{\tilde{P}}}{d\zeta}(\zeta_0)\varepsilon \quad (19.22)$$

$$\underline{\tilde{P}}(\zeta_0 = \pm 1.38 \dots) = 0 \quad (19.23)$$

$$\left| \frac{d\underline{\tilde{P}}}{d\zeta}(|\zeta_0| = 1.38 \dots) \right| = 25. \quad (19.24)$$

Thus one gets from (19.17) and (19.23–24)

$$\varepsilon \simeq 2 \cdot 10^{-40}. \quad (19.25)$$

Thus we need an approximation of ζ_0 up 40-digit arithmetics. We see that cosmological terms coming from the Nonsymmetric Kaluza–Klein (Jordan–Thiry) Theory are very small, but not zero, and that they are easily controllable by μ and ζ parameters.

Let us consider a self-interaction potential for a quintessence field for our contemporary epoch (which is the same as for the second de Sitter phase). One gets from

cosmological terms $\lambda_{c0}(\Psi)$

$$\lambda_{c0}(\Psi) = -\frac{1}{2} (\beta e^{2\Psi} - |\gamma|) e^{n\Psi} \quad (19.26)$$

$$\begin{aligned} U(q_0) = & -\frac{|\gamma|}{2(n+2)} \left(\frac{n|\gamma|}{(n+2)\beta} \right)^{\frac{n}{2}} \exp \left(\frac{nm_{\text{pl}}}{2\sqrt{2\pi|\overline{M}|}} q_0 \right) \\ & \times \left(n \exp \left(\frac{m_{\text{pl}}}{\sqrt{2\pi|\overline{M}|}} q_0 \right) - (n+2) \right) \end{aligned} \quad (19.27)$$

or

$$U(q_0) = -\frac{1}{4} \lambda_{c0} \exp \left(\frac{nm_{\text{pl}}}{2\sqrt{2\pi|\overline{M}|}} q_0 \right) \left(n \exp \left(\frac{m_{\text{pl}}}{\sqrt{2\pi|\overline{M}|}} q_0 \right) - (n+2) \right) \quad (19.27a)$$

where λ_{c0} is a cosmological constant for our contemporary epoch and

$$\Psi = \Psi_0 + \frac{m_{\text{pl}}}{2\sqrt{2\pi|\overline{M}|}} q_0 = \Psi_0 + \overline{\beta} q_0 = \Psi_0 + \varphi. \quad (19.28)$$

For λ_{c0} is very small ($10^{-52} \frac{1}{\text{m}^2}$), this interaction is very small. For $n = 14 (= \dim H = \dim G2)$ one gets

$$U(q_0) = -\frac{1}{2} \lambda_{c0} \exp \left(\frac{7m_{\text{pl}}}{\sqrt{2\pi|\overline{M}|}} q_0 \right) \left(7 \exp \left(\frac{m_{\text{pl}}}{\sqrt{2\pi|\overline{M}|}} q_0 \right) - 8 \right). \quad (19.29)$$

Even in the potential $U(q_0)$ we have exponential terms, the strength of the interactions is negligible for small value of q_0 . The interesting point in our theory is the effective gravitational constant (depending on a scalar field Ψ). Let us describe it by a quintessence field q_0 . One gets

$$G_{\text{eff}} = G_0 e^{-(n+2)\Psi}. \quad (19.30)$$

After some calculations one finds

$$G_{\text{eff}} = G_0 \left(\frac{(n+2)\beta}{n|\gamma|} \right)^{n+2} \exp \left(-\frac{n+2}{2} \frac{m_{\text{pl}}}{\sqrt{2\pi|\overline{M}|}} q_0 \right) \quad (19.31)$$

or

$$G_{\text{eff}} = \frac{G_0}{\lambda_{c0}^2} \cdot \frac{4(n+2)^2 \beta^2}{n^2} \exp \left(-\frac{n+2}{2} \frac{m_{\text{pl}}}{\sqrt{2\pi|\overline{M}|}} q_0 \right). \quad (19.31a)$$

It is easy to see that the rôle of a Newton's constant G_N is played by

$$G_N = \frac{G_0}{\lambda_{c0}^2} \left(\frac{2(n+2)\beta}{n} \right)^2. \quad (19.32)$$

This is the reason that we put a constant G_0 in the formula (19.30).

One gets

$$G_0 = G_N \lambda_{c0}^2 \left(\frac{n}{2(n+2)\beta} \right)^2. \quad (19.33)$$

For

$$\beta = \alpha_s^2 m_{\text{pl}}^2 \tilde{R}(\tilde{\Gamma}) = \alpha_s^2 \frac{\tilde{R}(\tilde{\Gamma})}{l_{\text{pl}}^2} \quad (19.34)$$

one gets

$$G_N = \frac{G_0 \alpha_s^4}{(\lambda_{c0} l_{\text{pl}}^2)^2} \left(\frac{2(n+2)\tilde{R}(\tilde{\Gamma})}{n} \right)^2 \quad (19.35)$$

or

$$G_0 = G_N \left(\frac{n}{2(n+2)\tilde{R}(\tilde{\Gamma})} \right)^2 \left(\frac{\lambda_{c0} l_{\text{pl}}^2}{\alpha_s^2} \right)^2. \quad (19.36)$$

Let us connect to G_0 a new Planck's mass m_{pl}^0 . In terms of this mass and ordinary m_{pl} mass one gets

$$m_{\text{pl}}^0 = m_{\text{pl}} \frac{2(n+2)\tilde{R}(\tilde{\Gamma})}{n} \frac{\alpha_s^2}{\lambda_{c0} l_{\text{pl}}^2} \quad (19.37)$$

or

$$m_{\text{pl}} = m_{\text{pl}}^0 \frac{n}{2(n+2)\tilde{R}(\tilde{\Gamma})} \frac{\lambda_{c0} l_{\text{pl}}^2}{\alpha_s^2}. \quad (19.38)$$

If we take our simplified model with $\tilde{R}(\tilde{\Gamma})$ given by Eq. (19.19)

$$\alpha_s^2 = \alpha_{\text{em}} = \frac{1}{137}, \quad n = 14,$$

we get

$$m_{\text{pl}}^0 = m_{\text{pl}} \frac{32(2\mu^3 + 7\mu^2 + 5\mu + 20)}{7(\mu^2 + 4)^2} \frac{\alpha_{\text{em}}}{\lambda_{c0} l_{\text{pl}}^2} \quad (19.39)$$

or

$$G_0 = G_N \left(\frac{7(\mu^2 + 4)^2}{32(2\mu^3 + 7\mu^2 + 5\mu + 20)} \right)^2 \left(\frac{\lambda_{c0} l_{\text{pl}}^2}{\alpha_{\text{em}}} \right)^2. \quad (19.40)$$

Moreover we can connect unknown constant G_0 (m_{pl}^0) with an energy scale of electro-weak interactions m_{EW} in a way suggested and developed here (Sec. 16).

It is enough to use the formula (19.10) and we get

$$m_{\text{pl}}^0 = m_{\text{EW}} \left(\frac{7\kappa}{8} \right)^{24} \frac{32(2\mu^3 + 7\mu^2 + 5\mu + 20)}{7(\mu^2 + 4)^2} \frac{\alpha_{\text{em}}}{\lambda_{c0} l_{\text{pl}}^2}. \quad (19.41)$$

If we take for λ_{c0} the known value of the cosmological constant (Eq. (19.4)) for $l_{\text{pl}} = 4 \cdot 10^{-32}$ cm and $\kappa = 6.19$ we get

$$m_{\text{pl}}^0 = 8.12 \cdot 10^{132} \cdot \frac{2\mu^3 + 7\mu^2 + 5\mu + 20}{(\mu^2 + 4)^2} m_{\text{EW}}. \quad (19.42)$$

Moreover we see that μ must be very close to the root of the polynomial $2\mu^3 + 7\mu^2 + 5\mu + 20$ (see (16.199)) and we get

$$m_{\text{pl}}^0 = 2 \cdot 10^{130} (2\mu^3 + 7\mu^2 + 5\mu + 20) m_{\text{EW}}. \quad (19.43)$$

If we take 70-digit approximation of the root of the polynomial (19.20) and considered value of $W(\mu)$ (Eq. (19.18)) obtained here, we get

$$m_{\text{pl}}^0 = 4.6 \cdot 10^{64} m_{\text{EW}} \quad (19.44)$$

or

$$m_{\text{pl}}^0 = 4.6 \cdot 10^{64} \frac{m_{\text{EW}}}{m_{\text{pl}}} m_{\text{pl}} = 1.5 \cdot 10^{48} m_{\text{pl}} = 3.2 \cdot 10^{43} \text{g} = 1.6 \cdot 10^{10} M_{\odot} \quad (19.45)$$

where $M_{\odot} = 1.99 \cdot 10^{33}$ g is a Solar mass. (This is a mass of our Galaxy.)

Moreover the present critical density of matter in the Universe is

$$\rho_{c,0} = 2.775 h^{-1} \times 10^{11} \frac{M_{\odot}}{(h^{-1} \text{Mpc})^3}. \quad (19.46)$$

So we can find a volume containing m_{pl}^0 . One gets

$$V_0 = \frac{m_{\text{pl}}^0}{\rho_{c,0}} = 5 \cdot 10^{-3} h^2 (\text{Mpc})^3. \quad (19.47)$$

Thus the size of this volume is of order

$$L_0 \simeq \sqrt[3]{h^2} 0.17 \text{Mpc}. \quad (19.48)$$

Taking under consideration the fact that our calculations are quite rough (for example κ could be bigger a little, or $W(\tilde{\mu})$ a little bigger), we come to the conclusion that m_{pl}^0 is of order of the mass of our visible Universe (a volume of size of 10^4 Mpc, see the second point of Ref. [129]). In this way it seems natural to suppose that

$$m_{\text{pl}}^0 = M_U \quad (19.49)$$

where M_U is the total mass of the visible Universe. However, this intriguing conjecture demands many investigations.

Moreover it is in a real spirit of Dirac's large number hypothesis and can give a link between cosmology and fundamental interactions theory. Thus if we suppose (19.49) we get

$$G_0 = G_U = \frac{1}{M_U^2}. \quad (19.50)$$

Let us come back to the equation for an effective gravitational constant (Eq. (19.32)):

$$G_{\text{eff}} = G_N \exp \left(-\frac{n+2}{2} \frac{m_{\text{pl}}}{\sqrt{2\pi|\overline{M}|}} q_0 \right). \quad (19.51)$$

q_0 —a quintessence scalar field—possesses a mass and because of this it has a finite range with Yukawa type behaviour

$$q_0 = \frac{\alpha}{R} \exp \left(-\frac{R}{r_0} \right) \quad (19.52)$$

where

$$r_0 = \frac{2}{\sqrt{n(n+2)}\sqrt{\lambda_{c0}}} \quad \text{and } \alpha \text{ is a positive constant} \quad (19.53)$$

or taking for $n = 14$ and for λ_{c0} Eq. (19.4),

$$r_0 = 3 \cdot 10^{23} \text{ m} = 10 \text{ Mpc}. \quad (19.54)$$

The formula (19.52) is valid for $R \simeq r_0$. Thus we cannot observe scalar gravitational radiation from closed binary sources and the quadrupole radiation formula is satisfied for a gravitational field.

Moreover for $R < r_0$ we should take under consideration a full self-interaction potential of a field q_0 (Eq. (19.29)). Moreover, because of the constant λ_{c0} in front of the formula (19.29) this is negligible. Because of this we can repeat some considerations from Sections 10 and 14 coming back to the field Ψ and consider different sources of mass for this field due to interaction with the matter. In this way in both cases

$$G_{\text{eff}} = G_N. \quad (19.55)$$

However, we can expect some small effects on short distances.

Let us notice that in the previous approximation we consider a weak field, it means $|q_0| \ll 1$. This is different from small field. A small field considered below is such that q_0 is small in a usual sense. It is negative. Moreover it can be big in the sense of the absolute value.

Let us calculate a radiation density in the moment when the second de Sitter phase starts. It means the radiation released after the phase transition. One gets

$$\rho_r = \lambda_{c1}(\Psi_1) - \lambda_{c0}(\Psi_0) = \lambda_{c0}(\Psi_0) \left(\frac{H_0^2}{H_1^2} - 1 \right). \quad (19.56)$$

Using Eqs (18.103a) and (18.111a) one gets

$$\begin{aligned} \rho_r &= \lambda_{c0}(\Psi_0) \\ &\times \left(\frac{n^2 |\tilde{P}|^2 + |\tilde{P}| \sqrt{n^2 |\tilde{P}|^2 + 4(n^2 - 4)A\tilde{R}(\tilde{I})} + 4A\tilde{R}(\tilde{I})(n+2)}{2^{(n+2)/2} |\tilde{P}|^{(n+2)/2} n^{n/2}} \right. \\ &\quad \left. \times \left(n |\tilde{P}| + \sqrt{n^2 |\tilde{P}|^2 + 4(n^2 - 4)A\tilde{R}(\tilde{I})} \right)^{(n-2)/2} - 1 \right), \end{aligned} \quad (19.57)$$

$$\lambda_{c0}(\Psi_0) = 6H_1^2 = 2 \left(\frac{m_{\tilde{A}}}{m_{\text{pl}}} \right)^n \frac{m_{\tilde{A}}^2}{\alpha_s^{2(n+1)}} \cdot \frac{|\tilde{P}|^{(n+2)/2} n^{n/2}}{(n+2)^{(n+2)/2} (\tilde{R}(\tilde{I}))^{n/2}}. \quad (19.58)$$

The density ρ_r is of course an effective radiation density, because we adsorbe into ρ_r a factor with an effective gravitational constant.

In our simplified model we get

$$\begin{aligned} \rho_r &= \Lambda \left(\frac{98g^2(\zeta, \mu) + (\mu^2 + 4)\sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} + 32H(\zeta, \mu)}{g^8(\zeta, \mu)(\mu^2 + 4)^2 \cdot 2^{11} \cdot 7^{14}} \right. \\ &\quad \times \left(7g(\zeta, \mu) + \sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} \right)^6 \\ &\quad \left. \times \left(\ln \left(|\zeta| + \sqrt{\zeta^2 + 1} \right) + 2\zeta^2 + 1 \right)^6 - 1 \right), \end{aligned} \quad (19.59)$$

$$\begin{aligned} \Lambda &= \left(\frac{m_{\tilde{A}}}{m_{\text{pl}}} \right)^{14} \cdot \frac{m_{\tilde{A}}^2}{\alpha_s^{36}} \cdot \frac{7^{14}}{2^{24}} \\ &\times \frac{g^8(\zeta, \mu)(\mu^2 + 4)^8}{(2\mu^3 + 7\mu^2 + 5\mu + 20)^7 \left(\ln \left(|\zeta| + \sqrt{\zeta^2 + 1} \right) + 2\zeta^2 + 1 \right)^8}. \end{aligned} \quad (19.60)$$

In the formulas (19.59–60) we should put $m_{\tilde{A}} = m_{\text{EW}}$ and $\alpha_s^2 = \alpha_{\text{em}} = \frac{1}{137}$. Let us notice that Λ is our contemporary cosmological constant given by (19.4). It gives a scale for ρ_r .

The functions $g(\zeta, \mu)$, $h(\zeta, \mu)$, $H(\zeta, \mu)$ are given by Eqs (18.143), (18.144a), (18.144b), (18.145). The numerical factor in front of Λ can be calculated and we get

$$\Lambda = 1.04 \cdot 10^{-178} m_{\text{EW}}^2 a(\zeta, \mu) \quad (19.61)$$

where

$$a(\zeta, \mu) = \frac{g^8(\zeta, \mu)(\mu^2 + 4)^8}{(2\mu^3 + 7\mu^2 + 5\mu + 20)^7 \left(\ln \left(|\zeta| + \sqrt{\zeta^2 + 1} \right) + 2\zeta^2 + 1 \right)^8}. \quad (19.62)$$

The calculated radiation density evolves in time according to the theory developed in Sec. 14.

In order to find some influence of q_0 (quintessence) field on the value of the effective gravitational constant we consider a field equation for the scalar q_0 field in empty space. One gets

$$\left(\frac{\partial^2 q_0}{\partial t^2} - \vec{\nabla}^2 q_0\right) + \tilde{\varepsilon} \bar{\alpha} \exp(n \bar{\beta} q_0) (\exp(2 \bar{\beta} q_0) - 1) = 0 \quad (19.63)$$

where

$$\bar{\alpha} = \frac{\lambda_{c0} n(n+2) m_{\text{pl}}^2}{16\pi \cdot \bar{M}} \quad (19.64)$$

$$\bar{\beta} = \frac{m_{\text{pl}}}{2\sqrt{2\pi|\bar{M}|}} \quad (19.65)$$

$$\tilde{\varepsilon} = \text{sgn } \bar{M}, \quad \tilde{\varepsilon}^2 = 1. \quad (19.66)$$

Let us consider a static, spherically symmetric case. In the spherical coordinates one gets

$$0 = \frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{dq_0}{dr} \right) - \tilde{\varepsilon} \bar{\alpha} \exp(n \bar{\beta} q_0) (\exp(2 \bar{\beta} q_0) - 1) \quad (19.67)$$

where $q_0 = q_0(r)$ is a function of r only. In order to treat this equation it is easier to come back to the old variable $\varphi = \bar{\beta} q_0$ (see Eq. (19.28)). One gets

$$\frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{d\varphi}{dr} \right) - \frac{1}{4} \tilde{\varepsilon} \bar{\lambda}_{c0} \exp(n\varphi) (\exp(2\varphi) - 1) = 0. \quad (19.68)$$

We change the independent variable r into τ

$$\frac{1}{\tau^2} \frac{d}{d\tau} \left(\tau^2 \frac{d\varphi}{d\tau} \right) - \tilde{\varepsilon} \exp(n\varphi) (\exp(2\varphi) - 1) = 0 \quad (19.69)$$

where

$$r = \frac{2}{\sqrt{\bar{\lambda}_{c0}}} \tau \quad (19.70)$$

and

$$\bar{\lambda}_{c0} = \frac{n(n+2)\lambda_{c0}}{8\pi\bar{M}} m_{\text{pl}}^2. \quad (19.71)$$

We consider Eq. (19.69) in two regions:

- 1) for small fields φ ,
- 2) for large fields φ .

In the first region we get

$$\frac{1}{\tau^2} \frac{d}{d\tau} \left(\tau^2 \frac{d\varphi}{d\tau} \right) + \tilde{\varepsilon} e^{n\varphi} = 0. \quad (19.72)$$

In the second region we get

$$\frac{1}{\tau^2} \frac{d}{d\tau} \left(\tau^2 \frac{d\varphi}{d\tau} \right) - \tilde{\varepsilon} e^{(n+2)\varphi} = 0. \quad (19.73)$$

Let us notice that both equations have similar nature and can be reduced to the equation

$$\frac{1}{x^2} \frac{d}{dx} \left(x^2 \frac{dy}{dx} \right) + \varepsilon \tilde{\varepsilon} e^y = 0 \quad (19.74)$$

where in the first region $\varepsilon = 1$,

$$y = n\varphi, \quad (19.75)$$

$$x = \sqrt{n} \tau, \quad (19.76)$$

and in the second region $\varepsilon = -1$,

$$y = (n+2)\varphi, \quad (19.77)$$

$$x = \sqrt{n+2} \tau, \quad (19.78)$$

We can transform (19.74) into

$$x \frac{d^2 y}{dx^2} + 2 \frac{dy}{dx} + \varepsilon \tilde{\varepsilon} x e^y = 0 \quad (19.79)$$

which is the celebrated Emden-Fowler equation known in the theory of gaseous spheres (see [131]). Let us notice that the first region (small fields) means large distances and the second region (large fields) means small distances.

In this way we should consider Eq. (19.79) in the region of small and large x . In the case of $\varepsilon \tilde{\varepsilon} = 1$ the equation (19.74) has an exact solution

$$y = \ln \left(\frac{2}{x^2} \right). \quad (19.80)$$

Let us apply this to both regions (remembering that $\tilde{\varepsilon}$ in both cases has a different sign).

One gets in the first region

$$q_0 = - \frac{2\sqrt{2\pi|\overline{M}|}}{m_{\text{pl}} n} \ln \left(\frac{r}{\frac{2\sqrt{2}}{\sqrt{n\overline{\lambda}_{c0}}}} \right) \quad (19.81)$$

and

$$G_{\text{eff}} = G_N \left(\frac{r}{\frac{2\sqrt{2}}{\sqrt{n\overline{\lambda}_{c0}}}} \right)^{(n+2)/n}. \quad (19.82)$$

In the second region

$$q_0 = -\frac{2\sqrt{2\pi|\overline{M}|}}{m_{\text{pl}}(n+2)} \ln \left(\frac{r}{\frac{2\sqrt{2}}{\sqrt{(n+2)\overline{\lambda}_{c0}}}} \right) \quad (19.83)$$

and

$$G_{\text{eff}} = G_N \left(\frac{r}{\frac{2\sqrt{2}}{\sqrt{(n+2)\overline{\lambda}_{c0}}}} \right). \quad (19.84)$$

In this way we get an interesting prediction for the behaviour of the strength of gravitational interactions. In this very special solution G_{eff} is going to zero if $r \rightarrow 0$ and to infinity if $r \rightarrow \infty$.

Let us come to Eq. (19.79) supposing $\varepsilon\tilde{\varepsilon} = 1$. Thus we get

$$x \frac{d^2 y}{dx^2} + 2 \frac{dy}{dx} + x e^y = 0. \quad (19.85)$$

Using an exact solution (19.80) we write

$$y = y_1 + \tilde{y} \quad (19.86)$$

and consider Eq. (19.85) for large x .

In this way we get an approximate solution (given by Chandrasekhar [131])

$$y = \ln \left(\frac{2}{\eta^2} \right) + \frac{A}{\sqrt{\eta}} \cos \left(\frac{\sqrt{7}}{2} \ln \eta \right) - 2 \ln \bar{\delta}, \quad |A| \ll 1, \quad (19.87)$$

where $\eta = \frac{x}{\bar{\delta}}$, A and $\bar{\delta}$ are integration constants, $\bar{\delta} > 0$. In this way we get in the first region

$$G_{\text{eff}} = \overline{G}_{\text{eff}} \cdot \exp \left(-\frac{\overline{A}}{\sqrt{\eta}} \cos \left(\frac{\sqrt{7}}{2} \ln \eta \right) \right) \quad (19.88a)$$

where $\overline{A}$ is a constant ($|\overline{A}| \ll 1$) and

$$\eta = \frac{\sqrt{n\overline{\lambda}_{c0}} \bar{\delta}}{2\sqrt{2}} r, \quad (19.88b)$$

$\overline{G}_{\text{eff}}$ is given by the formula (19.82).

In the second region

$$G_{\text{eff}} = \overline{\overline{G}}_{\text{eff}} \cdot \exp \left(-\frac{\overline{\overline{A}}}{\sqrt{\eta}} \cos \left(\frac{\sqrt{7}}{2} \ln \eta \right) \right) \quad (19.88c)$$

where $\overline{\overline{A}}$ is a constant ($|\overline{\overline{A}}| \ll 1$), $\overline{\overline{G}}_{\text{eff}}$ is given by the formula (19.84) and

$$\eta = \frac{\sqrt{(n+2)\overline{\lambda}_{c0}}\overline{\delta}}{2\sqrt{2}} r. \quad (19.88d)$$

In this way we have very interesting non-Newtonian behaviour of G_{eff} for large distances. Let us notice that the length scale is completely arbitrary, because it is given by an integration constant $\overline{\delta}$.

Let us consider Eq. (19.63) in Cartesian coordinates supposing flat symmetry for a quintessence field $q_0 = q_0(z, t)$ (nonstatic). One gets

$$\left(\frac{\partial^2 q_0}{\partial t^2} - \frac{\partial^2 q_0}{\partial z^2} \right) - \tilde{\varepsilon} \overline{\alpha} \exp(n\overline{\beta} q_0) (\exp(2\overline{\beta} q_0) - 1) = 0. \quad (19.89)$$

Let us change dependent and independent variables to ξ, η, φ :

$$z = \frac{2}{\sqrt{\overline{\lambda}_{c0}}} \xi \quad (19.90)$$

$$t = \frac{2}{\sqrt{\overline{\lambda}_{c0}}} \eta \quad (19.91)$$

$$\varphi = \overline{\beta} q_0. \quad (19.92)$$

One gets

$$\left(\frac{\partial^2 \varphi}{\partial \eta^2} - \frac{\partial^2 \varphi}{\partial \xi^2} \right) - \tilde{\varepsilon} e^{n\varphi} (e^{2\varphi} - 1) = 0. \quad (19.93)$$

Eq. (19.93) is an equation for flat scalar (quintessence) waves in our theory. Let us consider it for large and small field φ (as before).

In this way one gets the equation

$$\left(\frac{\partial^2 y}{\partial T^2} - \frac{\partial^2 y}{\partial x^2} \right) + \varepsilon \tilde{\varepsilon} e^y = 0, \quad (19.94)$$

where in the region of small field φ we have $\varepsilon = 1$ and

$$y = n\varphi \quad (19.95)$$

$$x = \sqrt{n} \xi \quad (19.96)$$

$$T = \sqrt{n} \eta \quad (19.97)$$

and in the region of large field φ , $\varepsilon = -1$ and

$$y = (n+2)\varphi \quad (19.98)$$

$$x = \sqrt{n+2} \xi \quad (19.99)$$

$$T = \sqrt{n+2} \eta. \quad (19.100)$$

Eq. (19.94) is the famous Liouville equation which can be transformed via a Bäcklund transformation into a two-dimensional wave equation and afterwards solved exactly. The general solution depends on two arbitrary functions f and g of one variable, sufficiently regular. It is possible to consider several problems for this equation: Cauchy initial problem, Darboux problem and Goursat problem.

The general solution of (19.94) looks like ($\varepsilon\tilde{\varepsilon} = -1$)

$$y(T, x) = \ln \left[\frac{2g'(x-T)f'(x+T)}{(g(x-T) + f(x+T))^2} \right] \quad (19.101)$$

where g' and f' are derivatives of g and f .

Thus one gets in the first region (small field)

$$q_0(t, z) = \frac{2\sqrt{2\pi\overline{M}}}{m_{\text{pl}}n} \cdot \ln \left[\frac{2g'(\frac{z-t}{a})f'(\frac{z+t}{a})}{(g(\frac{z-t}{a}) + f(\frac{z+t}{a}))^2} \right], \quad (19.102)$$

$$a = \frac{2\sqrt{2}}{\sqrt{n\overline{\lambda}_{c0}}}, \quad (19.103)$$

and

$$G_{\text{eff}} = G_N \left(\frac{(g(\frac{z-t}{a}) + f(\frac{z+t}{a}))^2}{2g'(\frac{z-t}{a})f'(\frac{z+t}{a})} \right)^{(n+2)/n}. \quad (19.104)$$

In the second region (large field)

$$q_0(t, z) = \frac{2\sqrt{2\pi\overline{M}}}{m_{\text{pl}}(n+2)} \cdot \ln \left[\frac{2g'(\frac{z-t}{b})f'(\frac{z+t}{b})}{(g(\frac{z-t}{b}) + f(\frac{z+t}{b}))^2} \right], \quad (19.105)$$

$$b = \frac{2\sqrt{2}}{\sqrt{(n+2)\overline{\lambda}_{c0}}}, \quad (19.106)$$

and

$$G_{\text{eff}} = G_N \cdot \frac{(g(\frac{z-t}{b}) + f(\frac{z+t}{b}))^2}{2g'(\frac{z-t}{b})f'(\frac{z+t}{b})}. \quad (19.107)$$

In this way we get a spatio-temporal pattern of changing the effective gravitational constant for small and large field regions. In both cases we have $\varepsilon\tilde{\varepsilon} = -1$. However, in the small field region we have $\varepsilon = 1$ and because of this $\tilde{\varepsilon} = -1$. In the case of large field region $\varepsilon = -1$ and $\tilde{\varepsilon} = 1$. In order to be in line with our assumptions we should consider in the first case such functions f and g that the expression in (19.105) is small and for the second case vice versa.

Let us consider Eq. (19.63) in cylindrical coordinates supposing cylindrical symmetry for the field q_0 , $q_0 = q_0(\rho)$. One gets

$$\frac{1}{\rho} \left(\rho \frac{dq_0}{d\rho} \right) - \varepsilon \bar{\alpha} \exp(n \bar{\beta} q_0) (\exp(2 \bar{\beta} q_0) - 1) = 0. \quad (19.108)$$

This equation can be transformed into

$$\frac{1}{\tau} \frac{d}{d\tau} \left(\tau \frac{d\varphi}{d\tau} \right) - \tilde{\varepsilon} \exp(n\varphi) (e^{2\varphi} - 1) = 0, \quad (19.109)$$

$$\rho = \frac{2}{\sqrt{\lambda_{c0}}} \tau. \quad (19.110)$$

As usual we consider Eq. (19.109) in two regions for small and large fields.

In the first region

$$\frac{1}{\tau} \frac{d}{d\tau} \left(\tau \frac{d\varphi}{d\tau} \right) + \tilde{\varepsilon} e^{n\varphi} = 0. \quad (19.111)$$

In the second region we get

$$\frac{1}{\tau} \frac{d}{d\tau} \left(\tau \frac{d\varphi}{d\tau} \right) - \tilde{\varepsilon} e^{(n+2)\varphi} = 0. \quad (19.112)$$

Both equations can be reduced to the equation

$$\frac{1}{x} \frac{d}{dx} \left(x \frac{dy}{dx} \right) + \varepsilon \tilde{\varepsilon} e^y = 0, \quad (19.113)$$

where in the first region $\varepsilon = 1$ and

$$y = n\varphi \quad (19.114)$$

$$x = \sqrt{n} \tau \quad (19.115)$$

and in the second region $\varepsilon = -1$ and

$$y = (n+2)\varphi \quad (19.116)$$

$$x = \sqrt{n+2} \tau. \quad (19.117)$$

We can transform (19.113) into

$$x \frac{d^2 y}{dx^2} + \frac{dy}{dx} + \varepsilon \tilde{\varepsilon} x e^y = 0 \quad (19.118)$$

which is the equation considered in [132] for $\varepsilon \tilde{\varepsilon} = 1$.

Following H. Lemke we write down a solution to Eq. (19.118) in a compact form in three cases (concerning an integration constant introduced by H. Lemke). We adopt his solutions to our problem.

1) $C = \kappa^2 > 0$, κ —arbitrary positive number:

$$y(x) = -2 \ln \left(\frac{x}{\bar{a}} \left(\left(\frac{\bar{a}}{x} \right)^\kappa + \left(\frac{x}{\bar{a}} \right)^\kappa \right) \right) + \ln(4\kappa^2 \bar{a}^2) \quad (19.119)$$

where $\bar{a} > 0$ is an arbitrary constant.

2) $C = -\omega^2 < 0$, ω is an arbitrary positive number:

$$y(x) = -2 \ln(2x \sin(\omega \ln x + \delta)) + \ln(4\omega^2) \quad (19.120)$$

where δ is an arbitrary constant.

3) $C = 0$:

$$y(x) = -2 \ln \left(\frac{x}{\bar{a}} \ln \left(\frac{x}{\bar{a}} \right) \right) - 2 \ln \bar{a} \quad (19.121)$$

where $\bar{a} > 0$ is an arbitrary constant.

Using these solutions we write down a spatial dependence of G_{eff} in the case of small and large fields. In the case of small fields one gets

$$1) \quad G_{\text{eff}} = G_N (4\kappa^2 \bar{a}^2)^{(n+2)/n} \frac{(1 + r^{2\kappa})^{2(n+2)/n}}{r^{2(\kappa-1)(n+2)/n}} \quad (19.122)$$

where

$$r = \frac{1}{2\bar{a}} \sqrt{n\bar{\lambda}_{c0}} \rho. \quad (19.123)$$

$$2) \quad G_{\text{eff}} = \frac{G_N (\bar{\delta})^{2(n+2)/n}}{(\omega^2)^{(n+2)/n}} r^{2(n+2)/n} (\sin(\omega \ln r))^{2(n+2)/n} \quad (19.124)$$

where

$$r = \frac{1}{2\bar{\delta}} \sqrt{n\bar{\lambda}_{c0}} \rho \quad (19.125)$$

$$\omega \ln \bar{\delta} = -\delta \quad (19.126)$$

$$3) \quad G_{\text{eff}} = G_N \bar{a}^{2(n+2)/n} r^{2(n+2)/n} (\ln r)^{2(n+2)/n} \quad (19.127)$$

and r is given by Eq. (19.123).

In the case of large field one gets:

$$1) \quad G_{\text{eff}} = G_N (4\kappa^2 \bar{a}^2) \frac{(1 + r^{2\kappa})^2}{r^{2(\kappa-1)}} \quad (19.128)$$

where

$$r = \frac{1}{2\bar{a}} \sqrt{(n+2)\bar{\lambda}_{c0}} \rho. \quad (19.129)$$

$$2) \quad G_{\text{eff}} = G_N \left(\frac{\bar{\delta}}{\omega} \right)^2 r^2 (\sin(\omega \ln r))^2 \quad (19.130)$$

where

$$r = \frac{1}{2\bar{\delta}} \sqrt{(n+2)\bar{\lambda}_{c0}} \rho, \quad \ln \bar{\delta} = -\frac{\delta}{\omega}. \quad (19.131)$$

$$3) \quad G_{\text{eff}} = G_N \bar{a}^2 r^2 (\ln r)^2 \quad (19.132)$$

and r is given by Eq. (19.129).

Let us notice that in that spatial dependence for large and small field we have κ and $\bar{a}$ ($\bar{\delta}$, ω) as integration constants. In this way integration constants induce a power law of this dependence and also a scale. For sufficiently big n ($n > 14$) there is no significant difference between both cases. It means the quintessence field behaves everywhere as for large field case (in these solutions of course). It is evident that the spatial dependence (in cylindrical symmetry case) of G_{eff} goes to some kind of the fifth force. However, we have to do not with a universal law of Nature but rather with some kind of initial conditions. Thus the real dependence of G_{eff} on spatial coordinates can be obtained after averaging G_{eff} with respect to initial conditions. Let us describe it using Eq. (19.128).

Let us write G_{eff} in a form where κ and $\bar{a}$ are explicitly visible:

$$G_{\text{eff}} = G_N (4\kappa^2 \bar{a}^2) \cdot \frac{\left(1 + \left(\frac{x}{\bar{a}}\right)^{2\kappa}\right)^2}{\left(\frac{x}{\bar{a}}\right)^{2(\kappa-1)}}, \quad (19.133)$$

$$x = \frac{1}{2} \sqrt{(n+2)\bar{\lambda}_{c0}} \rho. \quad (19.134)$$

The constants $\bar{a}$ and κ are chosen randomly for they are dependent on initial conditions. Let μ be a measure defined on $(0, +\infty)^2$, positive and normalized to 1, i.e.

$$\int_0^{+\infty} \int_0^{+\infty} d\mu(\bar{a}, \kappa) = 1. \quad (19.135)$$

This measure gives an account how frequently we have to do with some initial conditions (i.e. with integration constants $\bar{a}$, κ). In this way an experimental dependence of G_{eff} on x should be obtained from

$$E(G_{\text{eff}}) = 4G_N \int_0^{+\infty} \int_0^{+\infty} \frac{\kappa^2 \bar{a}^2 \left(1 + \left(\frac{x}{\bar{a}}\right)^{2\kappa}\right)^2}{\left(\frac{x}{\bar{a}}\right)^{2(\kappa-1)}} d\mu. \quad (19.136)$$

It means it is an expectation value of G_{eff} with respect to the measure μ . μ need not be absolutely continuous with respect to the Lebesgue measure on $\mathbb{R}^2$. The formula (19.136) could give an account on some serious problems with comparisons of several measurements of G (a Newton constant). Maybe we have to do with different initial conditions for quintessence field. These initial conditions appear with different probabilities according to the measure $d\mu(\bar{a}, \kappa)$ going to an expectation value $E(G_{\text{eff}})$. The

second central moment of $d\mu(\bar{a}, \kappa)$ (if it exists) can express a deviation from the law described by $E(G_{\text{eff}})$.

In the simplest case we can suppose a continuous Gaussian distribution for κ only:

$$d\mu(\kappa) = \frac{1}{\sqrt{2\pi}\sigma} e^{-(\kappa-\kappa_0)^2/(2\sigma^2)} d\kappa.$$

In this case we have

$$\overline{G}_{\text{eff}}(x) = E(G_{\text{eff}}(x)) = \frac{G_N \bar{a}^2}{\sqrt{2\pi}\sigma} \int_{-\infty}^{+\infty} d\kappa e^{-(\kappa-\kappa_0)^2/(2\sigma^2)} \kappa^2 \frac{\left(1 + \left(\frac{x}{\bar{a}}\right)^{2\kappa}\right)^2}{\left(\frac{x}{\bar{a}}\right)^{2(\kappa-1)}}. \quad (19.136a)$$

One gets after some algebra

$$\begin{aligned} \overline{G}_{\text{eff}}(x) = & G_N \kappa_0 \exp\left(\bar{a} - \frac{\kappa_0^2}{2\sigma^2}\right) \left(\frac{x}{\bar{a}}\right)^2 \\ & \times \left[4\sigma^4 \exp\frac{(2\ln\left(\frac{x}{\bar{a}}\right)\sigma^2 + \kappa_0^2)^2}{2\sigma^2} \ln\left(\frac{x}{\bar{a}}\right) \right. \\ & + (2\sigma^2 + \kappa_0^4) \exp\frac{(2\ln\left(\frac{x}{\bar{a}}\right)\sigma^2 + \kappa_0^2)^2}{2\sigma^2} \\ & \left. - 4\sigma^4 \exp\frac{(\sigma + \kappa_0)^2}{2} \ln^2\left(\frac{x}{\bar{a}}\right) + 2(\sigma^2 + \kappa_0^2) \exp\frac{(\sigma + \kappa_0)^2}{2} \right]. \end{aligned} \quad (19.136b)$$

In the case of the solution (19.107) parametrized by two functions of one variable we can consider them as random variables parametrized by z and t . Moreover in this case we should consider a measure μ on an infinite-dimensional space of functions f and g , supposing that solution (19.107) is generalized to this space. Afterwards we can use as μ the Wiener or Gaussian measure in L^2 space. If we use for Gaussian distribution the normalized distribution $N(0, 1)$, the formula (19.136b) simplifies to

$$\begin{aligned} \overline{G}_{\text{eff}}(x) = & e^{\bar{a}} \sqrt{e} G_N \\ & \times \left[4 \left(\frac{x}{\bar{a}}\right)^{2(1+\ln(x/\bar{a}))} \ln\left(\frac{x}{\bar{a}}\right) + \left(\frac{x}{\bar{a}}\right)^{2(1+\ln(x/\bar{a}))} - 4 \ln^2\left(\frac{x}{\bar{a}}\right) + 2 \right]. \end{aligned} \quad (19.136c)$$

Let us consider three our cases (19.119), (19.120) and (19.121) for small and large fields cases.

The small field case is such that

$$e^\varphi < 1, \quad \text{i.e. } \varphi < 0. \quad (19.137)$$

The large field case is if

$$e^\varphi > 1, \quad \text{i.e. } \varphi > 0. \quad (19.138)$$

For (19.119) we have for the small case

$$f(r) > 1,$$

where

$$f(r) = r^{1-\kappa} + r^{\kappa+1}, \quad r = \left(\frac{x}{a}\right). \quad (19.139)$$

Let us consider two cases

$$0 < \kappa < 1 \quad \text{and} \quad \kappa > 1.$$

In the first case $f(r) \nearrow$ in $[0, +\infty)$ and we have simply $r > r_0$ where r_0 satisfies the equation

$$r_0^{2\kappa+1} + r_0 - 1 = 0. \quad (19.140)$$

In the second case $\kappa > 1$,

$$\lim_{r \rightarrow 0} f(r) = +\infty \quad \text{and} \quad \lim_{r \rightarrow \infty} f(r) = +\infty.$$

The function $f(r)$ has a minimum at

$$r_1 = \left(\frac{\kappa-1}{\kappa+1}\right)^{1/\kappa}. \quad (19.141)$$

Let us calculate $f(r_1)$.

$$f(r_1) = \left(\frac{\kappa+1}{\kappa-1}\right)^{\kappa-1/\kappa} + \left(\frac{\kappa-1}{\kappa+1}\right)^{\kappa+1/\kappa}. \quad (19.142)$$

It is easy to see that if $\kappa > 1$, then $\kappa - 1/\kappa > 0$. This means that

$$f(r_1) > 1 \quad (19.143)$$

and therefore $f(r) > 1$ for all $r > 0$. Thus simultaneously we get a solution for large field only if $\kappa < 1$, i.e.

$$r < r_0. \quad (19.144)$$

If $\kappa = 1$, we always have $f(r) \geq 1$ (i.e. only a small field).

Let us consider (19.120). In this case the small field condition reads

$$h(r) = r \sin(\omega \ln r) > 1, \quad \text{where } r = \frac{x}{\delta}. \quad (19.145)$$

First of all we need $h(r) > 0$. Let us observe that $|h(r)| \leq r$ for every $r > 0$. Next we see that the roots of $h(r)$ are the numbers

$$r_{0,k} = e^{k\pi/\omega}, \quad k = 0, \pm 1, \pm 2, \dots \quad (19.146a)$$

and

$$h(r) > 0 \quad \text{if} \quad r_{0,2k} < r < r_{0,2k+1}, \quad k = 0, \pm 1, \pm 2, \dots \quad (19.146b)$$

The maximum of the function $h(r)$ in the interval $(r_{0,2k}, r_{0,2k+1})$ is greater than

$$h(r_{0,2k+1/2}) = e^{2k\pi/\omega} \cdot e^{\pi/(2\omega)}, \quad (19.147)$$

but smaller than $r_{0,2k+1}$. Thus the maxima are smaller than 1 for negative k and greater than 1 if $k = 0, 1, 2, \dots$. It means that in each interval $(r_{0,2k}, r_{0,2k+1})$, $k = 0, 1, 2, \dots$, there exist two numbers $r_{3,2k}$ and $r_{2,2k}$ such that $r_{3,2k} < r_{2,2k}$,

$$h(r_{3,2k}) = h(r_{2,2k}) = 1 \quad (19.148)$$

and

$$h(r) > 1 \quad \text{if} \quad r_{3,2k} < r < r_{2,2k}. \quad (19.149)$$

The condition for large fields, $0 < h(r) < 1$, is satisfied if

$$r_{0,2k} < r < r_{0,2k+1}, \quad k = -1, -2, \dots \quad (19.150)$$

or

$$r_{0,2k} < r < r_{3,2k}, \quad \text{or} \quad r_{2,2k} < r < r_{0,2k+1}, \quad k = 0, 1, 2, \dots \quad (19.151)$$

In the third case, i.e. Eq. (19.121), one has for the small field case

$$r \ln r > 1 \quad (19.152)$$

where $r = \frac{x}{a}$. Let $r_4 \ln r_4 = 1$, $r_4 > 1$. We have $r > r_4 = 1.7632 \dots$. In the large field case

$$0 < r < r_4 = 1.7632 \dots \quad (19.153)$$

Thus we have in general large fields on large distances.

In this way we have solutions for large and small distances. One can try to connect them to get a solution for all distances. However in this case it is necessary to be very careful, for our solutions depend on some integration constants which can be different for both asymptotic regions.

Let us consider Eq. (19.63) in two special cases:

- I $q_0 = q(z)$ — static and depending only on z ;
- II $q_0 = q_0(t)$ — non-static and spatially constant.

Let us consider also these cases for small and large fields φ . In all of these cases we come to the following equation

$$\frac{d^2 y}{dx^2} + \varepsilon_1 \varepsilon \tilde{e} e^y = 0 \quad (19.154)$$

where $\varepsilon_1 = 1$ for case I and $\varepsilon_1 = -1$ for case II.

Eq. (19.154) can easily be reduced to the integral

$$x - x_0 = \frac{\varepsilon_3}{\sqrt{2}} \int \frac{dy}{\sqrt{\varepsilon_2 \omega^2 - \eta e^y}} \quad (19.155)$$

where $\eta = \varepsilon_1 \varepsilon \tilde{e}$, $\eta^2 = 1$, $C = 2\varepsilon_2 \omega^2$, $\omega \geq 0$, $\varepsilon_2^2 = 1$ is an integration constant, $\varepsilon_3^2 = 1$ and x_0 also is an integration constant.

After some calculation we get the following solutions:

$$\text{A.} \quad y(x) = 2 \left[\ln \omega - \ln \left| \sinh \left(\frac{\sqrt{2}(x-x_0)\omega}{2} \right) \right| \right], \quad \eta = -1, \varepsilon_2 = 1 \quad (19.156)$$

$$\text{B.} \quad y(x) = 2 \left[\ln \omega - \ln \left| \sinh \left(\frac{\sqrt{2}(x-x_0)\omega}{2} \right) \right| \right], \quad \eta = 1, \varepsilon_2 = 1 \quad (19.157)$$

where

$$\frac{\sqrt{2}(x-x_0)\omega}{2} > \ln(1 + \sqrt{2}) \quad (19.158)$$

or

$$\frac{\sqrt{2}(x-x_0)\omega}{2} < \ln(\sqrt{2} - 1). \quad (19.159)$$

$$\text{C.} \quad y(x) = 2 \left[\ln \omega - \ln \left| \cos \left(\frac{\sqrt{2}(x-x_0)\omega}{2} \right) \right| \right], \quad \eta = -1, \varepsilon_2 = -1. \quad (19.160)$$

Let us apply these solutions to our problems. First of all let us consider a static configuration with z dependence only. In this case $\varepsilon_1 = 1$ and $\eta = \varepsilon\tilde{\varepsilon}$. For the small field case one gets, $\varepsilon = 1, \eta = \tilde{\varepsilon}$,

$$G_{\text{eff}} = \frac{G_N}{\omega^{2(n+2)/n}} |\sinh p|^{(n+2)/n} \quad (19.161)$$

where

$$p = \frac{\omega(z-z_0)}{4} \sqrt{n\bar{\lambda}_{c0}}. \quad (19.162)$$

For the large field case we get, $\varepsilon = -1, \eta = -\tilde{\varepsilon}$,

$$G_{\text{eff}} = \frac{G_N}{\omega^2} |\cos p| \quad (19.163)$$

where

$$p = \frac{\omega(z-z_0)}{4} \sqrt{(n+2)\bar{\lambda}_{c0}}. \quad (19.164)$$

In this case $\tilde{\varepsilon} = 1$ for $\eta = -1$.

In a nonstatic configuration $\varepsilon_1 = -1$ and $\eta = -\varepsilon\tilde{\varepsilon}$. For the small field case ($\varepsilon = 1$), $\eta = -\tilde{\varepsilon}$,

$$G_{\text{eff}} = \frac{G_N}{\omega^{2(n+2)/n}} |\sinh q|^{(n+2)/n} \quad (19.165)$$

where

$$q = \frac{\varepsilon_3 \omega(t-t_0)}{4} \sqrt{n\bar{\lambda}_{c0}} \quad (19.166)$$

and analogically for the large field case ($\varepsilon = -1$), $\eta = \tilde{\varepsilon}$,

$$G_{\text{eff}} = \frac{G_N}{\omega^2} |\cos q|, \quad (19.167)$$

$$q = \frac{\omega(t - t_0)}{4} \sqrt{(n + 2)\bar{\lambda}_{c0}}. \quad (19.168)$$

In this case $\tilde{\varepsilon} = -1$ for $\eta = -1$.

Let us notice that in a static configuration for small field we have two possibilities for $\eta = -1$ (no condition on p) and $\eta = 1$ (conditions (19.158–159)). Thus without conditions we have $\tilde{\varepsilon} = -1$ and with conditions $\tilde{\varepsilon} = 1$. In a non-static configuration for small field we have vice versa $\tilde{\varepsilon} = 1$ without conditions and $\tilde{\varepsilon} = -1$ with conditions (19.158–159).

Let us come back to the Eq. (19.89) and consider it in a travelling wave scheme. In this way we have

$$q_0(z, t) = \tilde{q}(z - vt) \quad (19.169)$$

where v is a velocity of the travelling wave (a soliton), $|v| < 1$. Let us consider this equation in both regimes (for small and large fields). In this way we come to the expression

$$(1 - v^2) \frac{d^2 \chi}{d\xi^2} - \varepsilon \tilde{\varepsilon} e^\chi = 0 \quad (19.170)$$

where χ is a shape function of a soliton. Changing an independent variable from ξ to λ one gets

$$\frac{d^2 \chi}{d\lambda^2} - \varepsilon_1 \varepsilon \tilde{\varepsilon} e^\chi = 0 \quad (19.171)$$

where $\varepsilon_1 = -1$, i.e. we get Eq. (19.154) with $\eta = -\varepsilon \tilde{\varepsilon}$,

$$\lambda = \frac{\xi}{\sqrt{1 - v^2}}, \quad \xi = \sqrt{1 - v^2} \lambda. \quad (19.172)$$

In this way we adopt our solutions A, B, C in both regimes: small and large field (changing χ into λ). For small field we get ($\varepsilon = 1$, $\eta = -\tilde{\varepsilon}$)

$$q_0(z, t) = \frac{2}{\beta n} [\ln \omega - \ln |\sinh p|] \quad (19.173)$$

where

$$p = \frac{\omega \sqrt{n \bar{\lambda}_{c0}}}{4 \sqrt{1 - v^2}} (z - vt). \quad (19.174)$$

For large field we get ($\varepsilon = -1$, $\eta = \tilde{\varepsilon}$)

$$q_0(z, t) = \frac{2}{\beta(n + 2)} [\ln \omega - \ln |\cos p|] \quad (19.175)$$

where

$$p = \frac{\omega \sqrt{(n+2)\bar{\lambda}_{c0}}}{4\sqrt{1-v^2}} (z - vt). \quad (19.176)$$

For (19.173) we have $\eta = -\tilde{\varepsilon}$ and because of this $\tilde{\varepsilon} = 1$ without any conditions and if $\tilde{\varepsilon} = -1$ we have conditions (19.158–159). In the case of the formula (19.174) $\eta = -1$ and $\tilde{\varepsilon} = -1$.

We can write down formulas for G_{eff} in the soliton case

$$G_{\text{eff}} = \frac{G_N}{\omega^{2(n+2)/n}} |\sinh p|^{(n+2)/n} \quad (19.177)$$

and p is given by the formula (19.174) (with or without conditions (19.158–159)). This is of course a small field case.

In the large field case

$$G_{\text{eff}} = \frac{G_N}{\omega^2} |\cos p|, \quad (19.178)$$

and p is given by the formula (19.176). In this case $\tilde{\varepsilon} = -1$. (Let us notice that this is a case of SO(3) group in our theory.)

Let us notice that conditions (19.158–159) can be considered as conditions for small field in z or t domains. Let us notice that in our solutions concerning a behaviour of an effective gravitational constant we get completely arbitrary length or time scale (given by integration constants). In this way a spatial or time dependence of G_{eff} can be (except the solution (19.80) and simultaneously the approximate solution in the case of spherical symmetry) such that G_{eff} can be really constant on distances (or times) accessible in experiments. Only a statistical approach mentioned here can give a light on this dependence to compare it with an experiment.

In order to connect our results to the ordinary gravitational physics we consider again Eq. (19.67) in small field regime for initial conditions $\varphi(0) = 0$ and $\frac{d\varphi}{dx}(0) = 0$. The first condition means that we want to have $G_{\text{eff}}(0) = G_N$ and the second that the quintessence field does not grow quickly. The problem cannot be solved analytically. Moreover R. Emden in the first point of Ref. [131] did it for us. We quote here his results adopted to our notation ($\varepsilon = +1$ and $\tilde{\varepsilon} = -1$).

x	$-y$	e^y	G_{eff}/G_N
0.00	0.00000	1.00000	1.00000
0.25	0.01037	0.98969	0.98823
0.50	0.04113	0.95971	0.95409
0.75	0.09113	0.91290	0.90109
1.00	0.15903	0.85296	0.83380
1.25	0.24225	0.78486	0.75816
1.50	0.33847	0.71285	0.67920
1.75	0.44488	0.64090	0.60143
2.00	0.55967	0.57140	0.52749
2.50	0.80584	0.44671	0.39813
3.00	1.06226	0.34537	0.29670
3.50	1.31937	0.26730	0.22138
4.00	1.57071	0.20790	0.16611
4.50	1.81246	0.16325	0.12601
5.00	2.04264	0.12968	0.09686
6.00	2.46598	0.08493	0.05971
7.00	2.84160	0.05833	0.03887
8.00	3.17489	0.04180	0.02656
9.00	3.47128	0.03108	0.01893
10.00	3.73646	0.02384	0.01398
100	8.59506	0.000175	$5.0854 \cdot 10^{-5}$
1000	13.09847	0.000002	$3.0683 \cdot 10^{-7}$

where

$$x = \frac{1}{2} \sqrt{n \bar{\lambda}_{c0}} r \cong \left(\frac{r}{10 \text{ Mpc}} \right), \quad (19.179)$$

$$\frac{G_{\text{eff}}}{G_N} = (e^y)^{(n+2)/n} = (e^y)^{8/7}. \quad (19.180)$$

We take $n = 14$.

It is easy to see that for large n

$$\frac{G_{\text{eff}}}{G_N} = e^y. \quad (19.181)$$

It is easy to see that on a distance of 1 Mpc G_{eff} does not differ from G_N . Even on a distance of 10 Mpc it is about 10% smaller. Thus in the Solar System Newtonian gravitational physics does not change. Even on the level of a galaxy this change is minimal and cannot be observed. Moreover, there is an important conclusion: on distances about 200 Mpc the strength of gravitational interactions is about 10^{-5} times this on short distances measured in the Solar System (and for 10^3 Mpc of 10^{-7}). It is

hard to tell how it influences a mass of a cluster of galaxies if we realize that from any observational data only a product GM has been obtained (not M).

From the other side on distances of 100 Mpc the strength of gravitational interactions is very weak (not only because of the distance). Thus if we consider clusters of galaxies as substrat particles in cosmology then they do not interact.

Let us consider Eq. (19.63) in Cartesian coordinates for two-dimensional static case (i.e. $\frac{\partial}{\partial z} = 0$, $\frac{\partial}{\partial t} = 0$). One gets

$$\left(\frac{\partial^2 q_0}{\partial x^2} + \frac{\partial^2 q_0}{\partial y^2} \right) - \tilde{\varepsilon} \bar{\alpha} \exp(n \bar{\beta} q_0) (\exp(2 \bar{\beta} q_0) - 1) = 0 \quad (19.182)$$

(where $\bar{\alpha}$, $\bar{\beta}$ are given by formulas (19.64) and (19.65)).

As usual, we come to the formula

$$\left(\frac{\partial^2 \varphi}{\partial x_1^2} + \frac{\partial^2 \varphi}{\partial x_2^2} \right) - \tilde{\varepsilon} e^{n\varphi} (e^{2\varphi} - 1) = 0 \quad (19.183)$$

where

$$\left. \begin{matrix} x \\ y \end{matrix} \right\} = \frac{2}{\sqrt{\lambda_{c0}}} x_i, \quad i = 1, 2. \quad (19.183a)$$

We consider Eq. (19.183) for small and large fields and we get

$$\left(\frac{\partial^2 \chi}{\partial z_1^2} + \frac{\partial^2 \chi}{\partial z_2^2} \right) - \varepsilon \tilde{\varepsilon} e^\chi = 0 \quad (19.184)$$

where as usual for a small field $\varepsilon = 1$ and

$$\chi = n\varphi, \quad (19.185)$$

$$z_i = \sqrt{n} x_i \quad (19.186)$$

and for a large field $\varepsilon = -1$ and

$$\chi = (n+2)\varphi, \quad (19.187)$$

$$z_i = \sqrt{n+2} x_i. \quad (19.188)$$

Thus we come to the equation known as Liouville equation

$$\Delta \chi = e^\chi \quad (19.184a)$$

if $\varepsilon \tilde{\varepsilon} = 1$.

This equation can be explicitly solved. First of all we change independent variables into

$$Z = \frac{1}{\sqrt{2}}(z_1 + iz_2) \quad (19.189)$$

$$\chi(Z) = -\ln\left(\frac{1}{2}(1 - |g|^2)\right) + \frac{1}{2} \ln \left| \frac{dg}{dZ} \right| \quad (19.190)$$

where g is an arbitrary analytic function on a complex plane Z .

In this way we get for the small field case

$$G_{\text{eff}} = G_N \left(\frac{1 - |g(Z)|^2}{2} \right)^{(n+2)/n} \left(\left| \frac{dg}{dZ}(Z) \right| \right)^{-(n+2)/(2n)} \quad (19.191)$$

where

$$Z = \sqrt{\frac{\bar{\lambda}_{c0}}{2n}} \cdot (x + iy). \quad (19.192)$$

In the large field case

$$G_{\text{eff}} = G_N \left(\frac{1 - |g(Z)|^2}{2} \right) \left(\left| \frac{dg}{dZ}(Z) \right| \right)^{-1/2} \quad (19.193)$$

where

$$Z = \sqrt{\frac{\bar{\lambda}_{c0}}{2(n+2)}} \cdot (x + iy). \quad (19.194)$$

Eqs (19.191) and (19.193) can have very interesting behaviour for $\frac{dg}{dZ}$ could have some singularities. The physical interpretation of these singularities can be very interesting.

Similarly as for Eqs (19.104) and (19.107) we can consider an expectation value of G_{eff} with respect to some kind of normalized measure for a space of analytic functions on the complex plane (respectively chosen).

It seems that an assumption all the energy of a quintessence is stored as quintessence particles is unrealistic. Let us suppose that only a fraction of this energy is stored as particles. Let this fraction be η ,

$$0 < \eta < 1. \quad (19.195)$$

η can be a function of time and even of a space-point (locally).

Let us suppose that a gas of quintessence particles is a perfect gas governed by the Clapeyron equation. Thus we have

$$\frac{p_1}{\rho_1} = \frac{K_B T}{m_0} = \frac{T}{T_0}, \quad T_0 = \frac{m_0}{K_B} \simeq 0.11 \text{ } ^\circ\text{K} \quad (19.196)$$

(if we take $m_0 \simeq 10^{-5} \text{ eV}$). T is the temperature of a gas.

One gets

$$\rho_1 = \eta \rho_Q = \eta \rho \quad (19.197)$$

$$p = p_Q + p_1 = p_Q + \eta \frac{T}{T_0} \rho = -\rho(1 - \eta) + \rho \frac{T}{T_0} \eta. \quad (19.198)$$

Eventually one gets

$$p = \rho \left(\eta \left(1 + \frac{T}{T_0} \right) - 1 \right) \quad (19.199)$$

$$p = w\rho \quad (19.200)$$

$$w = \eta \left(1 + \frac{T}{T_0} \right) - 1. \quad (19.201)$$

Now we can calculate an isothermic speed of sound in a quintessence.

$$C_1^2 = \frac{p}{\rho} = w = \eta \left(1 + \frac{T}{T_0} \right) - 1. \quad (19.202)$$

For

$$0 < C_1^2 < 1 \quad (19.203)$$

we get

$$\frac{T_0}{T + T_0} < \eta < \min \left[\frac{2T_0}{T_0 + T}, 1 \right]. \quad (19.204)$$

Let us remind to the reader that an isothermic sound is appropriate for low frequency of acoustic waves (we have to do with this sound in astrophysics). In general we have to do with so called adiabatic sound. In order to calculate a speed of an adiabatic sound in a quintessence we should find an analogue for a Poisson adiabat for our equation of state. One gets supposing that an internal energy of a quintessence is an energy of one-atomic gas of quintessence particles

$$dU + p dV = 0 \quad (19.205)$$

or

$$\frac{3}{2} dT = \frac{d\rho}{\rho} \left(\eta - 1 + \frac{T}{T_0} \eta \right) \quad (19.206)$$

and finally

$$\frac{p}{\rho^\kappa} = \text{const.} \quad (19.207)$$

$$\kappa = \frac{2\eta + 3}{3} \quad (19.208)$$

and a speed of an adiabatic sound simply reads

$$C_2 = \sqrt{\frac{\kappa p}{\rho}} = \sqrt{\kappa \left(\eta \left(1 + \frac{T}{T_0} \right) - 1 \right)}. \quad (19.209)$$

It is interesting to ask what kind of a polytrope κ represents. Let us remind to the reader that in general

$$\kappa = \frac{c_p - c}{c_v - c} \quad (19.210)$$

where c is a specific heat of a polythroe in mind.

One gets in our case

$$c = \frac{3(\eta - 1)}{2\eta} \bar{R} \quad (19.211)$$

where $\bar{R}$ is a universal gas constant.

Thus we found both speeds of sound in a quintessence for low frequency and for high frequency of sound. The measurement of both speeds can help us to find η and T . One gets

$$\eta = \frac{3}{2} \left(\left(\frac{C_2}{C_1} \right)^2 - 1 \right) \quad (19.212)$$

and

$$T = \frac{T_0}{3(C_2^2 - C_1^2)} (C_1^2(2C_1^2 + 5) - 3C_2^2). \quad (19.213)$$

Let us notice that a gas of quintessence particles is quite cold. Moreover these particles are highly relativistic. For the temperature $T \simeq T_0 = 0.11$ °K one sees that a speed of a quintessence particle is about 0.91 of the speed of light. Moreover we can consider lower temperatures. It is interesting to notice that the speed of sound is of the same order.

Thus if we want to have both speeds smaller than a speed of light,

$$0 < C_i^2 < 1, \quad i = 1, 2, \quad (19.214)$$

we get

$$\eta > \frac{1 + \sqrt{13 + 12t}}{1 + t} = f(t) \quad (19.215)$$

where

$$t = \frac{T}{T_0}. \quad (19.216)$$

The condition

$$t = \frac{T}{T_0} > 13 \quad (19.217)$$

guarantees that $f(t) < 1$. Moreover for $t = 14$ one gets

$$\eta \geq 0.96357. \quad (19.218)$$

This seems quite interesting, however maybe too much. Let us calculate a mean scattering length of quintessence particles for such a big $\eta = 0.96357$

$$l_{\text{scattering}} = \frac{1}{\eta \sigma n} = 1.0378 \cdot l. \quad (19.219)$$

We have still to do with a very dense gas of quintessence particles.

It seems that it would be reasonable to repeat these calculations using a different equation of state for quintessence particles gas. In particular we can consider a gas of quintessence particles as a massless boson gas (with spin zero). In this case the equation of state looks:

$$p = p_Q + p_1 = -(1 - \eta)\rho + \frac{4\eta\sigma\rho}{3} T^4 \quad (19.220)$$

and an adiabat equation

$$dU = \bar{c}_v dT + p dV = 0 \quad (19.221)$$

where

$$\bar{c}_v = 16\sigma T^3, \quad (19.222)$$

σ is the Stefan-Boltzmann constant.

One easily integrates

$$\frac{p}{\rho^{\bar{\kappa}}} = \text{const.} \quad (19.223)$$

where

$$\bar{\kappa} = \frac{3 + \eta}{3}. \quad (19.224)$$

We have as before two kinds of sound: an isothermic sound

$$\bar{C}_1 = \sqrt{\frac{p}{\rho}} = \sqrt{\eta - 1 + \frac{4\eta\sigma}{3} T^4} \quad (19.225)$$

and an adiabatic sound

$$\bar{C}_2 = \sqrt{\bar{\kappa} \left(\eta - 1 + \frac{4\eta\sigma}{3} T^4 \right)} = \sqrt{\frac{\bar{\kappa} p}{\rho}}. \quad (19.226)$$

From $0 < \bar{C}_i^2 < 1$, $i = 1, 2$, one gets

$$\frac{1}{1 + (T/\bar{T}_0)^4} < \eta < \min \left[1, \frac{2}{1 + (T/\bar{T}_0)^4} \right] \quad (19.227)$$

and

$$\eta > \frac{-3t^4 - 1 + \sqrt{9t^8 + 18t^4 + 13}}{2(1 + t^4)} = \bar{f}(t) \quad (19.228)$$

with the condition

$$t = \frac{T}{\bar{T}_0} > \frac{1}{\sqrt{2}} \simeq 0.707. \quad (19.229)$$

This condition guarantees that

$$\bar{f}(t) < 1 \quad (19.230)$$

where $T_0^4 = \frac{3}{4\sigma}$ or

$$\bar{T}_0 = \frac{1}{K_B} \sqrt[4]{\frac{360}{\pi}} \simeq \frac{3.27}{K_B}, \quad (19.231)$$

K_B is the Boltzmann constant.

From the measurements of $\bar{C}_1$ and $\bar{C}_2$ we can obtain as before η and T . One gets:

$$\eta = 3 \left(\left(\frac{\bar{C}_2}{\bar{C}_1} \right)^2 - 1 \right) \quad (19.232)$$

and

$$T = \frac{\bar{T}_0}{\sqrt[4]{3}} \left(\frac{4\bar{C}_1^2 + \bar{C}_1^4 - 3\bar{C}_2^2}{\bar{C}_2^2 - \bar{C}_1^2} \right)^{1/4}. \quad (19.233)$$

Recently many papers have appeared concerning the speed of sound in a quintessence (see Ref. [133]). In some of them the authors propose to measure this speed.

There are interesting propositions to include primordial gravitational waves to fluctuations of a quintessence and vice versa.

We consider a mass of a quintessence particle, a speed of sound in a quintessence and several solutions to quintessence equations coming to the interesting behaviour of an effective gravitational constant (see [135]). Some of issues of our theory are considered in self-interacting Brans-Dike theory [134].

The fraction η can be connected with the part of an energy density of quintessence field Q in such a way that it is a fraction of fluctuation energy density around an equilibrium Q_0 . In this way the q_0 field which can evolve due to an evolution of Q and due to fluctuations of primordial gravitational waves will be a source of a gas of thermalized particles. It seems that an approach with a boson equation of state is more appropriate to consider.

It is interesting to consider a fraction of a dark energy stored as boson particles as a dark matter in the Universe.

Now we can also calculate a mass of a skewon particle in a linear approximation from Sec. 17. Let us consider Eq. (17.77) for a field $F_{\mu\rho\gamma}$ given by Eq. (17.78). The constant in the formula (17.77) is a square of mass of a skewon particle (in a linear approximation of course). This is of course our λ_{c0} . Thus we have

$$m_{\text{skewon}} = \sqrt{\lambda_{c0}} \simeq 10^{-5} \text{ eV}. \quad (19.234)$$

Using Eqs (19.3) and (19.234) we get a relation between a skewon mass and a quintessence mass particle

$$m_0 = \frac{1}{2} \sqrt{n(n+2)} m_{\text{skewon}}. \quad (19.234a)$$

Let us consider a primordial spectrum of gravitational waves in our approach. This spectrum is flat for our Hubble constant is really constant

$$P_{\text{grav}} = \frac{2}{M_{\text{pl}}^2} \left(\frac{H_0}{2\pi} \right)^2 \quad (19.235)$$

(see Ref. [110]). Thus

$$n_{\text{gr}} = 1. \quad (19.236)$$

Now we start to examine quintessence fluctuations caused by fluctuations of a metric. In order to do it we follow Ref. [110] to perturb a spatial part of a metric

$$ds^2 = R^2(\tau) \left[-d\tau^2 + \left(\delta_{ij} + 2\tilde{E}_{ij}^T \right) dx^i dx^j \right], \quad (19.237)$$

τ is a conformal time.

If we expand Einstein equations for (19.237) up to linear terms, we get

$$\frac{d^2 E_{ij\vec{k}}^T}{d\tau^2} + 2RH_K \frac{dE_{ij\vec{k}}^T}{d\tau} + k^2 E_{ij\vec{k}}^T = 0, \quad K = 0, 1, \quad (19.238)$$

where $\tilde{E}_{ij}^T$ is a spatial perturbation of the metric, $R(\tau)$ is a scale factor depending on a conformal time, $E_{ij\vec{k}}^T$ is a $\vec{k}$ Fourier component $\tilde{E}_{ij}^T$

$$\begin{aligned} \tilde{E}_{ij}^T &= \frac{1}{(2\pi)^{3/2}} \int E_{ij\vec{k}}^T e^{-i\vec{k}\vec{r}} d^3\vec{k} \\ \vec{k} &= (k_1, k_2, k_3), \quad |\vec{k}|^2 = k^2, \quad \vec{r} = (x, y, z). \end{aligned} \quad (19.239)$$

The important quantity is an amplitude of gravitational wave corresponding to E_{ij}^T , i.e.

$$h_{ij\vec{k}} = RE_{ij\vec{k}}^T. \quad (19.240)$$

H_K is a Hubble constant. We take $K = 1$. The relation between a conformal time τ and ordinary time t is given by

$$R(\tau) = -\frac{1}{H_1\tau}, \quad -\infty < \tau < 0, \quad (19.241)$$

$$R(t) = R_1 e^{H_1 t}. \quad (19.242)$$

For $h_{ij\vec{k}}$ one gets

$$\frac{d^2 h_{ij\vec{k}}}{d\tau^2} - \frac{2}{\tau^2} \frac{dh_{ij\vec{k}}}{d\tau} + k^2 h_{ij\vec{k}} = 0. \quad (19.243)$$

This equation can be easily solved. For further investigations we need only

$$h_{\vec{k}} = h_{i\vec{k}}^i \quad (19.244)$$

$$h_{\vec{k}} = A_{\vec{k}}(\tau|k| \cos(\tau|k|) - \sin(\tau|k|)) + B_{\vec{k}}(\tau|k| \sin(\tau|k|) + \cos(\tau|k|)) \quad (19.245)$$

and especially

$$\frac{dh_{\vec{k}}}{d\tau} = A_{\vec{k}}\tau \sin(\tau|k|) + B_{\vec{k}}\tau \cos(\tau|k|) \quad (19.246)$$

where $A_{\vec{k}}$ and $B_{\vec{k}}$ are constants.

We need also a quintessence field time dependence (e.g. in a slow roll approximation). We use our solution from Sec. 14 (see Eq. (14.416)), making some simplifications and changing t into τ :

$$\Psi = \ln \left(\sqrt{\frac{n+2}{n}} \sqrt{\frac{\beta}{|\gamma|}} \right) + \ln \left(\frac{1 - C(-\tau)^{\tilde{\kappa}}}{1 + C(-\tau)^{\tilde{\kappa}}} \right). \quad (19.247)$$

Moreover what we really need is a derivative of Ψ .

$$\Psi' = \frac{d\Psi}{d\tau} = \frac{2C\tilde{\kappa}(-\tau)^{\tilde{\kappa}-1}}{1 - C^2(-\tau)^{2\tilde{\kappa}}} \quad (19.248)$$

where

$$\tilde{\kappa} = \frac{\overline{B}\sqrt{2n}}{H_1} \quad (19.248a)$$

$$C = e^{\overline{B}\sqrt{2n}t_0} (R_1 H_1)^{\overline{B}\sqrt{2n}/H_1} = e^{\overline{B}\sqrt{2n}t_0} (R_1 H_1)^{\tilde{\kappa}}. \quad (19.248b)$$

Now we proceed to a quintessence fluctuation equation (see Refs [111,133]):

$$\delta\ddot{Q}_{\vec{k}} + 3H_1\delta\dot{Q}_{\vec{k}} + (c_s^2 k^2 + R^2 U''(Q)) \delta Q_{\vec{k}} = \frac{1}{2\overline{\beta}} h'_k \cdot \Psi'. \quad (19.249)$$

One gets

$$\begin{aligned} & \delta\ddot{Q}_{\vec{k}} + 3H_1\delta\dot{Q}_{\vec{k}} + (c_s^2 k^2 + R^2 U''(Q)) \delta Q_{\vec{k}} \\ &= \frac{C\tilde{\kappa}}{\overline{\beta}} \left(A_{\vec{k}} \frac{(-\tau)^{\tilde{\kappa}-1} \sin(\tau|k|)}{1 - C^2(-\tau)^{2\tilde{\kappa}}} + B_{\vec{k}} \frac{(-\tau)^{\tilde{\kappa}-1} \cos(\tau|k|)}{1 - C^2(-\tau)^{2\tilde{\kappa}}} \right) \\ &= \frac{C\tilde{\kappa}}{\overline{\beta}} C_{\vec{k}} \frac{\sin(\tau|k| + \delta_{\vec{k}})}{1 - C^2(-\tau)^{2\tilde{\kappa}}} (-\tau)^{2\tilde{\kappa}}. \end{aligned} \quad (19.250)$$

$\overline{\beta}$ is a normalization constant in a definition of $Q = \frac{\Psi}{\overline{\beta}}$, $\frac{1}{\overline{\beta}^2} = \frac{8\pi|\overline{M}|}{m_{\text{pl}}^2}$ (see Eqs (14.139) and (19.28)).

Let us consider $\delta Q_{\vec{k}}$ as a Fourier component of a q_0 field subject to the following initial conditions:

$$\frac{dq_{0\vec{k}}(0)}{d\tau} = 0 = q_{0\vec{k}}(0). \quad (19.251)$$

In this case one gets

$$\frac{d^2 q_{0\vec{k}}}{d\tau^2} + 3H_1 \frac{dq_{0\vec{k}}}{d\tau} + \left(c_s^2 k^2 + \frac{m_0^2}{H_1^2 \tau^2} \right) q_{0\vec{k}} = \frac{(-\tau)^{\tilde{\kappa}-1} C\tilde{\kappa}}{\overline{\beta} (1 - C^2(-\tau)^{2\tilde{\kappa}})} C_{\vec{k}} \sin(\tau|k| + \delta_{\vec{k}}), \quad (19.252)$$

c_s is a speed of sound in a quintessence, $C_{\vec{k}}$ and $\delta_{\vec{k}}$ are constants. Let us notice that $\tau = 0$ corresponds to $t \rightarrow \infty$. Thus in some sense we have asymptotic conditions for gravitational waves.

For further convenience it is good to change τ into $-\tau$. In this way we get

$$\frac{d^2 q_{0\vec{k}}}{d\tau^2} - 3H_1 \frac{dq_{0\vec{k}}}{d\tau} + \left(c_s^2 k^2 + \frac{m_0^2}{H_1^2 \tau^2} \right) q_{0\vec{k}} = - \frac{C_{\vec{k}} \tilde{\kappa} \tau^{\tilde{\kappa}-1}}{\beta (1 - C^2 \tau^{2\tilde{\kappa}})} C_{\vec{k}} \sin(\tau|k| - \delta_{\vec{k}}), \quad (19.253)$$

$$\tau \in (0, +\infty), \quad \tau = \frac{1}{R_1 H_1} e^{-H_1 t}. \quad (19.254)$$

The fluctuations (the field q_0) of a quintessence Q are driven by gravitational waves and a quintessence field Q . It is easy to see that we are interested in solutions for small τ (large t). In this case one gets

$$\frac{d^2 q_{0\vec{k}}}{d\tau^2} - 3H_1 \frac{dq_{0\vec{k}}}{d\tau} + \frac{m_0^2}{H_1^2 \tau^2} q_{0\vec{k}} = - \frac{C_{\vec{k}} \tilde{\kappa} \tau^{\tilde{\kappa}-1}}{\beta} C_{\vec{k}} \sin(\tau|k| - \delta_{\vec{k}}). \quad (19.255)$$

Let us calculate a constant $\tilde{\kappa}$. One gets

$$\begin{aligned} \tilde{\kappa} &= \frac{\overline{B} \sqrt{2n}}{H_1} = \left(\frac{n}{n+2} \right)^{n/4} \left(\frac{|\gamma|}{\beta} \right)^{n/4} \frac{\sqrt{\overline{M}}}{m_{\text{pl}}} \cdot 4(n+2)|\gamma|^{1/2} \\ &= 8\sqrt{3\pi n}(n+2)\sqrt{\overline{M}} \left(\frac{n}{n+2} \right)^{n/4} \left(\frac{m_{\tilde{A}}}{m_{\text{pl}}} \right)^{(n+2)/2} \frac{|\tilde{P}|^{1/2}}{\alpha_s^{(n+1)}} \left(\frac{|\tilde{P}|}{\tilde{R}(\tilde{\Gamma})} \right)^{n/2}. \end{aligned} \quad (19.256)$$

If we use our simplified model with $n = 14$, $\alpha_s^2 = \alpha_{\text{em}}$, $m_{\tilde{A}} = m_{\text{EW}}$, we get

$$\tilde{\kappa} \cong 10^{92}. \quad (19.257)$$

Thus this is really a large number.

Moreover for small τ we have

$$q_{o\vec{k}}(\tau) \simeq 0, \quad \frac{dq_{0\vec{k}}}{d\tau} \simeq 0. \quad (19.258)$$

In this way we arrive to an equation

$$\frac{d^2 q_{0\vec{k}}}{d\tau^2} = - \frac{C_{\vec{k}} \tilde{\kappa}}{\beta} (B_{\vec{k}} k \tau^{\tilde{\kappa}} + A_{\vec{k}} \tau^{\tilde{\kappa}-1}) \quad (19.259)$$

and finally

$$q_{0\vec{k}}(\tau) = - \frac{B_{\vec{k}} |k| C_{\vec{k}} \tilde{\kappa}}{\beta (\tilde{\kappa} + 2)(\tilde{\kappa} + 1)} \tau^{\tilde{\kappa}+2} - \frac{A_{\vec{k}} C_{\vec{k}} \tilde{\kappa}}{\beta (\tilde{\kappa} + 1) \tilde{\kappa}} \tau^{\tilde{\kappa}+1} \quad (19.260)$$

or (taking under consideration that $\tilde{\kappa}$ is very large)

$$q_{0\vec{k}}(\tau) = -\frac{C}{\beta\tilde{\kappa}} (B_{\vec{k}}k\tau^{\tilde{\kappa}} + A_{\vec{k}}\tau^{\tilde{\kappa}-1}) \quad (19.261)$$

$$q_{0\vec{k}}(\tau) = -\frac{C}{\beta\tilde{\kappa}} \left(A_{\vec{k}}(R_1 H_1)^{-\overline{B}\sqrt{2n}/H_1} e^{-\overline{B}\sqrt{2nt}} \right. \\ \left. + k B_{\vec{k}}(R_1 H_1)^{-((\overline{B}\sqrt{2n}/H_1)-1)} e^{-\overline{B}\sqrt{2nt}} \cdot e^{H_1 t} \right) \quad (19.262)$$

$$q_0(t, \vec{r}) = \int q_{0\vec{k}}(t) e^{i\vec{k}\vec{r}} d\vec{k}. \quad (19.263)$$

One gets

$$q_0(t, \vec{r}) = -\frac{C}{\beta\tilde{\kappa}} \left(g(\vec{r}) \left(\frac{H_1}{R_1} \right)^{\overline{B}\sqrt{2n}/H_1} e^{-\overline{B}\sqrt{2nt}} \right. \\ \left. + f(\vec{r}) \left(\frac{H_1}{R_1} \right)^{(\overline{B}\sqrt{2n}/H_1)-1} e^{-2\overline{B}\sqrt{2nt}} \cdot e^{H_1 t} \right) \quad (19.264)$$

where

$$f(\vec{r}) = \int A_{\vec{k}} e^{i\vec{k}\vec{r}} d^3\vec{k} \quad (19.265)$$

$$g(\vec{r}) = \int |k| B_{\vec{k}} e^{i\vec{k}\vec{r}} d^3\vec{k}. \quad (19.266)$$

$g(\vec{r})$ and $f(\vec{r})$ characterize a spatial dependence of gravitational waves background. For large t we have to do with $g(\vec{r})$, i.e.

$$q_0(t, \vec{r}) = -\frac{C}{\beta\tilde{\kappa}} g(\vec{r}) (R_1 H_1)^{-((\overline{B}\sqrt{2n}/H_1)-1)} \cdot e^{-\overline{B}\sqrt{2nt}} e^{H_1 t}. \quad (19.267)$$

In this way one gets from an energy-momentum tensor for q_0

$$T_{\mu\nu} = \partial_\mu q_0 \cdot \partial_\nu q_0 - \frac{1}{2} \eta_{\mu\nu} (\partial^\alpha q_0 \cdot \partial_\alpha q_0 + m_0^2 q_0^2) \quad (19.268)$$

$$\rho_{q_0} = T_{44} = \dot{q}_0^2 - \frac{1}{2} \left(\dot{q}_0^2 - |\vec{\nabla} q_0|^2 - m_0^2 q_0^2 \right) \\ = \frac{1}{2} q_0^2 \left(\left(H_1 - \overline{B}\sqrt{2n} \right)^2 + |\vec{\nabla} \ln g(r)|^2 + m_0^2 \right) \\ = \frac{\rho_{\text{gr}} 2n \overline{B}^2 R_1^2 \left(\left(H_1 - \overline{B}\sqrt{2n} \right)^2 + |\vec{\nabla} \ln g(r)|^2 + m_0^2 \right)}{\beta^2 e^{2\overline{B}\sqrt{2nt}}} e^{2H_1 t} e^{\overline{B}\sqrt{2nt_0}} \quad (19.269)$$

where

$$\rho_{\text{gr}} = \frac{1}{2} g^2 \quad (19.270)$$

is an energy density of primordial gravitational waves.

For $\tilde{\kappa} = \frac{\overline{B}\sqrt{2n}}{H_1}$ is a large number,

$$\overline{B}\sqrt{2n} \gg H_1 \quad (19.271)$$

and ρ_{q_0} is going to zero if $t \rightarrow \infty$. This density will be frozen (no dependence on time) at an end of the second de Sitter phase $t = t_{\text{end}}^{\text{II}}$, $t_0 = t_{\text{initial}}^{\text{II}}$. In this way one gets

$$\frac{\rho_{q_0}}{\rho_{\text{gr}}} = \frac{2n\overline{B}^2 R_1^2 \left((H_1 - \overline{B}\sqrt{2n})^2 + |\vec{\nabla} \ln g(r)|^2 + m_0^2 \right)}{\overline{\beta}^2 \exp\left(\frac{2\overline{B}\sqrt{2n}N_1}{H_1}\right)} e^{2N_1} \quad (19.272)$$

where N_1 is an amount of inflation during the second de Sitter phase. If $g(\vec{r}) = \text{const.}$, ρ_{q_0} is isotropic and homogeneous.

The function $g(\vec{r})$ can be written in a form

$$g(\vec{r}) = g_0 + \delta g(\vec{r}) \quad (19.273)$$

where $\delta g(\vec{r})$ is a small deviation and $g_0 = \text{const.}$ One gets

$$\vec{\nabla} (\ln g(\vec{r})) \simeq \frac{1}{g_0} \vec{\nabla} \delta g(\vec{r}). \quad (19.274)$$

In this way one writes

$$\rho_{q_0} = \rho_{q_0}^0 + \delta \rho_{q_0} \quad (19.275)$$

where

$$\begin{aligned} \rho_{q_0}^0 &= \frac{2n\overline{B}^2 R_1^2 \rho_{\text{gr}} \left((H_1 - \overline{B}\sqrt{2n})^2 + m_0^2 \right)}{\overline{\beta}^2 \exp\left(\frac{2\overline{B}\sqrt{2n}N_1}{H_1}\right)} e^{2N_1} \\ \delta \rho_{q_0} &= |\vec{\nabla} g(\vec{r})|^2 \frac{2n\overline{B}^2 R_1^2 \rho_{\text{gr}}}{\overline{\beta}^2 \exp\left(\frac{2\overline{B}\sqrt{2n}N_1}{H_1}\right)} e^{2N_1} \end{aligned} \quad (19.276)$$

According to our ideas the energy ρ_{q_0} should be stored as a gas of quintessence particles. The measurement of a low frequency speed of sound and a high frequency speed of sound can give us η and its spatial variation. In this way we get also a square length of a gradient of $\delta g(\vec{r})$. If we suppose that ρ_{gr} is stored as a gas of gravitons (massless particles with two polarization states) then one can write

$$p_{\text{gr}} = \rho_{\text{gr}} \left(\frac{T_{\text{gr}}}{T_0} \right)^4 \quad (19.277)$$

(see Eq. (19.231)).

Probably we should also consider a gas of skewons. However, this is a different story. Taking into account (19.235) and (19.236) we should expect $\delta \rho_{q_0}$ as extremely small and we can neglect it in our theory. Moreover, ρ_{gr} is very important. Let us consider Eq. (19.253) for these values of τ for which the variable t is close to $t_{\text{end}}^{\text{I}}$ (the

end of the first de Sitter phase or beginning of the second de Sitter phase). In this way one writes

$$\tau = \frac{1}{H_1 R_1} e^{-H_1 t_{\text{end}}^I} + \xi = \tau_0 + \xi, \quad 0 < \xi \ll 1. \quad (19.278)$$

Taking into account that ξ is very small we arrive to the following equation

$$\frac{d^2 q_{0\vec{k}}}{d\xi^2} - 3H_1 \frac{dq_{0\vec{k}}}{d\xi} + \left(c_s^2 k^2 + \frac{m_0^2}{H_1^2 \tau_0^2} \right) q_{0\vec{k}} = -\frac{C_{\vec{k}} \tau_0}{2\xi \beta} \sin(\xi|k| + \bar{\delta}_{\vec{k}}) \quad (19.279)$$

where

$$\bar{\delta}_{\vec{k}} = \tau_0 |k| - \delta_{\vec{k}}. \quad (19.280)$$

We have also initial conditions

$$\frac{dq_{0\vec{k}}}{d\xi}(0) = q_{0\vec{k}}(0) = 0. \quad (19.281)$$

Eq. (19.279) can be solved by a Laplace transform method in the case of $\bar{\delta}_{\vec{k}} = 0$.

Let

$$c_s^2 k^2 + \frac{m_0^2}{H_1^2 \tau_0^2} = b^2. \quad (19.282)$$

One gets

$$s^2 \bar{q}_{0\vec{k}}(s) - 3H_1 s \bar{q}_{0\vec{k}}(s) + b^2 \bar{q}_{0\vec{k}}(s) = -\frac{C_{\vec{k}} \tau_0}{2\beta} \text{arctg}\left(\frac{|k|}{s}\right) \quad (19.283)$$

$$\bar{q}_{0\vec{k}}(s) = -\frac{C_{\vec{k}} \tau_0}{2\beta} \frac{\text{arctg}\left(\frac{|k|}{s}\right)}{s^2 - 3H_1 s + b^2} \quad (19.284)$$

and

$$q_{0\vec{k}}(\xi) = -\frac{C_{\vec{k}} \tau_0}{2\beta} \int_0^\infty \frac{\text{arctg}\left(\frac{|k|}{s}\right) e^{s\xi}}{s^2 - 3H_1 s + b^2} ds. \quad (19.285)$$

However, we cannot get any compact form of this solution. Moreover for we are looking for a solution around zero, for $\bar{\delta}_{\vec{k}} = 0$ we get

$$-\frac{C_{\vec{k}} \tau_0}{2\xi} \sin(\xi|k|) \simeq -\frac{C_{\vec{k}} \tau_0 |k|}{2}. \quad (19.286)$$

In this case the Laplace transform method is successful:

$$q_{0\vec{k}}(\xi) = -\frac{C_{\vec{k}} \tau_0 |k|}{2\beta} \int_0^\infty \frac{e^{s\xi}}{s(s^2 - 3H_1 s + b^2)} ds \quad (19.287)$$

$$q_{0\vec{k}}(\xi) = -\frac{C_{\vec{k}} \tau_0 |k|}{2\beta} \left(\frac{1}{b^2} + \frac{e^{s_1 \xi}}{s_1(s_1 - s_2)} + \frac{e^{s_2 \xi}}{s_2(s_2 - s_1)} \right) \quad (19.287a)$$

where s_1 and s_2 are roots of the polynomial

$$s^2 - 3H_1s + b^2 = 0. \quad (19.288)$$

Moreover ξ is small. Thus we write

$$e^{s_i \xi} \cong 1 + s_i \xi, \quad i = 1, 2, \quad (19.289)$$

and get

$$q_{0\vec{k}}(\xi) = -\frac{C_{\vec{k}}\tau_0|k|}{2b^2\bar{\beta}} \left(1 + \frac{3H_1}{\sqrt{\Delta}}\right) \quad (19.290)$$

if

$$\Delta = 9H_1^2 - 4b^2 = 9H_1^2 - 4\left(c_s^2 k^2 + \frac{m_0^2}{H_1^2 \tau_0^2}\right) > 0. \quad (19.291)$$

If $\Delta < 0$, one gets

$$q_{0\vec{k}}(\xi) = -\frac{C_{\vec{k}}\tau_0|k|}{2b^2\bar{\beta}} \left(1 + \exp\left(\frac{3}{2}H_1\xi\right) \left(\frac{3H_1}{\omega} \sin(\omega\xi) - \cos(\omega\xi)\right)\right) \quad (19.292)$$

where

$$4\omega^2 = -\Delta = 4\left(c_s^2 k^2 + \frac{m_0^2}{H_1^2 \tau_0^2}\right) - 9H_1^2 > 0. \quad (19.293)$$

For small ξ one gets

$$q_{0\vec{k}}(\xi) = -\frac{3C_{\vec{k}}\tau_0|k|H_1}{4b^2\bar{\beta}} \xi, \quad (19.294)$$

the dependence on t is as follows:

$$\xi = \frac{-1}{R_1 H_1} (e^{-H_1 t} - e^{-H_1 t_{\text{end}}^{\text{I}}}). \quad (19.295)$$

Moreover t is close to $t_{\text{end}}^{\text{I}}$ and we get

$$\xi \simeq \frac{e^{-H_1 t_{\text{end}}^{\text{I}}}}{R_1} (t - t_{\text{end}}^{\text{I}}) \quad (19.296)$$

for t very close to $t_{\text{end}}^{\text{I}}$.

In this way we get fluctuations of a quintessence in the moment of the first order phase transition of the first de Sitter phase to the second de Sitter phase.

Finally one gets

$$\begin{aligned} q_{0\vec{k}}(t) = & -\frac{C_{\vec{k}}|k|e^{-H_1 t_{\text{end}}^{\text{I}}}}{2H_1 R_1 \left(c_s^2 |k|^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^{\text{I}}}\right) \bar{\beta}} \\ & \times \left(1 + \frac{3H_1}{\sqrt{9H_1^2 - 4(c_s^2 |k|^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^{\text{I}}})}}\right) \end{aligned} \quad (19.297)$$

for

$$|k| < \frac{1}{2c_s} \sqrt{9H_1^2 - 4R_1^2 e^{2H_1 t_{\text{end}}^I} m_0^2} = k_0 \quad (19.298)$$

with a condition

$$3H_1 > 2R_1 e^{H_1 t_{\text{end}}^I} m_0 \quad (19.299)$$

and

$$q_{0\vec{k}}(t) = -\frac{3C_{\vec{k}} e^{-2H_1 t_{\text{end}}^I} (t - t_{\text{end}}^I) |k|}{4R_1^2 \left(c_s^2 k^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^I} \right) \bar{\beta}}. \quad (19.300)$$

for $k > k_0$.

One finds

$$\begin{aligned} q_0(\vec{r}, t) = & -\frac{e^{-H_1 t_{\text{end}}^I}}{2\bar{\beta}R_1} \left(\frac{1}{H_1} \int_{|k| < k_0} d^3\vec{k} e^{i\vec{k}\vec{r}} \frac{C_{\vec{k}} |k|}{\left(c_s^2 k^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^I} \right)} \right. \\ & \times \left(1 + \frac{3H_1}{\sqrt{9H_1^2 - 4 \left(c_s^2 k^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^I} \right)}} \right) \\ & \left. + \frac{3(t - t_{\text{end}}^I)}{2R_1} e^{-H_1 t_{\text{end}}^I} \int_{|k| > k_0} d^3\vec{k} e^{i\vec{k}\vec{r}} \frac{C_{\vec{k}} |k|}{\left(c_s^2 k^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^I} \right)} \right) \end{aligned} \quad (19.301)$$

Equation (19.301) gives us space-time fluctuations of a quintessence field after a phase transition from the first to the second de Sitter phase. Let us notice that (19.297) is singular for $k \rightarrow k_0$.

In an approach developed here we consider Eq. (19.250) with zero initial conditions in various approximations. Moreover there is a different approach. In this approach we write a solution to Eq. (19.250)

$$q_{0\vec{k}}(t) = \tilde{q}_{0\vec{k}}(t) + \bar{q}_{0\vec{k}}(t) \quad (19.302)$$

where $\tilde{q}_{0\vec{k}}(t)$ is a solution of a homogeneous equation (a general integral) and $\bar{q}_{0\vec{k}}(t)$ is a special solution to inhomogeneous equation. This different approach consists in taking $\tilde{q}_{0\vec{k}}(t) \equiv 0$ and considering only $\bar{q}_{0\vec{k}}(t)$. In this case we get nonzero initial conditions for quintessence fluctuations. However, this approach is not unambiguous. Our approach is unambiguous. The zero initial conditions are in some sense distinguished.

Let us calculate an energy density of this scalar field. In order to do this let us write $q_0(\vec{r}, t)$ in a form

$$q_0(\vec{r}, t) = A(\vec{r}) + B(\vec{r})(t_{\text{end}}^I - t) \quad (19.303)$$

where

$$A(\vec{r}) = -\frac{e^{-H_1 t_{\text{end}}^{\text{I}}}}{2\bar{\beta}R_1 H_1} \int_{|\vec{k}| < k_0} d^3\vec{k} e^{i\vec{k}\vec{r}} \frac{C_{\vec{k}}|k|}{(c_s^2 k^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^{\text{I}}})} \times \left(1 + \frac{3H_1}{\sqrt{9H_1^2 - 4(c_s^2 k^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^{\text{I}}})}} \right) \quad (19.304)$$

$$B(\vec{r}) = -\frac{3e^{-2H_1 t_{\text{end}}^{\text{I}}}}{4\bar{\beta}R_1^2} \int_{|\vec{k}| > k_0} d^3\vec{k} e^{i\vec{k}\vec{r}} \frac{C_{\vec{k}}|k|}{(c_s^2 k^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^{\text{I}}})}. \quad (19.305)$$

One gets from the Eq. (19.268)

$$\rho_{q_0} = T_{44} = \frac{1}{2} (C(\vec{r}) + D(\vec{r})(t_{\text{end}}^{\text{I}} - t) + E(\vec{r})(t_{\text{end}}^{\text{I}} - t)^2) \quad (19.306)$$

where

$$C(\vec{r}) = B^2(\vec{r}) + m_0^2 A^2(\vec{r}) + |\vec{\nabla}(A(\vec{r}))|^2 \quad (19.307)$$

$$D(\vec{r}) = 2 \left(\vec{\nabla}(A(\vec{r})) \cdot \vec{\nabla}(B(\vec{r})) + m_0^2 A(\vec{r})B(\vec{r}) \right) \quad (19.308)$$

$$E(\vec{r}) = \left(|\vec{\nabla}(B(\vec{r}))|^2 + m_0^2 B^2(\vec{r}) \right). \quad (19.309)$$

All these formulae are satisfied for t close to $t_{\text{end}}^{\text{I}}$. In general ρ_{q_0} is anisotropic and inhomogeneous. For $t_{\text{end}}^{\text{I}} - t$ is very small we get

$$\rho_{q_0} = \frac{1}{2} (C(\vec{r}) + D(\vec{r})(t_{\text{end}}^{\text{I}} - t)). \quad (19.310)$$

It is interesting to calculate a time $t(\vec{r})$ for which ρ_{q_0} is zero ($t(\vec{r})$ depends on a space point). One gets (neglecting gradients of $A(\vec{r})$ and $B(\vec{r})$)

$$t(\vec{r}) = t_{\text{end}}^{\text{I}} + \frac{1}{2m_0^2} \left(\bar{f}(\vec{r}) + \frac{m_0^2}{\bar{f}(\vec{r})} \right) \quad (19.311)$$

where

$$\bar{f}(\vec{r}) = \frac{3H_1}{2R_1} \cdot \frac{e^{-H_1 t_{\text{end}}^{\text{I}}} \int_{|\vec{k}| > k_0} d^3\vec{k} e^{i\vec{k}\vec{r}} \frac{C_{\vec{k}}|k|}{(c_s^2 k^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^{\text{I}}})}}{\int_{|\vec{k}| < k_0} d^3\vec{k} e^{i\vec{k}\vec{r}} \frac{C_{\vec{k}}|k|}{(c_s^2 k^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^{\text{I}}})} \left(1 + \frac{3H_1}{\sqrt{9H_1^2 - 4(c_s^2 k^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^{\text{I}}})}} \right)}}. \quad (19.312)$$

The conclusion from these calculations is that after a very short time an energy density of quintessence fluctuations is going to zero (almost zero) and probably is frozen on a very low level. A time of this process depends on a space point. Moreover, after

sufficiently long time (which is really very short) we get a homogeneity and isotropy, as far as our rough calculations advise us. Let us come back to Eq. (19.253). Our solutions obtained here are for large t and t close to $t_{\text{end}}^{\text{I}}$. One can try to match them. It seems reasonable to suppose that $\rho_{q_0}^0$ (see Eq. (19.276)) is equal to this very low level energy density, mentioned above.

The quintessence field $q_0(\vec{r}, t)$ considered above can be a source of fluctuations of an effective gravitational constant G_{eff} .

$$G_{\text{eff}} = G_N \exp \left(-(n+2)\bar{\beta}q_0(\vec{r}, t) \right) \quad (19.313)$$

(see Eq. (19.28)). If we use Eq. (19.267), we get

$$G_{\text{eff}} = G_N \exp \left(- \frac{\sqrt{2}(n+2)\sqrt{\rho_{q_0}}\bar{\beta}}{\sqrt{((H_1 - \bar{B}\sqrt{2n})^2 + |\vec{\nabla} \ln g(\vec{r})|^2 + m_0^2)}} \right) \quad (19.314)$$

and we connect contemporary fluctuations of an effective gravitational constant to an energy density of quintessence fluctuations. Taking a quintessence field from (19.303) we get

$$G_{\text{eff}} = G_N \exp \left(-(n+2)\bar{\beta} \left(A(\vec{r}) + B(\vec{r})(t_{\text{end}}^{\text{I}} - t) \right) \right). \quad (19.315)$$

In this way an effective gravitational constant depends on a speed of a sound in a quintessence.

In order to get some comparison let us consider a different approximation of a quintessence evolution given by Eqs (14.402) and (14.405),

$$Q(t) = Q_0 + \frac{\varphi_0}{\beta} \sin(\omega_0 t) \quad (19.316)$$

where

$$\omega_0^2 = \frac{M_{\text{pl}}}{2\bar{M}} (n+2) \left(\frac{n}{n+2} \right)^{n/2} |\gamma| \left(\frac{|\gamma|}{\beta} \right)^{n/2} \quad (19.317)$$

or

$$\begin{aligned} \omega_0 &= \frac{m_{\tilde{A}}\sqrt{m_{\text{pl}}}}{2\alpha_s^{(n+1)/2}\sqrt{2H}\sqrt{|\bar{M}|}} \left(\frac{m_{\tilde{A}}}{m_{\text{pl}}} \right)^{n/2} (n+2)^{1/2} \\ &\times \left(\frac{n}{n+2} \right)^{n/4} |\tilde{P}|^{1/2} \left(\frac{|\tilde{P}|}{\tilde{R}(\tilde{r})} \right)^{n/4}. \end{aligned} \quad (19.318)$$

In terms of a conformal time τ one gets

$$Q(\tau) = Q_0 - \frac{\varphi_0}{\beta} \sin \left(\frac{\omega_0}{H_1} \ln(-\tau R_1 H_1) \right) \quad (19.319)$$

$$\frac{dQ}{d\tau} = -\frac{\varphi_0\omega_0}{\beta H_1} \cdot \frac{1}{\tau} \cos \left(\frac{\omega_0}{H_1} \ln(-\tau R_1 H_1) \right) \quad (19.320)$$

Let us notice that φ_0 is an arbitrary constant.

Using Eq. (19.246) and Eq. (19.249) one gets

$$\begin{aligned} \delta\ddot{Q}_{\vec{k}} - 3H_1\delta\dot{Q}_{\vec{k}} + \left(c_s^2|k|^2 + \frac{m_0^2}{H_1^2\tau^2}\right)\delta Q_{\vec{k}} \\ = \frac{\varphi_0\omega_0}{2H_1\bar{\beta}} (A_{\vec{k}}\sin(\tau|k|) - B_{\vec{k}}\cos(\tau|k|)) \cos\left(\frac{\omega_0}{H_1}\ln(\tau R_1 H_1)\right) \end{aligned} \quad (19.321)$$

where we change $\tau \rightarrow -\tau$, $\tau \in (0, +\infty)$.

Let us consider $\delta Q_{\vec{k}}$ as quintessence field $q_{0\vec{k}}$ (as before) and let us consider equation (19.321) for

$$\tau = \tau_0 + \xi = \frac{1}{H_1 R_1} e^{-H_1 t_{\text{end}}^{\text{I}}} + \xi, \quad 0 < \xi \ll 1 \quad (19.322)$$

(also as before).

We can consider Eq. (19.321) only for τ given by (19.322), for our approximation works only here.

Changing a variable from τ to ξ and doing some simplifications we get

$$\begin{aligned} \frac{d^2 q_{0\vec{k}}}{d\xi^2} - 3H_1 \frac{dq_{0\vec{k}}}{d\xi} + \left(c_s^2|k|^2 + \frac{m_0^2}{H_1^2\tau_0^2}\right) q_{0\vec{k}} \\ = -\frac{\varphi_0\omega_0}{2H_1\bar{\beta}} C_{\vec{k}} \cos(\omega_0 t_{\text{end}}^{\text{I}}) \sin(|k|\xi + \bar{\delta}_{\vec{k}}) \end{aligned} \quad (19.323)$$

where $\bar{\delta}_{\vec{k}} = \tau_0|k| - \delta_{\vec{k}}$ (as before).

For a very small ξ we get

$$\sin(|k|\xi + \delta_{\vec{k}}) \simeq \sin \delta_{\vec{k}} \quad (19.324)$$

and we get Eq. (19.279) for $\bar{\delta}_{\vec{k}} = 0$ and for small ξ , $C_{\vec{k}}$ is expressed in terms of $A_{\vec{k}}$ and $B_{\vec{k}}$,

$$C_{\vec{k}} = \sqrt{A_{\vec{k}}^2 + B_{\vec{k}}^2}. \quad (19.325)$$

In this way using a Laplace transform method for this equation with initial conditions (19.281) one gets

$$q_{0\vec{k}}(\xi) = -F \left(\frac{1}{b^2} + \frac{e^{s_1\xi}}{s_1(s_1 - s_2)} + \frac{e^{s_2\xi}}{s_2(s_2 - s_1)} \right) \quad (19.326)$$

where s_1 and s_2 are roots of the equation (19.288) and

$$F = \frac{\varphi_0\omega_0 C_{\vec{k}} \sin \bar{\delta}_{\vec{k}} \cos(\omega_0 t_{\text{end}}^{\text{I}})}{2H_1\bar{\beta}}. \quad (19.327)$$

In this way we get similar solutions as before

$$q_{0\vec{k}}(t) = -\frac{C_{\vec{k}} \sin \bar{\delta}_{\vec{k}} \cos(\omega_0 t_{\text{end}}^I) \varphi_0 \omega_0}{2\bar{\beta} H_1 \left(c_s^2 |k|^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^I} \right)} \times \left(1 + \frac{3H_1}{\sqrt{9H_1^2 - 4(c_s^2 |k|^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^I})}} \right) \quad (19.328)$$

for $|k|$ satisfying condition (19.298) and

$$q_{0\vec{k}}(t) = -\frac{3\varphi_0 \omega_0 C_{\vec{k}} \sin \bar{\delta}_{\vec{k}} \cos(\omega_0 t_{\text{end}}^I) e^{-H_1 t_{\text{end}}^I}}{4\bar{\beta} R_1 H_1 \left(c_s^2 |k|^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^I} \right)} (t - t_{\text{end}}^I) \quad (19.329)$$

for $k > k_0$ (see Eq. (19.298)).

In this way one gets

$$q_0(\vec{r}, t) = -\frac{\varphi_0 \omega_0 \cos(\omega_0 t_{\text{end}}^I)}{2\bar{\beta} H_1} \left(\int_{|k| < k_0} d^3 \vec{k} e^{i\vec{k}\vec{r}} \frac{C_{\vec{k}} \sin \bar{\delta}_{\vec{k}}}{\left(c_s^2 |k|^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^I} \right)} \times \left(1 + \frac{3H_1}{\sqrt{9H_1^2 - 4(c_s^2 |k|^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^I})}} \right) + \frac{(t - t_{\text{end}}^I)}{2R_1} e^{-H_1 t_{\text{end}}^I} \int_{|k| > k_0} d^3 \vec{k} e^{i\vec{k}\vec{r}} \frac{C_{\vec{k}} \sin \bar{\delta}_{\vec{k}}}{\left(c_s^2 |k|^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^I} \right)} \right) \quad (19.330)$$

Thus we get similar dependence on time as before (see Eq. (19.303)). Now with the functions

$$\bar{A}(\vec{r}) = -\frac{\varphi_0 \omega_0 \cos(\omega_0 t_{\text{end}}^I)}{2\bar{\beta} H_1} \int_{|k| < k_0} d^3 \vec{k} e^{i\vec{k}\vec{r}} \frac{C_{\vec{k}} \sin \bar{\delta}_{\vec{k}}}{\left(c_s^2 |k|^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^I} \right)} \times \left(1 + \frac{3H_1}{\sqrt{9H_1^2 - 4(c_s^2 |k|^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^I})}} \right) \quad (19.331)$$

$$\bar{B}(\vec{r}) = \frac{3\varphi_0 \omega_0 \cos(\omega_0 t_{\text{end}}^I)}{4\bar{\beta} R_1 H_1} e^{-H_1 t_{\text{end}}^I} \int_{|k| > k_0} d^3 \vec{k} e^{i\vec{k}\vec{r}} \frac{C_{\vec{k}} \sin \bar{\delta}_{\vec{k}}}{\left(c_s^2 |k|^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^I} \right)} \quad (19.332)$$

and

$$\begin{aligned} \bar{f}(\vec{r}) = & \frac{3e^{-H_1 t_{\text{end}}^I} \int_{|k|>k_0} d^3\vec{k} e^{i\vec{k}\vec{r}} \frac{C_{\vec{k}} \sin \bar{\delta}_{\vec{k}}}{(c_s^2 |k|^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^I})}}{2R_1 \int_{|k|<k_0} d^3\vec{k} e^{i\vec{k}\vec{r}} \frac{C_{\vec{k}} \sin \bar{\delta}_{\vec{k}}}{(c_s^2 |k|^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^I})}} \\ & \times \left(1 + \frac{3H_1}{\sqrt{9H_1^2 - 4(c_s^2 |k|^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^I})}} \right) \end{aligned} \quad (19.333)$$

in the place of $A(\vec{r})$, $B(\vec{r})$, $f(\vec{r})$ (see Eqs (19.306÷312) and a discussion below which is applicable here).

The general conclusion from our recent calculations is such that a behaviour of quintessence fluctuations does not depend on an evolution law of a quintessence for time close to t_{end}^I .

Let us consider Eq. (14.402) for $\bar{M} < 0$. In this case one gets

$$Q = Q_0 + \varphi_0 \sinh(\omega_0 t) \quad (19.334)$$

where ω_0 is given by Eq. (19.318). It is easy to see that we can repeat all the considerations given above, changing $\sin$ into $\sinh$ and $\cos$ into $\cosh$. In this way one gets

$$\begin{aligned} q_{0\vec{k}}(t) = & -\frac{C_{\vec{k}} \sin \bar{\delta}_{\vec{k}} \cosh(\omega_0 t_{\text{end}}^I) \varphi_0 \omega_0}{2\bar{\beta} H_1 (c_s^2 |k|^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^I})} \\ & \times \left(1 + \frac{3H_1}{\sqrt{9H_1^2 - 4(c_s^2 |k|^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^I})}} \right) \end{aligned} \quad (19.335)$$

for $|k|$ satisfying condition (19.298) and

$$q_{o\vec{k}}(t) = -\frac{3\varphi_0 \omega_0 C_{\vec{k}} \sin \bar{\delta}_{\vec{k}} \cosh(\omega_0 t_{\text{end}}^I) e^{-H_1 t_{\text{end}}^I}}{4\bar{\beta} R_1 H_1 (c_s^2 |k|^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^I})} (t - t_{\text{end}}^I) \quad (19.336)$$

for $k > k_0$ (see Eq. (19.298)).

One finds

$$\begin{aligned}
q_0(\vec{r}, t) = & -\frac{\varphi_0 \omega_0 \cosh(\omega_0 t_{\text{end}}^{\text{I}})}{2\beta H_1} \left(\int_{|\vec{k}| < k_0} d^3 \vec{k} e^{i \vec{k} \vec{r}} \frac{C_{\vec{k}} \sin \bar{\delta}_{\vec{k}}}{\left(c_s^2 |\vec{k}|^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^{\text{I}}} \right)} \right. \\
& \times \left(1 + \frac{3H_1}{\sqrt{9H_1^2 - 4(c_s^2 |\vec{k}|^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^{\text{I}}})}} \right) \\
& \left. + \frac{(t - t_{\text{end}}^{\text{I}})}{2R_1} e^{-H_1 t_{\text{end}}^{\text{I}}} \int_{|\vec{k}| > k_0} d^3 \vec{k} e^{i \vec{k} \vec{r}} \frac{C_{\vec{k}} \sin \bar{\delta}_{\vec{k}}}{\left(c_s^2 |\vec{k}|^2 + R_1^2 m_0^2 e^{2H_1 t_{\text{end}}^{\text{I}}} \right)} \right)
\end{aligned} \tag{19.337}$$

and consequently

$$\bar{\bar{A}}(\vec{r}) = \frac{\cosh(\omega_0 t_{\text{end}}^{\text{I}})}{\cos(\omega_0 t_{\text{end}}^{\text{I}})} \bar{A}(\vec{r}) \tag{19.338}$$

$$\bar{\bar{B}}(\vec{r}) = \frac{\cosh(\omega_0 t_{\text{end}}^{\text{I}})}{\cos(\omega_0 t_{\text{end}}^{\text{I}})} \bar{B}(\vec{r}) \tag{19.339}$$

$$\bar{\bar{f}}(\vec{r}) = \bar{f}(\vec{r}) \tag{19.340}$$

with the same conclusions as for a case with $\bar{M} > 0$. In this case an amount of inflation reads

$$\begin{aligned}
N_1 = \Delta t H_1 &= \frac{H_1}{\omega_0} \left| \text{arsinh} \left(\frac{1}{\varphi_0 n} \right) \right| = \frac{H_1}{\omega_0} \ln \left(\frac{1 + \sqrt{\varphi_0^2 n^2 + 1}}{\varphi_0 n} \right) \\
&= \frac{M_{\text{pl}}}{\sqrt{6|\bar{M}|} (n+2)} \ln \left(\frac{1 + \sqrt{1 + \varphi_0^2 n^2}}{\varphi_0 n} \right)
\end{aligned} \tag{19.341}$$

Let us give simple applications of our inflation theory (which is effectively similar to a hybrid inflation) to CMB anisotropy problem. In order to do this we remind to the reader some notions of CMB anisotropy theory:

$$\frac{\Delta T(\vec{e})}{T} = \sum_{l,m} a_{lm} Y_{l,m}(\vec{e}) \tag{19.342}$$

where $Y_{lm}(\vec{e}) = Y_{lm}(\theta, \varphi)$ are spherical harmonics and a_{lm} are multipole coefficients of CMB. $\frac{\Delta T(\vec{e})}{T}$ means fluctuations of a temperature of CMB measured in a direction of $\vec{e}$ parametrized by two angles θ and φ . A direction of a photon is $\vec{n} = -\vec{e}$.

After averaging one gets

$$\langle a_{lm} a_{l'm'}^* \rangle = \delta_{ll'} \delta_{mm'} C_l. \tag{19.343}$$

For a two point correlation function one gets

$$\left\langle \frac{\Delta T}{T}(\vec{e}_1) \cdot \frac{\Delta T}{T}(\vec{e}_2) \right\rangle_{\vec{e}_1 \cdot \vec{e}_2 = \cos \theta} = \frac{1}{4\pi} \sum_l (2l+1) C_l P_l(\cos \theta) \quad (19.344)$$

(see Refs [110] for more details) where $P_l(\cos \theta)$ is a Legendre polynomial. Thus in order to compare a theory with observations it is necessary to calculate C_l in some region of l (i.e., $2 < l < l_{\max}$). Usually we subtract effects of relative movement and $C_0 = C_1 = 0$.

C_l can have different origins coming from several types of perturbations (i.e. scalar, vector, tensor) (see Refs [110] for details). In our theory we have scalar perturbations due to inflation and tensor perturbations due to gravitational waves. In our three models of Higgs' field evolution we get a power spectrum of primordial scalar fluctuations (see Eqs (18.87), (18.69), (18.10)) and a power spectrum for tensorial fluctuations (gravitational waves fluctuations) Eq. (19.235). The tensor fluctuations are scale invariant

$$n_T = n_{\text{gr}} - 1 = 0.$$

In the case of scalar fluctuations we calculate n_s and $\frac{dn_s}{d \ln K}$ (see Eqs (18.14), (18.15), (18.72), (18.74), (18.90), (18.91)) and we find conditions for $n_s = 1$. Thus we get (under some conditions) a Harrison-Zel'dovich spectrum. Moreover $\frac{dn_s}{d \ln K}$ can be in general nonzero and it is possible to use it to compare with observations in order to falsificate some models. In general we get in three cases

$$P_R^s(K) = P_0^s K^{n_s-1}. \quad (19.345)$$

Thus via a standard procedure (see Refs [110]) one gets

$$C_l^s = \frac{(P_0^s)^2}{9} \cdot \frac{\Gamma(3-n_s)\Gamma(l-\frac{1}{2}+\frac{n_s}{2})}{2^{3-n_s}\Gamma^2(2-\frac{n_s}{2})\Gamma(l+\frac{5}{2}-\frac{n_s}{2})} \quad (19.346)$$

where $n_s \simeq 1$.

If we take $n_s = 1$ we get

$$l(l+1)C_l^s = \text{const.} \quad (19.347)$$

which is in an agreement with observational data. In the case of tensorial (gravitational waves) perturbations one gets

$$P^T(K) = P_0^{\text{gr}} K^{n_{\text{gr}}-1} = P_0^{\text{gr}} K^{n_T} \quad (19.348)$$

and by a standard procedure

$$C_l^T = C_l^{\text{gr}} = (P_0^{\text{gr}})^2 \cdot \frac{(l+2)!}{(l-2)!} \cdot \frac{\Gamma(6-n_T)\Gamma(l-2+\frac{n_T}{2})}{2^{6-n_T}\Gamma^2(\frac{7}{2}-n_T)\Gamma(l+4-\frac{n_T}{2})}. \quad (19.349)$$

For $n_T = 0$ or $n_{\text{gr}} = 1$ one gets

$$C_l^{\text{gr}} = (P_0^{\text{gr}})^2 \cdot \frac{8}{15\pi} \cdot \frac{l^2(l+1)^2}{(l+3)(l-2)} \quad (19.350)$$

(see Refs [110]). The singularity for $l = 2$ is not realistic, there is only an amplification for $l = 2$.

Let us notice that our power flat spectrum (scale-invariant) is a good starting point for every structure formation theory in the Universe. Using our full $P_R^s(K)$ spectras we can do more precise calculations for C_l^s . Especially in order to compare with future very precise cosmological observations (Planck satellite) we should take under consideration $\frac{dn_s}{d\ln K}$ and deviations from scale invariant spectrum, which has been reported from WMAP data. Some further developments of our $P_R(K)$ spectras are beyond the scope of this work for they strongly depend on details of cosmological models involved (see Refs [110]). In particular we can consider C_l^s and C_l^{gr} only for $l \lesssim 100$. For $l \gtrsim 100$, $\frac{\Delta T}{T}$ is dominated by acoustic oscillation (see Refs [110]). C_l^{gr} are very small for $l \geq 60$ (they decay on sub-horizon scales). Finally we write

$$C_l^s l(l+1) = \text{const.} \cong \left\langle \left(\frac{\Delta T}{T}(\theta_l) \right)^2 \right\rangle, \quad (19.351)$$

$$\theta_l = \frac{\pi}{l} \quad (19.352)$$

and

$$C_l^{\text{gr}} l(l+1) = (P_0^{\text{gr}})^2 \cdot \frac{8}{15\pi} \cdot \frac{l(l+1)}{(l+3)(l-2)}. \quad (19.353)$$

The behaviour of C_l for $l > 100$ is beyond the scope of this work (now).

The interesting point in future calculations is to consider an influence of quintessence fluctuations. Those fluctuations are of scalar nature. For they depend on gravitational waves fluctuations their influence on C_l^s is negligible.

We can consider some applications of a quintessence in astrophysics. First of all we can develop a formalism concerning a macroscopic flow of a quintessence gas (i.e., a relativistic hydrodynamics of a quintessence). For a quintessence field has an influence on an effective gravitational constant, such a flow can disturb gravitational interactions between galaxies or even stars. In this approach we should add some assumptions, e.g. that a quintessence energy density is an energy density of quintessence field in a statistical field theory. However, this is beyond a scope of this work.

Secondly, it is interesting to consider a Bose-Einstein condensation of quintessence particles. They are massive scalar particles. Under some assumptions (beyond this theory) we can find a wave function of a condensat. This wave function can be considered a field of a quintessence and afterwards enters the formula for G_{eff} . In this way Einstein-Bose condensation can influence effective gravitational interactions between galaxies and stars. This is also beyond a scope of the work.

As a typical astrophysical application we can consider a problem of N bodies with the quintessence field. In this problem N points interact gravitationally with an effective gravitational constant depending on a distance. The simplest case is of course a two bodies problem. However, this problem cannot be solved analytically. It can be considered numerically using some codes for the N bodies problem. This is also beyond a scope of our work. Moreover, some numerical solutions can be applied for galaxies movement in a cluster of galaxies.

In all of those problems we should consider quintessence fluctuations developed here as a source of quintessence gas and Bose-Einstein condensat. In the third problem we should also consider an interaction of a quintessence field with mass points.

It is interesting to consider a continuous distribution of a matter (a dust) under gravitational interaction and a quintessence. In this way we consider hydrodynamic equation coupled to a quintessence field in a Newtonian physics limit (i.e. selfgravitating system with G_{eff} depending on a quintessence).

Conclusions

In this paper we consider the Nonsymmetric Kaluza–Klein (Jordan–Thiry) Theory with spontaneous symmetry breaking and Higgs’ mechanism with a scalar field Ψ (see Introduction for a short description). We get inflation from this theory with a quintessence and several testable cosmological predictions. We find a dynamical model for a cosmological constant. In the models of the Universe we get several phase transitions of the second and of the first order. The real source of a quintessence is a scalar field Ψ . Due to its very unusual selfinteraction potential (coming from higher dimensions) it can proceed to the very unexpectable features. Of course this is not the end of the story. This is really a beginning.

Especially we should develop a supersymmetric (supergravitational) extension of our theory. In order to do this it is necessary to use a general formalism of supermanifolds [136] and super-Lie algebra theory [137,138]. We should develop Einstein-Kaufmann connections on supermanifolds and super-Yang-Mills fields based on super-Lie group. From physical point of view the formalism by P. Nath and R. Arnowitt [139] will be very helpful. On the other side we can use also a theory of symplectic structure on homogeneous spaces [140].

In that way we want to connect our approach to classical approach with supersymmetry and supergravity [141] with a geometric formulation [142] geometrizing fermion fields. Perhaps our approach is one of the phases of M-theory (see [143]) as a low energy theory.

References

- [1] Kaluza T., *Zum Unitätsproblem der Physik*, Sitzgsber. Preuss. Akad. Wiss., p. 966, (1921).
- [2] Klein O. Z., Phys. **37**, p. 895 (1926).
Klein O., *On the theory of charged fields*, in New theories in Physics. Conference organized in collaboration with International Union of Physics and the Polish Co-operation Committee, Warsaw, May 30th-June 3rd, 1938, published in Paris, p. 77, (1939).
- [3] Einstein A., *The meaning of relativity*, Appendix II, Fifth Edition, revised, p. 127 (Methuen and Co.: London, U.K.), (1951).
Jakubowicz A. and Klekowska J., *The necessary and sufficient condition for the existence of the unique connection of the two-dimensional generalized Riemannian space*, Tensor N.S **20** p. 72, (1969).
Chung K. T. and Lee Y. J., *Curvature Tensors and Unified Field Equations in SEX_n* , Int. Journal of Theor. Phys. **27**, p. 1083 (1988).
Chung K. T. and Hwang H. J., *Three and Five-Dimensional Considerations of the Geometry of Einstein's *g -Unified Field Theory*, Int. Journal of theor. Phys. **27**, p. 1105, (1988).
Shavokhina N. S., *Nonsymmetric Metric in Nonlinear Field Theory*, Preprint of the JINR, P2-86-685, Dubna 1986.
- [4] Kaufmann B., *Mathematical structure of the nonsymmetric field theory*, Helv. Phys. Acta. Suppl. (1956), p. 227;
Chung K. T., *Some Recurrence Relations for Einstein's Connection for 2-Dimensional Unified Theory of Relativity*, Acta Mathematica Hungarica **41** (1-2) p. 47 (1983).
- [5] Einstein A. and Kaufmann B., Ann. of Math., **59** (1954), p. 230;
Kaufmann B., Ann. of Math., **46** (1945), p. 578.
- [6] Einstein A., *A generalization of the relativistic theory of gravitation*, Ann. of Math., **46**, p. 578, (1945).
Einstein A. and Strauss E. G., *A generalization of the relativistic theory of gravitation II*, Ann. of Math., **47**, p. 731, (1946).
- [7] Kerner R., *Generalization of Kaluza-Klein theory for an arbitrary nonabelian gauge group*, Ann. Inst. H. Poincaré IX, p. 143, (1968).
- [8] Cho Y. M., *Higher dimensional unification of gravitation and gauge theories*, Journal of Math. Phys., **16**, p. 2029, (1975).
Cho Y. M. and Freund P. G. O., *Nonabelian gauge fields as Nambu-Goldstone fields*, Phys. Rev. **D12**, p. 1711, (1975).
- [9] Kopczyński W., *A fibre bundle description of coupled gravitational and gauge fields*, in Differential Geometrical Methods in Mathematical Physics (Springer, Verlag, Berlin, Heidelberg, New York), p. 462, (1980).
- [10] Kalinowski M. W., *Vanishing of the cosmological constant in nonabelian Klein-Kaluza theories*, Int. Journal of Theor. Phys., **22**, p. 385, (1983).
- [11] Thirring W., *Five dimensional theories and CP violation*, Acta Phys. Austriaca Suppl. IX (Wien, New York, Springer), p. 256, (1972).
- [12] Kalinowski M. W., *PC nonconservation and dipole electric moment of fermion in the Kaluza-Klein Theory*, Acta Phys. Austriaca **53**, p. 229, (1981).
- [13] Kalinowski M. W., *Spinor fields in nonabelian Klein-Kaluza theories*, Int. Journal of Theor. Physics, **23**, p. 131, (1984).

- [14] Kalinowski M. W., *3/2 spinor field in the Klein–Kaluza theory*, Acta Phys. Austriaca **55**, p. 197, (1983).
- [15] Kalinowski M. W., *Rarita–Schwinger in nonabelian Klein–Kaluza theories*, Journal of Phys. **A** (Math and Gen), **15**, p. 2441. 144, (1982).
- [16] Kalinowski M. W., *Gauge fields with torsion*, Int. Journal of Theor. Physics, **20**, p. 563, (1981).
- [17] Einstein A., *Zur Elektrodynamik bewegter Körper*, Ann. Phys. (Germany), **17**, p. 891, (1905).
- [18] Kalinowski M. W., *The nonsymmetric Kaluza–Klein Theory*. Journal of Math. Phys., **24**, p. 1835, (1983).
- Kalinowski M. W., *Nonsymmetric Fields theory and its Applications*, World Scientific, Singapore, New Jersey, London, Hong Kong, 1990.
- Kalinowski M. W., *Can we get Confinement from extra dimensions*, in Physics of Elementary Interactions World Scientific, Singapore, New Jersey, London, Hong Kong 1991 (ed. Z. Ajduk, S. Pokorski, A. K. Wróblewski).
- Kalinowski M. W., *Nonsymmetric Kaluza–Klein (Jordan–Thiry) in a general Nonabelian Case*, Int. Journal of Theor. Phys. **30** p. 281 (1991).
- Kalinowski M. W., *Nonsymmetric Kaluza–Klein (Jordan–Thiry) Theory in the Electromagnetic Case*, Int. Journal of Theor. Phys. **31** p. 611 (1992).
- Kalinowski M. W., *Geodetic equations in nonsymmetrics Jordan–Thiry theory*, I and II, Bull. Soc. Sci. Lettres Łódź 45 Ser. Rech. Deform. **19** p. 89 (1995), **19**, p. 117 (1995).
- Kalinowski M. W., *Deformation of Riemannian Geometry and Kaluza–Klein Theory*, Bull. Soc. Sci. Lettres Łódź **XXV** p. 113 (1998).
- [19] Kalinowski M. W., *On the nonsymmetric Jordan–Thiry theory*, Canadian Journal of Physics, **61**, p. 844, (1983).
- [20] Jordan P., *Schwerkraft und Weltal*, (Vieweg Verlag: Braunschweig, 1955).
- [21] Thiry Y., *Étude mathématique de equations d’une theorie unitaire a quinze variables de champ*, (thèse), (Gautiers-Villars, 1951).
- [22] Lichnerowicz A., *Theorie relativistes de la gravitation et de l’electromagnetisme*, (Masson: Paris), (1955).
- [23] Kalinowski M. W., *The nonsymmetric-nonabelian Kaluza–Klein Theory*, Journal of Physics A (Math. and Gen.), **16**, p. 1669, (1983).
- [24] Kalinowski M. W., *The nonsymmetric-nonabelian Jordan–Thiry Theory*, II Nuovo Cimento LXXXA, p. 47, (1984).
- [25] Kalinowski M. W., *Material sources in the nonsymmetric Kaluza–Klein theory*, Journal of Math. Phys., **25**, p. 1045, (1984).
- [26] Kalinowski M. W., *Spontaneous symmetry breaking and Higgs’ mechanism in the nonsymmetric Kaluza–Klein theory*, Ann. of Phys., **148**, (New York), p. 241, (1983).
- [27] Kalinowski M. W., *Spontaneous symmetry breaking and Higgs’ mechanism in the nonsymmetric Jordan–Thiry theory*, Fortschritte der Physik **34**, p. 361, (1986).
- [28] Kalinowski M. W. and Mann R. B., *Linear approximation in the nonsymmetric Kaluza–Klein theory*, Clasical and Quantum Gravity, **1**, p. 157, (1984).
- [29] Kalinowski M. W. and Mann R. B., *Linear approximation in the nonsymmetric Jordan–Thiry theory*, Nuovo Cimento **91B**, p. 67, (1986).
- [30] Kalinowski M. W. and Kunstatter G., *Spherically Symmetric solution in the nonsymmetric Kaluza–Klein theory*, Journal of Math. Phys., **25**, p. 117, (1984).
- [31] Mann R. B., *Exact solution of an algebraically extended Kaluza–Klein theory*, Journal of Math. Phys., **26**, p. 2308, (1985).
- [32] Kalinowski M. W., *PC nonconservation and a dipole electric moment of fermion in the nonsymmetric Kaluza–Klein theory*, Int. Journal of Theor. Physics, **26**, p. 21, (1987).
- [33] Kalinowski M. W., *Spinor field in the nonsymmetric-nonabelian Kaluza–Klein theory*, Int. Journal of Theor. Physics, **26**, p. 565, (1987).

- [34] Moffat J. W., *Generalized theory of gravitation and its physical consequences*, in Proceeding of the VII international School of Gravitation and Cosmology. Erice, Scilly ed. by V. de Sabbata (World Scientific Publishing Co.: Singapore), p. 127, (1982).
- [35] Kunstatter G., Moffat J. W. and Malzan J., *Geometrical interpretation of a generalized theory of gravitation*, Journal of Math. Phys., **24**, p. 886, (1983).
- [36] Hilbert D., Gott. Nachr., **12**, (1916).
- [37] Levi-Civita, Atti. R. Accad. Naz. Lincei Mam. Cl. Sci. Fiz. Mat. Nat. **26**, p. 311, (1917); Thiry Y., J. Math. Pure Appl., **30**, p. 275, (1951).
- [38] Lichnerowicz A., *Sur certains problèmes globaux relatifs au système des équations d'Einstein*, (Herman: Paris), 1939.
- [39] Einstein A. and Pauli W., Ann. Math., **44**, p. 131, (1943).
Einstein A., Rev. Univ. Nac. Tucuman, **2**, p. 11, (1941).
- [40] Werder R., *Absence of stationary solutions to Einstein–Yang–Mill’s equations*, Phys. Rev., **D25**, p. 2515, (1982).
Bartnik R. and McKinnon J., *Particlelike Solutions of the Einstein–Yang–Mill’s Equations*, Phys. Rev. Lett., **61**, p. 141, (1988).
- [41] Kunstatter G., *Finite Energy Electric Monopoles in an Extended theory of gravitations*, Journal of Math. Phys., **25**, p. 2691, (1984).
- [42] Roseveare N. T., *Mercury’s Perihelion: From Le Verier to Einstein*, Claredon Press Oxford, (1982).
- [43] Hlavaty V., *Geometry of Einstein’s Unified Field Theory*, Nordhoff-Verlag, Groningen 1957.
Tonnelat M. A., *Einstein’s Unified Field Theory*, Gordon and Breach Science Publishers, New York, London, Paris (1966).
- [44] Hill H. A., Bos R. J. and Goode P. R., *Preliminary determination of the quadrupole moment of the sun from Rotational splitting of global oscillations and its relevance to test of general relativity*, Physical Review Lett. **33**, p. 709, (1983).
Hill H. A., *Rotational splitting of Global Solar Oscillations and its Relevance to test of General Relativity*, Int. Journal of Theor. Physics **23**, p. 689, (1984).
Gough D. O., *Inertial rotation and gravitational quadrupole moment of the Sun*, Nature **298**, p. 334, (1982).
- [45] Moffat J. W., *Consequences of a New experimental Determination of the Quadrupole Moment of the Sun for Gravitational Theory*, Physical Rev. Lett. **50**, p. 709, (1983).
Campbell L. and Moffat J. B., *Quadrupole moment of the Sun and Planetary Orbits*, Astrophysical Journal **275**, p. L77, (1983).
- [46] Moffat J. W., *The orbit of Icarus as a test of a Theory of gravitation*, University of Toronto preprint, May (1982).
Campbell L., McDow J. C., Moffat J. W. and Vincent D., *The sun’s quadrupole moment and the perihelion precession of Mercury*, Nature **305**, p. 508, (1983).
- [47] Moffat J. W., *Generalized theory of gravitation*, Foundation of Physics **14**, p. 1217, (1984).
Moffat J. W., *Test of a theory of gravitation using the Data from the Binary Pulsar 1913+16*, University of Toronto Report, August 1981.
Krisher T. P., *Dipole gravitational radiation in the non-symmetric theory of Moffat*, Phys. Rev. **D32**, p. 329, (1985).
Will M. C., *Violation of weak equivalence principle in theories of gravity with nonsymmetric metric*, Phys. Rev. Lett. **62**, p. 369, (1989).
- [48] Moffat J. W., *Experimental Consequences of the Nonsymmetric Gravitation theory including the Apsidal motion of Binaries*, Lecture given at the conference on General Relativity and Relativistic Astrophysics, University of Dalhousie, Halifax, Nova Scotia, April 1985.
- [49] McDow J. C., *Testing the nonsymmetric theory of gravitation*, PhD thesis, University of Toronto, 1983.
Hoffman J. A., Masshal H. L. and Lewin W. G. H., Nature **271**, p. 630, (1978).

- [50] Bergman P. G., *Comments on the scalar-tensor theory*, Int. Journal of Theor. Physics, **1**, p. 52, (1968).
- [51] Trautman A., *Fibre bundles associated with space-time*, Rep. of Math. Phys. **1**, p. 29, (1970).
- [52] Utiyama R., *Invariant Theoretical Interpretation of Interaction*, Phys. Rev. **101**, p. 1597, (1956).
- [53] Stacey F. D., Tuck G. J., Moore G. J., Holding S. C., Goldwin B. D. and Zhou R., *Geophysics and the law of gravity*, Rev. of Modern Physics **59**, p. 157, (1987).
 Ander M. E., Goldman T., Hughs R. J. and Nieto M. M., *Possible Resolution of the Brookhaven and Washington Eötvös Experiments*, Phys. Rev. Lett. **60**, p. 1225, (1988).
 Eckhardt D. H., Jekeli C., Lazarewicz A. R., Romaides A. J. and Sands R. W., *Tower Gravity Experiments: Evidence for Non-Newtonian Gravity*, Phys. Rev. Lett. **60**, p. 2567, (1988).
 Moore G. I., Stacey F. D., Tuck G. J., Goodwin B. D., Linthorne N. P., Barton M. A., Reid D. M. and Agnew G. D., *Determination of the gravitational constant at an effective mass separation of 22 m*, Phys. Rev. **D38** p. 1023, (1988).
- [54] Fischbach E., Sudarsky D., Szafer A., Tolmadge C. and Arnson S. H., *Reanalysis of the Eötvös experiment*, Phys. Rev. Lett. **56**, p. 3, (1985).
- [55] Thieberg P., *Search for a Substance-Dependent Force with a New Differential Accelerometer*, Phys. Rev. Letters **58**, p. 1066, (1987).
- [56] Wesson P. S., *Does gravity change with a time?*, Physics Today **33**, p. 32, (1980).
- [57] Gillies G. T. and Ritter R. C., *Experiments on Variation of the Gravitational Constant using Precision Rotations*, in Precision Measurements and Fundamental Constants II ed. by B. N. Taylor and W. D. Phillips, Natl. Bur. Stand. (US) Spec. Publ. 617, p. 629. (1984).
- [58] Rayski J., *Unified theory and modern physics*, Acta Physica Polonica Vol. XXVIII, p. 89, (1965).
- [59] Kobayashi S. and Nomizu K., *Foundation of differential geometry*, Vol. I and Vol. II, New York, (1963).
 Kobayashi S., *Transformation Groups in differential geometry*, Springer Verlag, Berlin (1972).
- [60] Lichnerowicz A., *Théorie globale des connexions et de group d'holonomie*, Ed. Cremonese, Rome, (1955).
- [61] Hermann R., *Yang-Mills, Kaluza-Klein and the Einstein Program*, Math. Sci. Press, Brookline, Mass. 1978.
 Coquereaux R. and Jadczyk A., *Riemannian geometry, Fibre bundle, Kaluza-Klein theory and all that ...*, World Scientific Publishing, Singapore 1988.
- [62] Zalewski K., *Lecture on rotational group*, PWN, Warsaw 1987 (in Polish).
 Barut A. O., Rączka R., *Theory of group representations and applications*, PWN, Warsaw, (1980).
- [63] Moffat J. W., *Static spherically symmetric solution for the field of a charged particle in a new theory of gravity*, Phys. Rev. **D19**, p. 3562, (1978).
- [64] Kalinowski M. W., *Comment on the nonsymmetric Kaluza-Klein theory with material sources*, Zeitschrift für Physik C (Particles and Fields) **33**, p. 76, (1986).
- [65] Moffat J. W., *New theory of gravitation*, Phys. Rev. D **19**, p. 3557, (1979).
- [66] Moffat J. W., *Gauge invariance and string interactions in a generalized theory of gravitation*, Phys. Rev. D **23**, p. 2870, (1981).
- [67] Moffat J. W. and Woolgar E., *The Apsidal Motion of the Binary Star in the Nonsymmetric Gravitational Theory*, University of Toronto Report, 1984.
 Moffat J. W., *The Orbital motion of DI Hercules as a test of the Theory of Gravitation*, University of Toronto Report, 1984.
- [68] de Groot S. R. and Suttorp R. G., *Foundations of Electrodynamics*, North Holland Publishing Company, Amsterdam, 1972.
- [69] Plebański J., *Nonlinear Electrodynamics*, Nordita Copenhagen, 1970.

- [70] Kalinowski M. W., *On a generalization of the Einstein–Cartan theory and Kaluza–Klein theory*, Lett in Math. Phys. **5**, p. 489, (1981).
Kalinowski M. W., *Torsion and the Kaluza–Klein theory*, Acta Phys. Austriaca **27**, p. 45, 1958.
- [71] Wang H. C., *On Invariant connection on principal fibre bundle*, Nagoye Math. Journal **13**, p. 19, (1958).
- [72] Harnad J., Shnider S. and Vinet L., *Group actions on Principal Bundles and Invariance conditions for Gauge Fields*, Journal of Math. Phys., **21**, p. 2719, (1980).
- [73] Harnad J., Shnider S. and Tafel J., *Group actions on Principal Bundles and Dimensional Reduction*, Lett. in Math. Phys. **4**, p. 107, (1980).
- [74] Forgacs P. and Manton N. S., *Space-time Symmetries in gauge Theories*, Comm. Math. Phys. **72**, p. 15, (1980).
- [75] Manton N. S., *A New Six-Dimensional Approach to the Weinberg–Salam Model*, Nucl. Phys. B **158**, p. 141, (1979).
Pawłowski M. and Rączka R., *A Higgs’ free model for Fundamental Interactions, Part I: Formulation of the Model*, Preprint SINS – 1P/VIII/1995 ILAS/EP–3–1995.
Pawłowski M. and Rączka R., *A Higgs’ free model for Fundamental Interactions, Part II: Predictions for Electroweak observables*, Preprint SINS – 2P/VIII/1995 ILAS/EP–4–1995.
- [76] Trautman A., *Geometrical Aspect of Gauge Configurations*, in new Developments in Mathematical Physics, Acta Phys. Austriaca Suppl. XXIII, p. 401, (1981), Springer Verlag (1981).
- [77] Witten E., *Some Exact Multipseudo-Particle Solutions of Clasical Yang–Mill’s Theory*, Phys. Rev. Lett. **38**, p. 121, (1977).
Hooft G., *Magnetic monopoles in unified gauge theones*, Nuclear Phys. B: 79 p. 276 (1976).
- [78] Maclenburg W., *Geometrical unification of gauge and Higgs’ fields*, Triesre 1979 preprint JCTP JC/79/131.
- [79] Maclenburg W., *Towards a unifold theory for gauge and Higgs’ fields*, Trieste 1981 preprint JTCP JC/81/8.
- [80] Chapline G. and Manton N. S., *The Geometrical Significance of Certain Higgs’ Potentials: an Approach to grand Unification*, Nucl. Phys. B **184**, p. 391, (1981).
- [81] Sato K., *Cosmological Consequences on a First Order Phase Transition of a Vacuum*, Proceedings of the 4th Kyoto summer Institute ed. M. Konuma and T. Maskawe, World Scientific Publishing Co., Singapore (1981).
- [82] Linde A., *Coleman–Weinberg theory and inflationary universe scenerio*, Phys. Lett. **114B**, p. 431, (1982).
- [83] Linde A. D., *A new Inflationary Universe Scenario: a Possible Solution of the Horizon, Flatness, Homogeneity, Isotropy and Primordial Monopoles Problems*, Phys. Lett. **1088**, p. 389, (1982).
- [84] Sher M. A., *Effects of phase transitions of the evolution of the early universe*. Phys. Rev. D **22**, p. 2989, (1980).
- [85] Guth A. H., *Inflantionary universe: A possible solution to the horizon and flatness problems*, Phys. Rev. D **23**, p. 347, (1981).
- [86] Helgason S., *Differential Geometry, Lie groups and Symmetric Spaces*, Academic Press, New York 1987.
Helgason S., *Groups and Geometric Analysis*, Academic Press Inc., London, 1984.
Wolf J. A. and Gray A., *Homogeneous spaces defined by Lie group automorphisms I, II* Journal of Differential Geometry **2** p. 77, (1968), **2** p. 115, (1968).
Gray A., *Nearly Kählerian manifolds*, Journal of Differential Geometry, **4**, p. 283, (1970).
Штукарь В. Л., *Инвариантные связности и метрики на однородных пространствах, соответствующих глобальным тройкам*, Математические заметки Т. 26, стр. 449 (1979).

- [87] Cheeger J. and Ebin D. G., *Comparison Theorem in Riemannian Geometry*, North Holland/American Elsevier, New York, Amsterdam, Oxford (1975).
- [88] Coleman S. and Weinberg E., *Radiative Corrections as the Origin of the Spontaneous Symmetry Breaking*, Phys. Rev. D **7**, p. 1888, (1973).
- [89] Weinberg S., *Gauge hierarchies*, Phys. Lett. **82B** p. 387.
- [90] Moffat J. W., *Phase Transition in the Early Universe as a consequence of a Generalized Theory of Gravitation*, invited talk presented at the Conference of Differential Geometric Methods in Theoretical Physics, Trieste, Italy June 30-July 3, (1981).
- [91] Moffat J. W. and Vincent D., *The Early Universe in a Generalised Theory of Gravitation*, Can. Journal of Phys. **60**, p. 659, (1982).
- [92] Kalinowski M. W., *The minimal coupling scheme for Dirac's field in the nonsymmetric theory of gravitation*, Journal of Modern Physics **A1**, p. 185, (1981).
- [93] Langacker P., *Grand Unified Theories and Proton Decay*, Phys. Rep. **72**, p. 185, (1981).
- [94] Mann R. B. and Moffat J. W., *Post-Newtonian approximation of new theory of gravity*, Can J. Phys. **59**, p. 1592, (1981).
- [95] Moffat J. W., *Test Particle Motion in the Nonsymmetric Gravitation Theory*, Phys. Rev. **D35** p. 3733, (1987).
- [96] Moffat J. W., Savaria R. and Woolgar E., *Eötvös Test of the Nonsymmetric Gravitation Theory and the Fifth Force*, University of Toronto Report, 1986.
Moffat J. W. and Woolgar E., *Motion of masive bodies: Testing the nonsymmetric gravitation theory*, Phys. Rev. **D37**, p. 918, (1988).
Krisher T. P., Astrophys. J. (Lett) **320** p. 147, (1987).
Moffat J. W., *Cosmions in the Nonsymmetric Gravitational Theory*, Phys. Rev. **D39**, p. 474, (1989).
Mann R. B., Moffat J. W. and Palmer J. H., *Comment on "Violation of the Weak Equivalence Principle in Theories of Gravity with a Nonsymmetric Metric"*. MATPHYS-TH 89/01, U of T Report, Toronto 1989.
Krisher T. P., *Possible test at Jupiter of the nonsymmetric gravitational theory*, Phys. Rev. **D40** p. 1373, (1989).
- [97] Einstein A., *The Bianchi identities in the generalized theory of gravitation*, Can. J. Math. **2**, p. 120, (1950).
- [98] Kunstatter G. and Moffat J. W., *Conservation laws in a generalized theory of gravitation*, Phys. Rev. **D19**, p. 1084, (1979).
- [99] Rudjobing M., Ann Astrophys. **22**, p. 111, (1959).
Martynov D. Ya. and Khaliulin Kh. F., Astrophysics Space Sc. **71**, p. 147, (1980).
Popper D. M., Ap. J. **188**, p. 559, (1974).
Popper D. M., Ap. J. **254**, p. 209, (1982).
Guinan E. F. and Maloney F. P., 163-rd AAS Meeting, Las Vegas (Abstract);
Khaliulin Kh. F., Sov. Astron. **27**, p. 43, (1983).
Koch R. H., Ap. J. **183**, p. 275, (1973).
Koch R. H., Astron. J. **82**, p. 653, (1977).
- [100] Will C. M., *Theory and Experiment in Gravitational Physics*, Cambridge University Press, 1981.
- [101] Cornish N. J., Moffat J. M., *Non-singular gravity without black holes*, Journal of Math. Phys. **35**, p. 6628 (1994).
- [102] Cornish N. J., Moffat J. M., *A non-singular theory of gravity ?*, Phys. Lett. **B 336** p. 337.
- [103] Cornish N. J., *Analysis of the Nonsingular Wyman-Schwarzschild Metric*, Modern Physics Letters **A9**, p. 3629 (1994).
- [104] Bahcall N. A., Ostriker J. P., Perlmutter S., Steinhardt P. J., *The cosmic triangle: revealing the state of the Universe*, Science **284**, p. 1481 (1999).

- Wang L., Caldwell R. R., Ostriker J. P., Steinhardt P. J., *Cosmic concordance and quintessence*, arXiv: astro-ph/9901388v2.
- Wiltshire D. L., *Supernovae Ia, evolution and quintessence*, arXiv: astro-ph/0010443.
- Primack J. R., *Cosmological parameters*, Nucl. Phys. B (Proc. Suppl.) **87**, p. 3 (2000).
- Bennett C. L. et al., *First Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Determination of Cosmological Parameters*, arXiv: astro-ph/0302209v2.
- [105] Caldwell R. R., Dave R., Steinhardt P. J., *Cosmological imprint of an energy component with general equation of state*, Phys. Rev. Lett. **80**, p. 1582 (1998).
- Armeendariz-Picon C., Mukhanov V., Steinhardt P. J., *Dynamical solution to the problem of a small cosmological constant and late-time cosmic acceleration*, Phys. Rev. Lett. **85**, p. 4438 (2000).
- Maor J., Brustein R., Steinhardt P. J., *Limitations in using luminosity distance to determine the equation of state of the Universe*, Phys. Rev. Lett. **86**, p. 6 (2001).
- [106] Axenides M., Floratos E. G., Perivolaropoulos L., *Some dynamical effects of the cosmological constant*, Modern Physics Letters **A15**, p. 1541 (2000).
- [107] Vishwakarma R. G., *Consequences on variable Λ -models from distant type Ia supernovae and compact radio sources*, Class Quantum Grav. **18**, p. 1159 (2001).
- [108] Díaz-Rivera L. M., Pimentel L. O., *Cosmological models with dynamical Λ in scalar-tensor theories*, Phys. Rev. **D66**, p. 123501-1 (1999).
- González-Díaz P. F., *Cosmological models from quintessence*, Phys. Rev. **D62**, p. 023513-1 (2000).
- Frampton P. H., *Quintessence model and cosmic microwave background*, astro-ph/0008412.
- Dimopoulos K., *Towards a model of quintessential inflation*, Nucl. Phys. B (Proc. Suppl.) **95**, p. 70 (2001).
- Charters T. C., Mimoso J. P., Nunes A., *Slow-roll inflation without fine-tuning*, Phys. Lett. **B472**, p. 21 (2000).
- Brax Ph., Martin J., *Quintessence and supergravity*, Phys. Lett. **B468**, p. 40 (1999).
- Sahni V., *The cosmological constant problem and quintessence*, Class. Quantum Grav. **19**, p. 3435 (2002).
- [109] Liddle A. R., *The early Universe*, arXiv: astro-ph/9612093.
- Gong J. O., Stewart E. D., *The power spectrum for multicomponent inflation to second-order corrections in the slow-roll expansion*, Phys. Lett. **B538**, p. 213 (2002).
- Sasaki M., Stewart E. D., *A general analytic formula for the spectral index of the density perturbations produced during inflation*, Progress of Theoretical Physics **95**, p. 71 (1996).
- Nakamura T. T., Stewart E. D., *The spectrum of cosmological perturbations produced by a multicomponent inflation to second order in the slow-roll approximation*, Phys. Lett. **B381**, p. 413 (1996).
- Turok N., *A critical review of inflation*, Class. Quantum Grav. **19**, p. 3449 (2002).
- [110] Liddle A. R., Lyth D. H., *Cosmological Inflation and Large-Scale Structure*, Cambridge Univ. Press, Cambridge 2000.
- Peebles P. J. E., *Principles of Physical Cosmology*, Princeton Univ. Press, Princeton, 1993.
- Tsagas Ch. G., *Cosmological Perturbations*, arXiv: astro-ph/0201405v1.
- Durrer R., *The Theory of CMB Anisotropies*, arXiv: astro-ph/0109522v1.
- Bertschinger E., *Cosmological Perturbation Theory and Structure Formation*, arXiv: astro-ph/0101009v1.
- Brandenberger R. H., *Lectures on the Theory of Cosmological Perturbations*, arXiv: hep-th/0306071v1.
- Challinor A., *Anisotropies in the Cosmic Microwave Background*, arXiv: astro-ph/0403344v1.
- [111] Steinhardt P. J., Wang L., Zlatev I., *Cosmological tracking solutions*, Phys. Rev. **D59**, p. 123504-1 (1999).

- Steinhardt P. J., *Quintessential Cosmology and Cosmic Acceleration*, Web page <http://feynman.princeton.edu/~steinh>
- [112] Painlevé P., *Sur les équations différentielles du second ordre et d'ordre supérieur dont l'intégrale générale est uniforme*, Acta Mathematica **25**, p. 1 (1902).
- [113] Steigman G., *Primordial alchemy: from the big-bang to the present Universe*, arXiv: astro-ph/0208186.
Steigman G., *The baryon density through the (cosmological) ages*.
Scott J., Bechtold J., Dobrzycki A., Kul-Karni V. P., *A uniform analysis of the LY- α forest of $Z = 0 - 5$* , arXiv: astro-ph/0004155.
Hui L., Haiman Z., Zaldarriaga M., Alexander T., *The cosmic baryon fraction and the extragalactic ionizing background*, arXiv: astro-ph/0104442.
Balbi A., Ade P., Bock J., Borrill J., Boscaleri A., De Bernardis P., *Constraints on cosmological parameters from Maxima-1*, arXiv: astro-ph/0005124.
Jeff A. H. et al., *Cosmology from Maxima-1, BOOMERANG & COBE/DMR CMB observations*, arXiv: astro-ph/0007333.
- [114] Serna A., Alimi J. M., *Scalar-tensor cosmological models*, arXiv: astro-ph/9510139.
Serna A., Alimi J. M., *Constraints on the scalar-tensor theories of gravitation from primordial nucleosynthesis*, arXiv: astro-ph/9510140.
Navarro A., Serna A., Alimi J. M., *Search for scalar-tensor gravity theories with a non-monotonic time evolution of the speed-up factor*, Class. Quantum Grav. **19**, p. 4361 (2002).
Serna A., Alimi J. M., Navarro A., *Convergence of scalar-tensor theories toward General Relativity and primordial nucleosynthesis*, arXiv: gr-qc/0201049.
- [115] Kalinowski M. W., *Dynamics of Higgs' field and a quintessence in the nonsymmetric Kaluza-Klein (Jordan-Thiry) theory*, arXiv: hep-th/0306241.
Kalinowski M. W., *Cosmological models in the nonsymmetric Kaluza-Klein theory*, arXiv: astro-ph/0310745v3.
- [116] Kälbermann G., *Communication through an extra dimension*, Int. J. of Modern Phys. **A15**, p. 3197 (2000).
Visser M., *An exotic class of Kaluza-Klein model*, Phys. Lett. **B159**, p. 22 (1985).
Li Li-Xin, Gott III J. R., *Inflation in Kaluza-Klein theory: Relation between the fine-structure constant and the cosmological constant*, Phys. Rev. **D58**, p. 103513-1 (1998).
Squires E. J., *Dimensional reduction caused by a cosmological constant*, Phys. Lett. **B167**, p. 286 (1986).
Caldwell R., Langlois D., *Shortcuts in the fifth dimension*, Phys. Lett. **B511**, p. 129 (2001).
Loup F., Santos P. A., Martins da Silva Santos D., *Can Geodesics in Extra Dimensions Solve Cosmic Light Speed Limit?*, Gen. Rel. and Grav. **35**, p. 1849 (2003).
Loup F., Santos P. A., Martins da Silva Santos D., *The Dynamics of Chung-Freese Braneworld*, Gen. Rel. and Grav. **35**, p. 2035 (2003).
- [117] Dick R., *Brane worlds*, Class. Quantum Grav. **18**, p. R1 (2001).
Brax Ph., Falkowski A., Lalak Z., *Non-BPS branes of supersymmetric brane worlds*, Phys. Lett. **B521**, p. 105 (2001).
- [118] Randall L., Sundrum R., *An alternative to compactification*, arXiv: hep-th/9906064.
- [119] Arkani-Hamed N., Dimopoulos S., Dvali G., *The hierarchy problem and new dimensions at a milimeter*, Phys. Lett. **B429**, p. 263 (1998).
- [120] Kalinowski M. W., *A Warp Factor in the nonsymmetric Kaluza-Klein (Jordan-Thiry) theory*, arXiv: hep-th/0306157v3.
Kalinowski M. W., *Hierarchy of symmetry breaking in the nonsymmetric Kaluza-Klein (Jordan-Thiry) theory*, arXiv: hep-th/0307135v2.
Kalinowski M. W., *Higgs' Fields in the Nonsymmetric Field Theory*, Warsaw 1995.
- [121] Damour T., Deser S. and MacCarthy J., *Nonsymmetric gravity theories: Inconsistencies and a cure*, Phys. Rev. **D47** p. 1541 (1993).

- [122] Misner C. W., Thorne K. S. and Wheeler J. A., *Gravitation*, San Francisco: Freeman (1973).
Hulse R. A. and Taylor J. H., *Astrophys. J.* **195** p. L51 (1975).
- [123] Moffat J. W., *Nonsymmetric gravitational theory*, *J. Math. Phys.* **36**, p. 3722 (1995).
- [124] Cones A., *Noncommutative Geometry*, Academic Press Inc., London UK, San Diego California USA, 1994.
Martin C. P., Garcia-Bondia J. M. and Varilly J. C., *The Standard Model as a noncommutative geometry: the low-energy regime*, *Physics Reports* **294**, p. 363 (1998).
Jochun B. and Schücker T., *Yang–Mills–Higgs versus Connes–Lott*, *Comm. in Math. Physics* **178** p. 1 (1996).
Jochun B. and Schücker T., *A Left-Right Symmetric Model à la Connes–Lott*, *Lett. in Math. Physic* **32**, p. 153 (1994).
Coquereaux R., Häubling R., Papadopoulos N. A. and Scheck F., *Generalized gauge transformations and hidden symmetry in the standard model*, *Int. Journal of Modern Phys.* **7**, p. 2809 (1992).
Dubois-Violette M., Kerner R. and Madore J., *Noncommutative differential geometry of matrix algebras*, *J. Math. Phys.* **31**, p. 316 (1990).
Madore J., *An Introduction to Noncommutative Differential Geometry and Its Physical Application*, Cambridge University Press, Cambridge 1995.
- [125] Ashtekar A., *New variables for classical and quantum gravity*, *Phys. Rev. Lett.* **57**, p. 2244 (1986).
Ashtekar A., *New Hamiltonian formulation of general relativity*, *Phys. Rev.* **D36**, p. 1587 (1987).
Ashtekar A., Lewandowski J., Marolf D., Mourão J. and J., Thiemann T., *Quantization of diffeomorphism invariant theories of connections with local degrees of freedom*, *J. Math. Phys.* **36**, p. 6456 (1995).
Ashtekar A., Lewandowski J. and Sahlmann H., *Polymer and Fork representations for scalar field*, *Class. Quantum Gravity* **20**, L1–11 (2003).
Rovelli C., *Notes for a brief history of quantum gravity*, arXiv: gr-qc/0006061.
Gaul M., Rovelli C., *Loop quantum gravity and the meaning of diffeomorphism invariance*, *Lect. Notes Phys.* **541**, p. 277 (2000).
Thiemann T., *Lectures on Loop Quantum Gravity*, arXiv: gr-qc/0210094v1.
Thiemann T., *Quantum Spin Dynamics (QSD) : VII. Symplectic Structures and Continuum Lattice Formulations of Gauge Field Theories*, arXiv: hep-th/0005232v1, *Class. Quant. Grav.* **18**, p. 3293 (2001).
Ashtekar A., *A Generalized Wick Transform for Gravity*, arXiv: gr-gc/9511083v2, *Phys. Rev.* **D53**, p. 2865 (1996).
Ashtekar A., Lewandowski J., *Background Independent Quantum Gravity: A Status Report*, arXiv: gr-gc/0404018v1.
- [126] Donoghue J. F., *The Quantum Theory of General Relativity at Low Energies*, arXiv: gr-gc/9607039v1.
Donoghue J. F., *General relativity as an effective field theory: The leading quantum corrections*, *Phys. Rev.* **D50**, p. 3874 (1994).
Donoghue J. F., *Introduction to the Effective Field Theory. Description of Gravity*, arXiv: gr-gc/9512024v1.
Donoghue J. F., Torma T., *On the power counting of loop diagrams in general relativity*, arXiv: hep-th/9602121v1.
Bjerrum-Bohr N. E. J., Donoghue J. F., Holstein B. R., *Quantum corrections to the Schwarzschild and Kerr metrics*, *Phys. Rev.* **D68**, p. 084005 (2003).
Donoghue J. F., *Leading Quantum Correction to the Newtonian Potential*, *Phys. Rev. Lett.* **72**, p. 2996 (1994).

- Donoghue J. F., Torma T., *Infrared behaviour of graviton-graviton scattering* arXiv: hep-ph/9901156v1.
- Donoghue J. F., Holstein B. R., Garbrecht B., Konstandin T., *Quantum Corrections to Reissner-Nordström and Kerr-Newman metrics*, Phys. Lett. **B529**, p. 132 (2002).
- Bjerrum-Bohr N. E. J., Donoghue J. F., Holstein B. R., *Quantum gravitational corrections to the nonrelativistic scattering potential of two masses*, Phys. Rev. **D67**, p. 084033 (2003).
- [127] Colombeau J. F., *Differential Calculus and Holomorphy*, North-Holland Publishing Comp., Amsterdam 1982.
- Colombeau J. F., *New Generalized Functions and Multiplication of Distribution*, North-Holland Publishing Comp., Amsterdam 1984.
- Colombeau J. F., *Elementary Introduction to New Generalized Functions*, North-Holland Publishing Comp., Amsterdam 1985.
- Colombeau J. F., *Multiplication of Distributions*, Springer-Verlag, Berlin 1992 (Lecture Notes in Mathematics 1532).
- Charpentier E., *Sur l'élimination des "infinis" en théorie quantique des champs: la régularisation zeta à l'épreuve de l'interprétation de Colombeau ou vice versa*, Dissertationes Mathematicae CCCLXXXIII, Warszawa 1999.
- [128] www.cmbfast.org
www.cmbeasy.org
Dorau M., *CMBEASY: an Object Oriented Code for the Cosmic Microwave Background*, arXiv: astro-ph/0302138v1.
- [129] Kalinowski M. W., *On some cosmological consequences of the Nonsymmetric Kaluza-Klein (Jordan-Thiry) Theory*, arXiv: hep-th/0307289v2.
- Gott III J. R. et al., *A Map of the Universe*, arXiv: astro-ph/0310571v1.
- [130] Kalinowski M. W., *A short note on a total amount of inflation*, arXiv: hep-th/0307299v2.
- [131] Emden R., *Gaskugeln: Anwendungen der mechanischer Wärme*, B. G. Teubner, Leipzig und Berlin 1907.
- Chandrasekhar S., *Introduction to the Study of Stellar Structure*, New York, Dover 1939.
- [132] Lemke H., *Über die Differentialgleichungen, welche den Gleichgewichtszustand eines gasförmigen Himmelskörpers bestimmen, dessen Teile gegeneinander nach dem Newtonschen Gesetze gravitieren*, Journal für Mathematik, Bd. 142, Heft 2, S. 118 (1913).
- [133] Steinhardt P. J., *The quintessential introduction to dark energy*, Lecture given at *The search of dark matter and dark energy in the Universe*, The Royal Society, January 22–23, 2003.
- DeDeo S., Caldwell R. R., Steinhardt P. J., *Effects of the sound speed of quintessence on the microwave background and large scale structure*, arXiv: astro-ph/0301284v2.
- Erickson J. K., Caldwell R. R., Steinhardt P. J., Armendriz-Picon C., Mukhanov V., *Measuring the speed of sound of quintessence*, arXiv: astro-ph/0112438v1.
- Rahul D., Caldwell R. R., Steinhardt P. J., *Sensitivity of the cosmic microwave background anisotropy to initial conditions in quintessence cosmology*, arXiv: astro-ph/0206372v2.
- Caldwell R. R., Steinhardt P. J., *The imprint of gravitational waves in models dominated by a dynamical cosmic scalar field*, arXiv: astro-ph/9710062v1.
- [134] Chakraborty S., Chakraborty N. C., Debnath U., *A quintessence problem in self-interacting Brans-Dicke theory*, arXiv: gr-gc/0305103v1.
- Chakraborty S., Chakraborty N. C., Debnath U., *Quintessence problem and Brans-Dicke theory*, arXiv: gr-qc/0306040v1.
- [135] Kalinowski M. W., *Comment on a quintessence particle mass in the Kaluza-Klein Theory and properties of its field*, arXiv: astro-ph/0310300v6.
- [136] De Witt B., *Supermanifolds*, second edition, Cambridge University Press, Cambridge 1992.
- [137] Kac V. G., *A sketch of Lie superalgebra*, Comm. in Math. Phys. **53**, p. 31 (1977).
- [138] Kac V. G., *Lie superalgebras*, Advances in Math. **26**, p. 8 (1977).

- [139] Nath P., Arnowitt R., Chamseddine A. H., *Applied $N = 1$ Supergravity*, World Scientific, London, Singapore, Hongkong 1984.
- [140] Zwart Ph. B., Boothby W. M., *On compact homogeneous symplectic manifolds*, Ann. Inst. Fourier Grenoble **30**, p. 129 (1980).
Tralle A., Oprea J., *Symplectic manifolds with non-Kähler structure*, Springer-Verlag, Berlin-Heidelberg 1997.
- [141] Wess J., Bagger J., *Supersymmetry and Supergravity*, second edition revised and expanded, Princeton University Press, Princeton, New Jersey 1992.
- [142] Binétruy P., Girardi G., Grimm R., *Supergravity couplings: A geometric formulation*, Phys. Reports **343**, p. 255 (2001).
- [143] Polchinski J., *String Theory*, vol. I+II, Cambridge University Press, Cambridge, Mass. 1998.

Acknowledgement

We would like to thank University of Łódź, Poland, for a partial support by Badania Własne UŁ Nr. 505/485 on early stages of development of this work. We thank also Higher Vocational State School in Chełm for a continuous support during the academic year 2002/2003 under own scientific programme conducted in the Institute for Mathematics and Computer Science.

